1
2
3 ASSESSMENT OF QUALITY CHANGES IN FROZEN SARDINE (*Sardina*
4 *pilchardus*) BY FLUORESCENCE DETECTION
5
6
7
8
9

10
11 Santiago P. Aubourg*, Carmen G. Sotelo and Ricardo Pérez- Martín
12
13
14

15 Instituto de Investigaciones Marinas (CSIC)
16 C/ E. Cabello, 6
17 36208-VIGO (Spain)
18
19
20
21
22

23 *Correspondent:

24 Phone: +34 86 231930

25 Fax: +34 86 292762

26 E-mail: saubourg@iim.csic.es
27

ABSTRACT

The formation of fluorescent compounds was tested as a quality assessment during the frozen storage of sardine at -18°C (up to 24 months) and -10°C (up to 120 days). The fluorescence ratio between two excitation/emission maxima (393/463 nm and 327/415 nm) was studied in the aqueous (δF_{aq}) and organic (δF_{or}) extracts after Bligh and Dyer extraction of the white muscle. Fluorescence results were compared to common quality indices (total volatile base-nitrogen, TVB-N; conjugated dienes; thiobarbituric acid, TBA-i; free fatty acids, FFA). The δF_{aq} showed good correlations with the storage time ($r = 0.80$ and $r = 0.72$, at -18 and -10°C , respectively) and the TBA-i ($r = 0.92$ and $r = 0.81$). The principal component analysis grouped the δF_{aq} with quality indices that are sensitive for the assessment of fish damage during the frozen storage at both temperatures (TBA-i and FFA at -18°C ; BVT-N, TBA-i and FFA at -10°C). According to the present results, fluorescence detection of interaction compounds in the aqueous phase can provide an accurate method to assess quality differences during the frozen storage of fish.

Key Words: Fluorescence, frozen storage, interaction compounds, lipid damages, sardine.

Running Title: Fluorescence in frozen sardine

INTRODUCTION

Canned fish and other marine species are products of large economic importance in many countries. Many of the problems encountered with poor quality canned fish can be related with the quality of the raw material (1). Frozen storage is the most utilized method in canneries for preserving fish prior to canning. Fish species employed possess a high lipid content (2, 3), which in addition have a high proportion of polyunsaturated fatty acids (PUFA) typical of marine lipids (4). During the storage period enzymatic and non enzymatic lipid oxidation (5, 6) can become a very important factor responsible for fish damage. A close relationship has been found between the lipid damages and the quality of the raw material employed for canning, and as a result with the quality of the final canned product (7, 8).

Many methods have been used to measure lipid oxidation in foods as a means for determining the degree of damage (9, 10). However, some difficulties were found with common methods when quality has to be assessed because oxidation products are unstable and tend to react with biological amino constituents (proteins, peptides, free amino acids and phospholipids), leading to interaction compounds (11-14).

The analysis of the above interaction products by fluorescence detection has become a complementary method to the other more sophisticated measurements for assessing lipid damage (15-18). Recent studies have measured the fluorescent properties of thermally treated fish at different excitation/emission maxima (19-21). Increments of lipid oxidation, time and temperature of processing produced a fluorescence shift towards higher wavelength maxima. The fluorescence ratio between two of these maxima (393/463 nm and 327/415 nm) showed an interesting correlation with fish quality after cooking and during chilling (20-22).

The present work was design to test the fluorescent compounds detection as a quality assessment index for frozen stored sardine. Two frozen storage temperatures (-18°C and -10°C) were studied. The fluorescence shift, measured as the above mentioned ratio, was studied together with common quality estimations (total volatile base-nitrogen, conjugated dienes, thiobarbituric acid index, free fatty acids).

MATERIALS AND METHODS

Raw material, freezing, frozen storage and sampling

Fresh sardines (*Sardina pilchardus*) were obtained in a local market. Upon arrival in our laboratory, individual fish were frozen at -40°C and then distributed into two storage conditions: -18°C and -10°C. At each storage temperature, sardines were divided into three batches that were analyzed separately along the whole experiment. Sardines stored at -18°C were sampled at month 0.5, 2, 4, 8, 12,

18 and 24 and those stored at -10°C were sampled at day 3, 10, 25, 60 and 120. In each batch of each storage temperature, analyses were performed on the homogenized white muscle from three individual sardines.

Water and lipid contents

Water content was determined by weight difference of the homogenized muscle (1-2g) before and after 24 hours at 105°C. Lipids were extracted by the Bligh and Dyer (23) method. Quantification was carried out according to Herbes and Allen (24).

Total volatile base-nitrogen (TVB-N) determination

TVB-N was measured by the Antonacopoulos (25) method with some modifications. Fish muscle (10g) was extracted with perchloric acid (6%) and made up to 50 mL. TVB-N content was determined by steam distillation of the acid extracts made alkaline to pH 13 with NaOH (20%), followed by titration of the distillate with 10 mM hydrochloric acid. Data were expressed as mg TVB-N/ 100g muscle.

Determination of lipid damages

Conjugated dienes (CD) formation was measured at 233 nm (10). Results are expressed according to the following formula (15): $CD = B \times V / w$, where B is the absorbance reading at 233 nm, V denotes the volume (mL) of the sample and w is the mass (mg) of the lipid sample.

The thiobarbituric acid index (TBA-i) (mg malondialdehyde / kg sample) was determined according to Vyncke (26).

Free fatty acids (FFA) content was determined by the Lowry and Tinsley (27) method based on complex formation with cupric acetate-pyridine. Results are expressed as g FFA / 100g lipids.

Fluorescence analysis

A Perkin-Elmer LS 3B fluorescence spectrophotometer was employed. Fluorescence formation was studied at 393/463 nm and 327/415 nm according to previous experiences (21, 22). The relative fluorescences (RF) were calculated as: $RF = F / F_{st}$, where F is the sample fluorescence at each excitation/emission maximum, and F_{st} is the corresponding fluorescence intensity of a quinine sulfate solution (1 µg/mL in 0.05 M H₂SO₄). The fluorescence shift (δF) was calculated as the ratio between both RF values: $\delta F = RF_{393/463 \text{ nm}} / RF_{327/415 \text{ nm}}$. The δF value was studied in the aqueous (δF_{aq}) and organic (δF_{or}) phases resulting from the lipid extraction (23); the ratio between both values ($\delta F_{or} / \delta F_{aq}$) was also evaluated.

1 Statistical analysis

2 The *Statistica* package (28) was employed. Data from the different damage measurements were
3 subjected to the ANOVA one-way method ($p < 0.05$), correlation analysis and factor analysis (principal
4 components) ($p < 0.05$ and $p < 0.01$). A varimax normalized rotation was employed for factor rotation.

8 RESULTS

9
10 Water contents ranged between 68% and 71% in all samples. No significant differences ($p < 0.05$)
11 were obtained as a result of temperatures of frozen storage (-18°C and -10°C). Lipid contents ranged
12 between 4% and 7% (wet basis). Little differences observed between samples, could be explained as a
13 result of lipid content variations in individual fishes, and not as a result of the frozen storage.

14 Quality measurements

15
16 Sardines stored at -18°C produced TVB-N during the storage period at a slower rate than those
17 stored at -10°C . A slight increase with the storage time was observed at -10°C while no clear tendency
18 was detected at -18°C . The amount of volatile amines nitrogen has been widely employed for the
19 estimation of fish quality during and after several processes (29, 30). In the case of frozen stored fish,
20 volatile amines result from the breakdown of either trimethylamine oxide (in the case of gadoid fish
21 possessing) or amino acids, leading to the accumulation of dimethylamine and NH_3 , respectively (31).
22 Since non gadoid fish do not show trimethylamine oxide demethylase activity (32), it might be expected
23 that the TVB-N formed during the frozen storage of sardines comes mainly from the deamination of amino
24 acids.

25 Primary oxidation products (conjugated dienes) showed some significant differences although a
26 definite pattern during the storage time at both frozen temperatures could not be observed (Tables 1 and
27 2). It is concluded that the CD content was not suitable as a quality measurement since dienes are
28 relatively unstable and capable of interacting with other constituents (21, 22, 33).

29 Secondary lipid oxidation products were measured by the TBA-i. TBA-i determined of this index at
30 -18°C (Table 1) showed a significant increase after 4 months of storage. Thereafter, progressive
31 increases were found along the whole storage. A similar behavior was obtained at -10°C (Table 2), where
32 a significant increase was obtained after 10 days of storage. According to previous experiences (34, 35),
33 the TBA-i has shown to be an accurate way of assessing quality changes during the frozen storage time.

34 Lipid hydrolysis was measured by the FFA. A progressive increase of the FFA content was
35 observed throughout the storage at both frozen temperatures (Tables 1 and 2). At -18°C increases were
36 detected after 0.5, 12 and 24 months of storage. At -10°C differences were observed after 3 and 25 days

of storage. According to previous experiences (34, 36), the FFA formation as a result of the lipid hydrolysis (triglycerides and phospholipids classes) has provided a suitable means for assessment of fish damage during the frozen storage.

Fluorescence measurements

The fluorescence shift measured in the organic phase resulting from the Bligh and Dyer (23) extraction (δF_{or}) of the different frozen samples was studied. Both storage temperatures (Tables 1 and 2) showed a progressive increase of the δF_{or} value with time, except for the end of the storage where a decrease was detected (24 months at -18°C and 120 days at -10°C). Significant increases were obtained after 0.5 and 18 months (at -18°C) and after 10 and 60 days (at -10°C).

However, measurement of fluorescence in the aqueous phase resulting from Bligh and Dyer (23) extraction (δF_{aq} value) showed a progressive increase along the whole storage at both temperatures (Tables 1 and 2). At -18°C significant increases were detected after 12, 18 and 24 months, while at -10°C increases were observed after 60 and 120 days.

In order to study the relative formation of fluorescent compounds that are soluble in organic solvents and water, the $\delta F_{or}/\delta F_{aq}$ ratio was evaluated. A similar behavior was obtained at both frozen temperatures (Tables 1 and 2). An increase in the $\delta F_{or}/\delta F_{aq}$ ratio was observed during the first steps of the storage, that was followed by a sharp decrease after 12 months (at -18°C) and after 25 days (at -10°C).

A similar behavior was observed by studying the $\delta F_{or}/\delta F_{aq}$ ratio during sardine storage at 0°C and 10°C (22). In the above work, during the first steps of the storage, fluorescent compounds responsible for the δF value were mostly lipid-soluble; however, as the lipid damage increased, these kinds of compounds became progressively more soluble in the aqueous phase.

Fluorescent compounds studies based on a single excitation/emission maximum have been mostly carried out on organic extracts (lipids) and have shown high correlations with sensory measurements and storage times (16, 17). However, experimental evidence has demonstrated that fluorescent substances formed from oxidized membrane lipids remain attached to the amino constituents resulting in compounds quite insoluble in organic solvents (37-39). According to our results based on fluorescence ratios, the analysis of the aqueous phase can provide a more accurate assessment of the fluorescent properties of the interaction compounds formed as a result of the fish damage.

Correlation and multivariate analyses

The different damage indices were tested for correlations with storage time and also with each other (Tables 3 and 4). Almost every index correlates well with the storage time (ST) at both temperatures, except for the CD which did not vary with storage time. Indices behave similarly at both storage temperatures. Two indices (TBA-i and FFA) correlate best with storage time at both temperatures and

according to previous experiences could be used as an indication of quality deterioration of frozen sardine (34-36).

Concerning the fluorescence measurements, both indices (δF_{or} and δF_{aq}) reflected good correlations with the storage time at both temperatures and showed to be strongly correlated with those lipid damage indices that also show variation with storage time (FFA and TBA-i). Correlation data are better for the δF_{aq} than for the δF_{or} , especially in the case of -18°C storage.

Principal component analysis showed that the 80.0 % and 80.9 % of the variability observed at -18 and -10°C , respectively, could be explained by two factors. These factor loadings are graphically displayed in Figures 1 and 2. Figure 1 (-18°C) shows that ST, FFA, TBA-i and δF_{aq} are grouped together at high loading in the Factor 1 axis, whereas their loading values for Factor 2 are low; meanwhile CD, TVB-N and δF_{or} have clearly higher loadings in Factor 2. In the case of -10°C (Figure 2) similar results can be concluded, with the exception that the TVB-N index has a high loading in Factor 1 axis and is grouped together with the ST, FFA, TBA-i and δF_{aq} .

DISCUSSION

In previous experiences related to chilling, cooking and canning lipid deterioration compounds (peroxides and carbonyls) have caused the formation of interaction compounds with fluorescent properties (20-22). Accelerated by the temperature, fluorescent compounds formed in the first stages of interaction led to the formation of other fluorescent compounds showing excitation/emission maxima at higher wavelengths than their precursors. The fluorescence ratio between two of these maxima (393/463 nm and 327/415 nm) provided an interesting method for quality assessment.

The aim of the present work was to study the fluorescence formation during the frozen storage and to test the fluorescence ratio as a way of assessing fish quality changes. Previous research on frozen storage (40, 41) had detected some fluorescence that could be related to the quality of product.

In the present experiment, fluorescence detection of interaction compounds was accomplished in both the aqueous and organic phases resulting from the lipid extraction. As in the case of chilling, the fluorescence analysis of the aqueous phase (δF_{aq}) provided better results than that of the organic one (δF_{or}). δF_{aq} showed a good correlation with the storage time and grouped together with indices that reflect fish damage at -18°C (TBA-i and FFA) and at -10°C (TBA-i, FFA and TVB-N).

The satisfactory correlation of fluorescence measurements with secondary lipid oxidation (TBA-i) is in accord with the general theory about fluorescent compounds formation as a result of interaction between carbonyl compounds (electrophilic molecules) formed during lipid oxidation and amine compounds (nucleophilic molecules) present in the fish muscle (13, 14, 42).

1 A good correlation between lipid hydrolysis (FFA) and lipid oxidation (TBA-i) has also been
2 obtained. Some previous experiences have proved the influence of the lipid hydrolysis on the lipid
3 oxidation (43, 44). According to the correlations obtained, a positive influence of the FFA content on the
4 fluorescent compounds formation can be inferred.

5 Results obtained in this work reinforce the role of fluorescence detection of interaction
6 compounds as a tool for quality assessment, especially when processing food with a high content of
7 unsaturated lipids and amino compounds. This is the case of marine species, that are known to contain a
8 high amounts of polyunsaturated fatty acids (4) and of free amino nitrogen compounds (45).

9
10
11
12 Acknowledgments

13 The authors thank *Mr. Oscar Bellón* and *Mr. Marcos Trigo* for technical assistance and the Xunta
14 de Galicia for financial support (*Project XUGA 402 01B93*).

REFERENCES

1. Cheftel, J. and H. Cheftel, *Introduction to the Biochemistry and Technology of Foods*. Editorial Acribia, Zaragoza (Spain), 1976.
2. Hale M. and T. Brown, Fatty acids and lipid classes of three underutilized species and change due to canning, *Marine Fish. Rev.* 45: 4-6 (1983).
3. Aubourg, S., C. Sotelo and J. Gallardo, Changes in flesh lipids and fill oils of albacore (*Thunnus alalunga*) during canning and storage, *J. Agric. Food Chem.* 38: 809-812 (1990).
4. Ackman, R., Fatty acids, in *Marine biogenic lipids, fats and oils*, edited by R. Ackman, CRC Press, Boca Raton, Florida (USA), Vol. 1, 1989, pp. 103-137.
5. Hsieh, R. and J. Kinsella, Oxidation of polyunsaturated fatty acids: mechanisms, products and inhibition with emphasis on fish, *Adv. Food Res. Nutr. Res.* 33: 233-341 (1989).
6. German B., H. Zhang and R. Berger, Role of lipoxygenases in lipid oxidation in foods, in *Lipid oxidation in food*, edited by A. St. Angelo, American Chemical Society, Washington, DC, USA, 1992, pp. 74-92.
7. Pearson, A., J. Love and F. Shorland, Warmed-over flavor in meat, poultry and fish, *Adv. Food Res.* 23: 2-61 (1977).
8. Pigott, G. and B. Tucker, Science opens new horizons for marine lipids in human nutrition, *Food Rev. Intern.* 3: 105-138 (1987).
9. Melton, S., Methodology for following lipid oxidation in muscle foods, *Food Technol.* 37: 105-111, 116, (1983).
10. Kim, R. and F. Labella, Comparison of analytical methods for monitoring autoxidation profiles of authentic lipids, *J. Lipid Res.* 28: 1110-1117 (1987).
11. Pokorný, J., Interactions of oxidized lipids with proteins, *Riv. Ital. Sostanze Grasse* 54: 389-393 (1977).
12. Gardner, H., Lipid hydroperoxide reactivity with proteins and amino acids: A review, *J. Agric. Food Chem.* 27: 220-229 (1979).
13. Leake, L. and M. Karel, Nature of fluorescent compounds generated by exposure of protein to oxidizing lipids, *J. Food Biochem.* 9: 117-136 (1985).
14. Kikugawa, K. and M. Beppu, Involvement of lipid oxidation products in the formation of fluorescent and cross-linked proteins, *Chem. Phys. Lipids* 44: 277-297 (1987).
15. Smith, G., M. Hole and S. Hanson, Assessment of lipid oxidation in Indonesian salted-dried Marine catfish (*Arius thalassinus*), *J. Sci. Food Agric.* 51: 193-205 (1990).
16. Maruf, F., D. Ledward, R. Neale and R. Poulter, Chemical and nutritional quality of Indonesian dried-salted mackerel (*Rastrelliger kanagurta*), *Int. J. Food Sci. Technol.* 25: 66-77 (1990).

- 1 17. Lubis, Z. and K. Buckle, Rancidity and lipid oxidation of dried-salted sardines, *Int. J. Food Sci. Technol.*
2 25: 295-303 (1990).
- 3 18. Miyagawa, K., K. Hirai and R. Takeoze, Tocopherol and fluorescence levels in deep-frying oil and their
4 measurement for oil assessment, *J. Am. Oil Chem. Soc.* 68: 163-166 (1991).
- 5 19. Aubourg, S., R. Pérez-Martín, I. Medina and J. Gallardo, Fluorescence formation by interaction of
6 albacore (*Thunnus alalunga*) muscle with acetaldehyde in a model system, *J. Agric. Food Chem.*
7 40: 1805-1808 (1992).
- 8 20. Aubourg, S., R. Pérez-Martín, I. Medina and J. Gallardo, Fluorescence formation during albacore
9 (*Thunnus alalunga*) thermal processing, *Z. Lebensm. Unters. Forsch.* 195: 332-335 (1992).
- 10 21. Aubourg, S., I. Medina and R. Pérez-Martín, A comparison between conventional and fluorescence
11 detection methods of cooking-induced damage to tuna fish lipids, *Z. Lebensm. Unters. Forsch.*
12 200: 252-255 (1995).
- 13 22. Aubourg, S., C. Sotelo and J. Gallardo, Quality assessment of sardines during storage by
14 measurement of fluorescent compounds, *J. Food Sci.* 62: 295-299 (1997).
- 15 23. Bligh, E. and W. Dyer, A rapid method of total extraction and purification, *Can. J. Biochem. Physiol.* 37:
16 911-917 (1959).
- 17 24. Herbes, S. and C. Allen, Lipid quantification of freshwater invertebrates: method modification for micro
18 quantification, *Can. J. Fish. Aquat. Sci.* 40: 1315-1317 (1983).
- 19 25. Antonacopoulos, N., Verbesserte apparatus zur quantitativer destillation wasserdampfllühtiger stoffe,
20 *Z. Lebensm. Unters. Forsch.* 113: 113-160 (1960).
- 21 26. Vyncke, W., Direct determination of the thiobarbituric acid value in trichloroacetic acid extracts of fish as
22 a measure of oxidative rancidity, *Fette Seifen Anstrichm.* 72: 1084-1087 (1970).
- 23 27. Lowry, R. and I. Tinsley, Rapid colorimetric determination of free fatty acids, *J. Amer. Oil Chem. Soc.*
24 53: 470-472 (1976).
- 25 28. Statsoft, *Statistica for Macintosh*, Statsift and its licensors, Tulsa, Oklahoma, USA, 1994.
- 26 29. Gallardo J., S. Aubourg and R. Pérez-Martín, Lipid classes and their fatty acids at different loci of
27 albacore (*Thunnus alalunga*): Effects of precooking, *J. Agric. Food Chem.* 37: 1060-1064 (1990).
- 28 30. Bennour M., A. El Marrakchi, N. Bouchriti, A. Hamama and M. El Ouadaa, Chemical and
29 microbiological assessments of mackerel (*Scomber scombrus*) stored in ice, *J. Food Protect.* 54:
30 784, 789-792 (1991).
- 31 31. Sotelo, C., C. Piñeiro and R. Pérez-Martín, Denaturation of fish proteins during frozen storage: role of
32 formaldehyde, *Z. Lebensm. Unters. Forsch.* 200: 1-2 (1995).
- 33 32. Rehbein, H., Relevance of trimethylamine oxide demethylase activity and haemoglobin content to
34 formaldehyde production and texture deterioration in frozen stored minced fish muscle, *J. Sci.*
35 *Food Agric.* 43: 261-276 (1988).
- 36 33. Shimasaki, H., O. Privett and I. Hara, Studies of the fluorescent products of lipid oxidation in aqueous
37 emulsion with glycine and on the surface of silica gel, *J. Am. Oil Chem. Soc.* 54: 119-123 (1977).

- 1 34. Quaranta, H. and S. Pérez, Chemical methods for measuring changes in freeze stored fish: A review,
2 *Food Chemistry* 11: 79-85 (1983).
- 3 35. Orlick, B., J. Oehlenschläger and W. Schreiber, Changes in lipids and nitrogenous compounds in cod
4 (*Gadus morhua*) and saithe (*Pollachius virens*) during frozen storage, *Arch. Fish. Wiss.* 41: 89-99
5 (1991).
- 6 36. de Koning, A. and T. Mol, Quantitative quality tests for frozen fish: soluble protein and free fatty acid
7 content as quality criteria for hake (*Merluccius merluccius*) stored at -18°C, *J. Sci. Food Agric.* 54:
8 449-458 (1991).
- 9 37. Shimasaki H., N. Ueta, H. Mowri and K. Inove, Formation of age pigment-like fluorescent substances
10 during peroxidation of lipids in model membranes, *Biochim. Biophys. Acta* 792: 123-129 (1984).
- 11 38. Iio, T. and K. Yoden, Fluorescence formation from hydroperoxide of phosphatidylcholine with amino
12 compound, *Lipids* 23: 65-67 (1988).
- 13 39. Hasegawa, K., Y. Endo and K. Fujimoto, Oxidative deterioration in dried fish model systems assessed
14 by solid sample fluorescence spectrometry, *J. Food Sci.* 57: 1123-1126 (1992).
- 15 40. Davies, H., Fluorescence of fish muscle: description and measurement of changes occurring during
16 frozen storage, *J. Sci. Food Agric.* 33: 1135-1142 (1982).
- 17 41. Davies H. and P. Reece, Fluorescence of fish muscle: causes of change occurring during frozen
18 storage, *J. Sci. Food Agric.* 33: 1143-1151 (1982).
- 19 42. Tappel, A., Measurement of and protection from in vivo lipid peroxidation, in *Free Radicals in Biology*,
20 edited by R. Ackman, Academic Press, New York (USA), 1980; Vol. 4, pp. 1-47.
- 21 43. Miyashita, K. and T. Takagi, Study on the oxidative rate and prooxidant activity of free fatty acids, *J.*
22 *Am. Oil Chem. Soc.* 63: 1380-1384 (1986).
- 23 44. Han, T. and J. Liston, Correlation between lipid peroxidation and phospholipid hydrolysis in frozen fish
24 muscle, *J. Food Sci.* 53: 1917-1918 (1988).
- 25 45. Navarro, P., Valor nutritivo del pescado: I. Pescado fresco, *Rev. Agroquím Tecnol. Aliment.* 31: 330-
26 342 (1991).
- 27

TABLE 1: Quality measurements* and fluorescence shifts* during frozen storage at -18°C**.

ST (months)	TVB-N	CD	TBA-i	FFA	δF_{or}	δF_{aq}	$\delta F_{or}/\delta F_{aq}$
0	25.6 a (0.67)	1.2 bc (0.12)	0.5 a (0.05)	1.1 a (0.14)	0.4 a (0.04)	0.4 a (0.02)	1.2 b (0.10)
0.5	24.2 a (0.26)	1.5 cd (0.14)	1.6 ab (0.28)	4.2 b (0.63)	1.7 b (0.20)	0.3 a (0.02)	5.1 d (0.33)
2	29.5 b (0.52)	0.9 ab (0.03)	2.8 ab (0.61)	3.0 b (0.33)	1.9 b (0.14)	0.3 a (0.02)	5.7 d (0.64)
4	27.2 ab (1.08)	1.2 bc (0.01)	5.0 bc (0.87)	4.1 b (0.20)	1.5 b (0.02)	1.2 ab (0.10)	1.3 b (0.13)
8	24.1 a (0.96)	0.8 ab (0.07)	3.9 ab (0.11)	4.0 b (0.16)	2.4 b (0.40)	1.0 ab (0.07)	2.3 c (0.30)
12	35.6 c (0.30)	2.2 e (0.37)	8.6 c (0.67)	7.1 c (0.37)	2.3 b (0.60)	2.4 b (0.33)	0.9 ab (0.15)
18	29.6 b (2.63)	1.9 de (0.14)	9.6 c (2.10)	6.4 c (0.60)	3.7 c (0.15)	5.9 c (0.42)	0.6 ab (0.04)
24	30.2 b (0.56)	0.7 a (0.03)	29.9 d (2.32)	11.8 d (0.63)	2.2 b (0.45)	12.6 d (1.28)	0.2 a (0.06)

* Mean values of three determinations. Values in the same column followed by different letters are significantly different ($p < 0.05$). Standard errors of the means are indicated in brackets.

** Abbreviations: ST (storage time), TVB-N (total volatile base-nitrogen), CD (conjugated dienes), TBA-y (thiobarbituric acid index), FFA (free fatty acids), δF_{or} (fluorescence shift in the organic phase), δF_{aq} (fluorescence shift in the aqueous phase)

TABLE 2: Quality measurements* and fluorescence shifts* during frozen storage at -10°C^{**} .

ST (days)	TVB-N	CD	TBA-i	FFA	δF_{or}	δF_{aq}	$\delta F_{\text{or}}/\delta F_{\text{aq}}$
0	25.6 b (0.67)	1.2 abc (0.12)	0.5 a (0.05)	1.1 a (0.14)	0.4 a (0.04)	0.4 a (0.02)	1.2 a (0.10)
3	28.5 c (1.13)	1.4 c (0.06)	1.9 ab (0.16)	3.8 b (0.47)	0.7 a (0.05)	0.3 a (0.02)	2.6 a (0.23)
10	22.2 a (0.52)	1.3 bc (0.16)	2.9 b (0.41)	3.1 ab (0.52)	2.0 b (0.20)	0.3 a (0.02)	8.1 c (1.36)
25	29.3 cd (0.45)	1.1 ab (0.02)	5.4 c (0.41)	6.9 c (0.40)	1.8 b (0.08)	0.4 a (0.05)	5.2 b (0.65)
60	28.4 c (0.93)	1.0 a (0.07)	5.2 C (0.20)	7.8 C (1.74)	3.1 C (0.59)	0.5 b (0.07)	5.7 B (0.93)
120	31.1 d (0.28)	1.4 c (0.09)	8.7 d (1.07)	8.5 c (0.07)	1.8 b (0.14)	2.2 c (0.05)	0.8 a (0.08)

* Mean values of three determinations. Values in the same column followed by different letters are significantly different ($p < 0.05$). Standard errors of the means are indicated in brackets.

** Abbreviations as specified in Table 1.

Table 3: Correlation matrix for different parameters*** (storage time and quality estimations) measured during sardine storage at -18°C.

	TVB-N	CD	TBA-i	FFA	δF_{or}	δF_{aq}
ST	0.52*	0.09	0.79**	0.82**	0.65*	0.80**
TVB-N		0.43*	0.36	0.46*	0.35	0.34
CD			-0.22	-0.10	0.23	-0.15
TBA-i				0.87**	0.35	0.92**
FFA					0.50*	0.83**
δF_{or}						0.36

* Significance: $p < 0.05$.

** Significance: $p < 0.01$.

*** Abbreviations as specified in Table 1.

Table 4: Correlation matrix for different parameters (storage time and quality estimations) measured during sardine storage at -10°C.

	TVB-N	CD	TBA-i	FFA	δF_{or}	δF_{aq}
ST	0.55*	-0.07	0.94**	0.88**	0.72*	0.72*
TVB-N		0.07	0.61*	0.71*	0.14	0.60*
CD			0.14	-0.03	-0.24	0.27
TBA-i				0.84**	0.56*	0.81**
FFA					0.73*	0.59*
δF_{or}						0.17

* Significance: $p < 0.05$.

** Significance: $p < 0.01$.

*** Abbreviations as specified in Table 1.

FIGURE LEGENDS

Figure 1: Principal component analysis for different parameters (storage time and quality indices) measured at -18°C storage (abbreviations as specified in Table 1).

Figure 2: Principal component analysis for different parameters (storage time and quality indices) measured at -10°C storage (abbreviations as specified in Table 1).