Seasonal changes in the diet of a critically endangered seabird and the importance of trawling discards

Joan Navarro · Maite Louzao · José Manuel Igual · Daniel Oro ·
Antonio Delgado · José Manuel Arcos · Meritxell Genovart ·
Keith A. Hobson · Manuela G. Forero

Abstract Pelagic seabirds obtain food from oceans where the availability of their prey changes rapidly both seasonally and spatially. Here, we investigated changes in the trophic habits of the critically endangered Balearic shearwater (*Puffinus mauretanicus*) through the breeding season and tested for dietary differences between sexes and age classes. We analysed $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in blood of adults during the pre-incubation, incubation and chick-rearing periods and of their chicks. Using a two-isotope mixing model, we estimated dietary contributions based on isotope values from potential prey species which included small pelagic

species available naturally and demersal fish species available only from trawling discards. Balearic shearwaters showed clear isotopic and dietary variation through the breeding season. During pre-incubation, breeding adults appeared to exploit demersal fish, whereas during the incubation and chick-rearing period, they fed mainly on pelagic anchovies (*Engraulis encrasicolus*) and pilchards (*Sardina pilchardus*). Similarly, chicks were fed mainly with anchovies, a resource with a high energetic value. This variation in the dietary habits of adult shearwaters during the breeding season was probably related to both natural and fishery-induced seasonal changes in the availability of potential prey species within their main feeding grounds. However, changes in the nutritional requirements of the shearwaters could also play an important role. Indeed, diet differed between sexes during pre-incubation: females fed less on trawling discards and more on small pelagic fish than males. This sexual segregation in diet could be the consequence of higher nutritional requirements of females during this period. Our study reveals the differential importance of both trawling discards and small pelagic fish species for a pelagic seabird depending on the breeding period and illustrates the importance of considering the entire breeding season when making inferences about the importance of specific prey in seabird dietary studies.

Communicated by R. Lewison.

J. Navarro (&) · M. G. Forero
Department of Conservation Biology,
Estación Biológica de Doñana, CSIC,
Av. Américo Vespucio, s/n, 41092 Sevilla, Spain
e-mail: joan@ebd.csic.es

M. Louzao
Helmholtz Centre for Environmental Research-UFZ,
Permoserstraße 15, 04318 Leipzig, Germany

J. M. Igual · D. Oro · M. Genovart
Institut Mediterrani d'Estudis Avançats, IMEDEA (CSIC-UIB),
C/Miquel Marqués 21, 07190 Esporles, Mallorca, Spain

A. Delgado
Estación Experimental del Zaidín, CSIC,
Profesor Albareda 1, Granada 18008, Spain

J. M. Arcos
Sociedad Española de Ornitología, SEO/BirdLife,
Delegación de Cataluña, C/Murcia 2, 08026 Barcelona, Spain

K. A. Hobson
Environment Canada, 11 Innovation Blvd.,
Saskatoon, SK S7N3H5, Canada

Introduction

Pelagic seabirds obtain their food from oceans where resources are patchily distributed and their abundance and availability changes seasonally (Brooke 2004). For this reason, their foraging efficiency and adaptation to these constraints are important components affecting their population dynamics (Lewis et al. 2006). During the breeding season,

when food demand is high, adult seabirds show flexible feeding strategies, which help them cope with unpredictable prey availability and to cover nutritional requirements of reproduction (Brooke 2004). Beyond natural variability of prey, seabirds also face the impact of human activities such as commercial fisheries. Fisheries can not only deplete marine fish stocks (Myers and Worm 2003) but also generate large volumes of discarded undersized fishes and non-target species that several opportunistic seabirds have learned to exploit (i.e. Oro 1999; Tasker et al. 2000; Montevecchi 2002; Becker and Beissinger 2006).

This situation is especially true in the western Mediterranean ecosystem, where trawling discards have an impact on the foraging behaviour, breeding performance and population dynamics of opportunistic seabirds, including endangered species (Oro 1999; Arcos and Oro 2002a; Oro et al. 2004b; Louzao et al. 2006b). The Balearic shearwater (*Puffinus mauretanicus*) is one of the rarest seabirds in the world, currently listed as Critically Endangered by the IUCN (Oro et al. 2004a; BirdLife International 2008). The species is endemic to the western Mediterranean and about 2,200 breeding pairs breed in the Balearic Archipelago (BirdLife International 2008). During the breeding season, the species mainly forages over the eastern Iberian continental shelf (Louzao et al. 2006a), a highly productive and spawning area for small pelagic fish such as anchovies (*Engraulis encrasicolus*) and pilchards (*Sardina pilchardus*) and where important fishing fleets operate (Abad et al. 1998; Agostini and Bakun 2002; Palomera et al. 2007). Although the Balearic shearwater appears to primarily forage on shoals of small pelagic fish, mainly anchovies and pilchards, different ecological models have also shown the importance of demersal and pelagic fish provided by trawling discards (Arcos and Oro 2002a; Abelló et al. 2003; Louzao et al. 2006b). Similarly, trawling discards also seem to improve the breeding performance of this and other endangered seabirds in the Mediterranean (Oro 1999; Louzao et al. 2006b). However, there are few data on the relative importance of both pelagic and demersal fish during the pre-laying and incubation period or on potential differences in diet related to sex or between adults and chicks.

Currently, the stable isotope approach can augment conventional means of dietary analysis since stable isotopes reflect time-integrated dietary records and thus eliminate some of the shortcomings of traditional dietary studies, particularly when used in combination with the stable isotope determination of potential prey (reviewed by Forero and Hobson 2003; Barrett et al. 2007). For a bird, the size of a shearwater, whole blood stable isotope values integrate dietary information over 3–4 weeks prior to sampling (Bearhop et al. 2002; Pearson et al. 2003). Thus, changes in dietary habits during the breeding season can be investigated by repeated blood sampling during breeding. Stable

isotopes of nitrogen ($\delta^{15}\text{N}$) are indicators of trophic positions as consumers are predictably enriched in ^{15}N relative to their food (Post 2002; Vanderklift and Ponsard 2003). Stable carbon isotope values ($\delta^{13}\text{C}$) give information on primary production and are useful to trace origin of the prey consumed (Forero and Hobson 2003). Furthermore, combining stable isotope values of the consumers and their prey, isotopic mixing models can be applied to obtain estimates of the relative contribution of each potential prey to the diet habits of the consumer (Phillips and Gregg 2001).

Here, our main objective was to examine the trophic ecology of the Balearic shearwater throughout the breeding season. Specifically, we investigated whether breeding shearwaters showed seasonal changes in diet and assessed whether there were age- (chicks vs. adults) and sex-related differences. Moreover, in order to disentangle the effect of fishing activity on the foraging ecology of shearwaters, we combined the analyses of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in blood of adults at different breeding stages with those of their potential prey items, both naturally available pelagic species and trawling discards.

Materials and methods

Fieldwork

This study was conducted in 2003 in Sa Cella (Mallorca, Illes Balears, Spain, 39°30'N; 2°33'E, Fig. 1), a colony of about 200 breeding pairs of Balearic shearwater (Oro et al. 2004a). During the breeding period, we sampled individual males and females at pre-incubation (February), incubation (March–April) and chick-rearing (May–June) periods (see Table 1 for number of males and females sampled at each breeding stage). We also sampled chicks at the end of the chick-rearing period (mid June). From each bird, we took 0.1 ml of blood that was preserved in 70% ethanol and further used for both molecular sexing and stable isotope analyses. We also collected samples of potential prey of shearwaters during May 2003 from an experimental oceanographic trawling cruise along the Iberian continental shelf (MEDITS—Mediterranean International Trawl Survey, Bertrand et al. 2002; Table 2). We focused on the Ebro Delta area because it is the main foraging ground for the Balearic shearwater during the breeding season (Arcos and Oro 2002a; Abelló et al. 2003; Louzao et al. 2006a). To select potential prey, we followed Arcos and Oro (2002a) who determined a feeding preference index for the Balearic shearwater selecting nine fish species and one cephalopod species commonly inhabiting the pelagic (4 species) and demersal sea compartment (6 species, Table 2). From each specimen, we sampled a muscle that was stored frozen at -21°C .

Fig. 1 Location of the breeding colony (Sa Cella, Mallorca, Spain) and the main foraging ground (Ebro Delta area) for the Balearic shearwater, as well the location of the main trawling ports. The picture (courtesy of J. M. Arcos) shows a Balearic shearwater following a Wshing boat

Table 1 Sample size (n) and mean $\pm$ standard deviation of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of Balearic shearwater throughout the breeding season

	n	$\delta^{15}\text{N}$ (‰)	$\delta^{13}\text{C}$ (‰)
Adults			
Pre-incubation			
Males	27	10.56 $\pm$ 0.52	17.69 $\pm$ 0.32
Females	12	10.01 $\pm$ 0.76	17.99 $\pm$ 0.25
Incubation			
Males	12	9.76 $\pm$ 0.57	18.86 $\pm$ 0.40
Females	9	10.12 $\pm$ 0.58	18.68 $\pm$ 0.53
Chick rearing			
Males	27	10.51 $\pm$ 0.21	19.23 $\pm$ 0.16
Females	16	10.26 $\pm$ 0.32	19.35 $\pm$ 0.23
Chicks			
Males	15	10.73 $\pm$ 0.24	20.01 $\pm$ 0.16
Females	27	10.51 $\pm$ 0.26	19.97 $\pm$ 0.22

Stable isotopes analyses

Ethanol was removed from blood samples prior to analysis by decanting and freeze-drying. For prey muscle, we removed two aliquots from each lyophilized muscle sample; one aliquot was immediately prepared for $\delta^{15}\text{N}$ analysis. The other underwent lipid extraction prior to $\delta^{13}\text{C}$ analysis (Sholto-Douglas et al. 1991; Schlechtriem et al. 2003). Lipids were extracted using the methods of Smedes (1999). All samples (about 0.5–1 mg) were combusted at 1,020°C using continuous-flow isotope-ratio mass spectrometry (CFIRMS) system by means of a Carlo Erba 1500NC elemental analyser interfaced with a Delta Plus XL mass spectrometer. All isotope abundances are expressed in δ -notation as parts per thousand (‰) deviation from the IAEA standards AIR ($\delta^{15}\text{N}$) and VPDB ($\delta^{13}\text{C}$). Based on within-run replicate measurements of lab standards (urea, shark cartilage), we estimated measurement

error to be ± 0.2 and $\pm 0.1\text{‰}$ for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values, respectively.

Statistical analyses

We tested the effect of breeding stage (pre-incubation, incubation and chick rearing) and sex of adults on stable isotope values using analysis of variance (ANOVA). As segregation in the diet between sexes could change depending on the breeding stage, interaction between sex and breeding stage was also tested in the model. We tested the effect of the age and sex of chicks (adults during chick rearing vs. male and female chicks) on stable isotope values by ANOVA. Similarly, we examined differences among prey groups using ANOVA. Post-hoc comparisons were tested with Tukey's test. Stable isotope values were previously log-transformed to meet normality. All statistical analyses were conducted by using SAS 9.1 software.

Isotope mixing model

Dietary composition of the Balearic shearwater at each breeding stage was estimated based on the isotopic values of potential prey. We created four different prey groups according to the dietary importance of each prey species for the Balearic shearwater (Arcos and Oro 2002a): anchovies, pilchards, pelagic Wsh (*Gadiculus argenteus* and *Micromesistius poutassou*) and demersal species (Table 2). Statistical differences in stable isotope values among these prey groups were tested by an ANOVA and Tukey post-hoc comparisons). Because pilchards and pelagic Wsh did not differ in their stable isotope values (see "Results"), we used only three clusters, anchovies, pelagic Wsh (including pilchards) and demersal Wsh, to further estimate the proportion of each potential prey consumed by the Balearic shearwater. We then created consumer end members for blood of shearwaters feeding exclusively on each of the

Table 2 Sample size (*n*) and mean $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of the potential prey species for the Balearic shearwater, sampled in the Ebro Delta area (western Mediterranean) during spring 2003

Species	Habitat	<i>n</i>	$\delta^{15}\text{N}$ (‰)	$\delta^{13}\text{C}$ (‰)
Fish				
Anchovy <i>Engraulis encrasicolus</i>	P	10	8.85 $\pm$ 0.48	$\bar{\delta}$ 19.96 $\pm$ 0.46
Blue whiting <i>Micromesistius poutassou</i>	P	10	8.90 $\pm$ 1.09	$\bar{\delta}$ 18.85 $\pm$ 0.86
Pilchard <i>Sardina pilchardus</i>	P	10	7.24 $\pm$ 1.46	$\bar{\delta}$ 18.91 $\pm$ 0.71
Silvery cod <i>Gadiculus argenteus</i>	P	9	6.81 $\pm$ 3.09	$\bar{\delta}$ 18.25 $\pm$ 0.37
Blotched picarel <i>Spicara Xexuosa</i>	D	10	9.76 $\pm$ 0.73	$\bar{\delta}$ 19.31 $\pm$ 0.44
Brown comber <i>Serranus hepatus</i>	D	9	10.68 $\pm$ 0.57	$\bar{\delta}$ 17.64 $\pm$ 0.47
Greater forkbeard <i>Phycis blennoides</i>	D	10	8.65 $\pm$ 0.42	$\bar{\delta}$ 18.44 $\pm$ 0.15
Jewel lanternfish <i>Lampanyctus crocodilus</i>	D	8	7.57 $\pm$ 0.56	$\bar{\delta}$ 18.74 $\pm$ 0.45
Tonguesole <i>Symphurus nigrescens</i>	D	8	11.41 $\pm$ 0.56	$\bar{\delta}$ 17.37 $\pm$ 0.29
Cephalopods				
Common squid <i>Alloteuthis subulata</i>	D	6	9.19 $\pm$ 1.61	$\bar{\delta}$ 17.62 $\pm$ 0.54

Habitat type for each prey species is indicated (*P* pelagic, *D* demersal) following FishBase (<http://www.fishbase.org>)

Fig. 2 Stable isotope values in blood ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) of breeding adults during pre-incubation (a), incubation (b), chick rearing (c) and chicks (d) of Balearic shearwater and their prey. White and grey circles indicate male and female shearwaters, respectively. Mean ($\pm$ SE) of each group of potential prey items are indicated as follow: anchovies (Willed diamond), pilchards (Willed square), pelagic fish (Willed circle, *Gadiculus argenteus*, *Micromesistius poutassou*) and demersal fish (Willed triangle, *Serranus hepatus*, *Lampanyctus crocodilus*, *Spicara Xexuosa*, *Phycis blennoides*, *Symphurus nigrescens*, and *Alloteuthis subulata*). Boxes represent the range of expected values in the consumer, considering 99% CI for prey and accounting for diet-blood isotopic fractionation factors of 1.7‰ for $\delta^{15}\text{N}$ and 0.9‰ for $\delta^{13}\text{C}$ (following Ramos et al. 2009). Partial labelling for expected dietary isotopic values (rectangular frames) is shown to enhance clarity

three isotopically different prey types (99% CI, Fig. 2) using diet-blood isotopic fractionation values between prey muscle and whole blood of 1.7‰ for $\delta^{15}\text{N}$ and 0.9‰ for $\delta^{13}\text{C}$. These values are derived from an isotopic study on the yellow-legged gull *Larus michahellis* from a breeding population that feed exclusively on pelagic fish (Ramos et al. 2009). We applied the Iso-Error dual-isotope, 3-source mixing model of Phillips and Gregg (2001).

(Willed triangle, *Serranus hepatus*, *Lampanyctus crocodilus*, *Spicara Xexuosa*, *Phycis blennoides*, *Symphurus nigrescens*, and *Alloteuthis subulata*). Boxes represent the range of expected values in the consumer, considering 99% CI for prey and accounting for diet-blood isotopic fractionation factors of 1.7‰ for $\delta^{15}\text{N}$ and 0.9‰ for $\delta^{13}\text{C}$ (following Ramos et al. 2009). Partial labelling for expected dietary isotopic values (rectangular frames) is shown to enhance clarity

Results

Seasonal changes in stable isotope values in Balearic shearwater

Stable isotope values of Balearic shearwater varied depending of the breeding stage (Table 1). During incubation, breeding adults showed lower $\delta^{15}\text{N}$ and higher $\delta^{13}\text{C}$ values

Table 3 Summary of the ANOVA tests examining variation in stable isotopes between breeding stages for 66 males and 37 females Balearic shearwaters

Parameter	EVect	$F_{[df]}$	P	Post-hoc
$\delta^{15}\text{N}$ (‰)	Sex	5.86 _[1,102]	0.17	
	Stage $\times$ sex	5.68 _[2,102]	<0.01	
	Stage	5.86 _[2,102]	<0.01	I $\times$ PI, CR
$\delta^{13}\text{C}$ (‰)	Sex	1.56 _[1,102]	0.21	
	Stage $\times$ sex	4.09 _[2,102]	0.02	
	Stage	208.13 _[2,102]	<0.001	PI $\times$ I, CR I $\times$ CR

Symbols designating stage combination in Tukey post-hoc test summaries are: *PI* pre-incubation; *I* incubation; *CR* chick rearing. Pairs of means differing significantly ($P = 0.05$) by Tukey test are linked with an 'x'

than during pre-incubation and chick rearing (Table 1, and see post-hoc comparison in Table 3). Although we did not detect a significant effect of sex throughout breeding on $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ isotopic values (Table 3), males showed higher $\delta^{15}\text{N}$ values than females during the pre-incubation (Table 1, $T = 2.48$, $df = 76$, $P = 0.01$). Male chicks had higher $\delta^{15}\text{N}$ values than female chicks (Table 1, $T = 2.55$, $df = 42$, $P = 0.01$), whereas $\delta^{13}\text{C}$ values did not differ between male and female chicks (Table 1, $T = 1.056$, $df = 42$, $P = 0.57$). During the chick-rearing period, adults had lower $\delta^{15}\text{N}$ (adults vs. male chicks: $T = 3.77$, $df = 56$, $P < 0.001$; adults vs. female chicks: $T = 2.55$, $df = 40$, $P = 0.01$) and higher $\delta^{13}\text{C}$ values than chicks (adults vs. male chicks: $T = 12.63$, $df = 56$, $P < 0.001$; adults vs. female chicks: $T = 1.056$, $df = 40$, $P = 0.57$; Table 1).

Isotopic values of potential prey

Overall, both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values differed among the four prey groups (Table 2; Fig. 2. For $\delta^{15}\text{N}$; $F_{2,97} = 9.74$, $P < 0.0001$. For $\delta^{13}\text{C}$; $F_{2,97} = 17.85$, $P < 0.0001$). Demersal species showed higher $\delta^{15}\text{N}$ values than pilchards and pelagic Wsh, and similar values as anchovies (Table 2; Fig. 2. Tukey test, $P < 0.0001$, < 0.0001 and $= 0.61$, respectively). Concerning $\delta^{13}\text{C}$ values, anchovies had lower values than demersal species, pilchards and pelagic Wsh (Table 2; Fig. 2, Tukey test, $P < 0.0001$, 0.0001 and 0.001 , respectively). Only pelagic Wsh and pilchards had similar $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values (Table 2; Fig. 2, $P = 0.67$ and 0.81 , respectively).

Dietary changes during breeding season inferred by isotopic mixing model

The proportion of each prey type in the diet of the Balearic shearwater changed throughout the breeding season

Fig. 3 Results of the Iso-Error mixing model (Phillips and Gregg 2001) for the percentage (mean $\pm$ SD) of each prey group in the diet of male and female Balearic shearwaters during the different breeding stages. The potential groups were; anchovies, pelagic Wsh (*Gadiculus argenteus*, *Micromesistius poutassou*, and *Sardina pilchardus*) and demersal Wsh (see Table 2)

(Fig. 3). During the pre-incubation period, the diet of adult males was composed mostly of demersal Wsh (58.5 $\pm$ 8.9%), followed of pelagic Wsh (29.0 $\pm$ 10.8%) and then anchovies (12.4 $\pm$ 6.7). In contrast, the diet of females during pre-incubation was composed mainly of pelagic Wsh (53.8 $\pm$ 17.8%), followed by anchovies and demersal Wsh in a similar proportion (23.6 $\pm$ 8.4 and 22.6 $\pm$ 13.9%, respectively). During incubation (Fig. 3), the model estimated that the diet of males and females was composed mainly of anchovies (57.1 $\pm$ 11.7 and 66.6 $\pm$ 13.6%, respectively) and pelagic Wsh (42.0 $\pm$ 16.5 and 33.8 $\pm$ 15.2%, respectively), whereas demersal Wsh were not present. During chick rearing, both sexes fed mainly on anchovy (males = 93.2 $\pm$ 10.1%, females = 79.3 $\pm$ 11.1%) and secondly on pelagic Wsh (less than 15%) and demersal species were not consumed. Finally, the model indicated that adults fed their male and female chicks mainly with anchovies (98.3 $\pm$ 15.3 and 88.9 $\pm$ 14.7%, respectively), and in much lower proportion with pelagic Wsh (1.7 $\pm$ 16.6 and 11.0 $\pm$ 15.3%).

Discussion

Stable isotope analyses revealed clear changes in the diet of the Balearic shearwater throughout the breeding season, and a subtler, apparent difference in the diet between males and females during the pre-incubation period. This variation in diet was probably related to complex interrelationships affecting natural- and fishery-induced seasonal changes in the availability of their potential prey and to nutritional requirements during reproduction. These results

illustrate the importance of considering the entire breeding season when conducting studies of trophic ecology of seabirds.

It is well documented that Wshing activity provides substantial food for opportunistic seabirds through the discarding of non-target demersal and pelagic species (e.g. Oro 1999; Furness and Tasker 2000; Tasker et al. 2000; Arcos and Oro 2002b; Votier et al. 2004). Similar to previous studies, we found that Balearic shearwaters clearly exploited this artiWcial resource during the breeding period, but were able to provide a quantitative estimate of the relative importance of this resource during the breeding season. During pre-incubation, breeding adults fed on demersal species, prey only available for surface feeders such as the Balearic shearwater through trawling discards (Arcos et al. 2008). Breeding shearwaters probably also obtained some of their pelagic Wsh and anchovies from the trawling discards in addition to natural-foraging behaviours since they can become available in addition to demersal species during Wshing operations at the Ebro Delta (Arcos and Oro 2002a, b; Louzao et al. 2006b).

So, why did breeding shearwaters apparently only feed on demersal species during pre-incubation? In the north eastern Iberian shelf, the main feeding area for Balearic shearwaters (Louzao et al. 2006a), the abundance of anchovies and pilchards changes seasonally during the year, showing lower abundances between February–March, coinciding with the pre-incubation period of Balearic shearwater (Lloret et al. 2004). Therefore, breeding birds probably included demersal species in their diet because they were relatively more available during this period. During April–June, coinciding with the incubation and chick-rearing period of the Balearic shearwater, the abundance of anchovies and pilchards in the Ebro Delta is higher than during the pre-incubation period (Lloret et al. 2004), supporting the hypothesis that breeding adults exploit the most available resource within their feeding ground. Moreover, the exploitation of anchovies and other pelagic Wsh (including pilchards) might help birds meet the high energetic requirements of breeding due to their high energetic content (Arcos and Oro 2002a; Grémillet et al. 2008). For instance, adults fed their chicks mainly with anchovies, a high energetic prey item (Arcos and Oro 2002a; Grémillet et al. 2008). Although both chicks and rearing adult shearwaters apparently consumed the same trophic resources, we found that chicks showed higher $\delta^{15}\text{N}$ values than adults rearing chicks. This isotopic diVerence could be related to physiological eVects related to the chick-growing process. Because chicks are growing, their increase in catabolism of protein stores could result in increases in $\delta^{15}\text{N}$ values (Navarro et al. 2007; Williams et al. 2007).

On the other hand, the lower $\delta^{13}\text{C}$ values of chicks compared with adults could be related to the high lipid reserves of the chicks that could depleted their blood $\delta^{13}\text{C}$ values (Phillips and Hamer 1999; Thompson et al. 2000; Williams et al. 2007).

The dietary habits of male and female shearwaters diVered during the pre-incubation period; females fed less on demersal Wsh than males and consumed a higher proportion of anchovies and pelagic Wsh. This sexual segregation in diet could be the consequence of diVerential nutritional requirements of males and females during pre-incubation relating to the egg formation (Williams 2005). In general, the nutritional value of demersal species in terms of lipids and proteins is lower than pelagic species (Arcos and Oro 2002a; Grémillet et al. 2008). For this reason, females, constrained by egg formation, probably need to feed on species of higher nutritional content (Grémillet et al. 2008). Previous studies suggested a positive relationship between hatching success and the availability of pelagic Wsh (Louzao et al. 2006b), and between egg size and trophic level of Balearic shearwaters (Louzao et al. 2008).

In conclusion, this study illustrates and reinforces the importance of both pelagic and demersal Wsh as trophic resources for the Balearic shearwater during the breeding period (Arcos and Oro 2002a, b; Louzao et al. 2006b; Arcos et al. 2008). During pre-incubation, when the natural abundance of prey of high-energy content such as anchovy and pilchards was low, breeding shearwaters exploited demersal Wsh species, an artiWcial resource only available for the shearwaters through trawling discards. Later, during incubation and chick rearing, the diet of both adults and chicks was composed exclusively of anchovies and pelagic Wsh, probably related to their higher natural abundance and their higher energetic contents. Furthermore, female shearwaters showed clearly diVerent trophic habits than males during pre-incubation, female shearwaters consumed less demersal Wsh and more pelagic Wsh species than male shearwaters during the pre-incubation probably linked to the egg formation requirements. Even though trawling discards represent a additional and easy (due to their predictability in space and time) foraging resource for shearwaters and the potential Wtness –and conservation- beneWts for birds, it is important to consider discards as an undesirable oVset of a non-selective, damaging human activity for marine ecosystems, and thus a phenomenon to be eradicated in the near future.

Acknowledgments We extend thanks to Isabel Afán and to Arsenio Granados for their Weldwork and stable isotope analyses, respectively. Fish samples were collected within the frame of the MEDITS project, with the help of Luis Gil de Sola, Pere Abelló and the scientists and crew at R/V Cornide de Saavedra (Spanish Institute of Oceanography, IEO). Research funds were provided by several grants from the Spanish

Government (P06-RNM-02362, REN2002-00450, BOS2003-01960 and CGL2006-04325/BOS). Maite Louzao was supported by a PhD fellowship of the Balearic Government and a post-doctoral fellowship of the Spanish Ministry of Education and Science (Ref. EX-2007-1148). Meritxel Genovart was funded by an I3P post-doctoral fellowship from the Spanish Ministry of Education and Science. Four referees contributed to improve a previous draft.

References

- Abad R, Miquel J, Iglesias M, Álvarez F (1998) Acoustic estimation of abundance and distribution of anchovy in the NW Mediterranean. *Sci Mar* 62:37–43
- Abelló P, Arcos JM, Gil de Sola L (2003) Geographical patterns of seabird attendance to a research trawler along the Iberian Mediterranean coast. *Sci Mar* 67:69–75
- Agostini VN, Bakun A (2002) 'Ocean triads' in the Mediterranean Sea: physical mechanisms potentially structuring reproductive habitat suitability (with example application to European anchovy, *Engraulis encrasicolus*). *Fish Oceanogr* 11:129–142
- Arcos JM, Oro D (2002a) Significance of fisheries discards for a threatened Mediterranean seabird, the Balearic shearwater *Pu Ynus mauretanicus*. *Mar Ecol Prog Ser* 239:209–220
- Arcos JM, Oro D (2002b) Significance of nocturnal purse seine fisheries for seabirds: a case study of the Ebro Delta (NW Mediterranean). *Mar Biol* 141:277–286
- Arcos JM, Louzao M, Oro D (2008) Ecosystem impacts and management in the Mediterranean: seabirds point of view. In: Nielsen JL, Dodson JJ, Friedland K, Hamon TR, Musick J, Verspoor E (eds) *Reconciling Fisheries with conservation: proceedings of the Fourth World Fisheries Congress*. American Fisheries Society, Symposium 49, Bethesda, pp 1471–1479
- Barrett RT, Camphuysen KCJ, Anker-Nilssen T, Chardine JW, Furness RW, Garthe S, Hüppop O, Leopold MF, Montevecchi WA, Veit RR (2007) Diet studies of seabirds: a review and recommendations. *ICES J Mar Sci* 64:1675–1691
- Bearhop S, Waldron S, Votier SC, Furness RW (2002) Factors that influence assimilation rates and fractionation of nitrogen and carbon stable isotopes in avian blood and feathers. *Physiol Biochem Zool* 75:451–458
- Becker BH, Beissinger SR (2006) Centennial decline in the trophic level of an endangered seabird after fisheries decline. *Conserv Biol* 20:470–479
- Bertrand JA, Gil de Sola L, Papaconstantinou C, Relini G, Souplet A (2002) The general specifications of the MEDIS surveys. *Sci Mar* 66:9–17
- BirdLife International (2008) *Pu Ynus mauretanicus*. IUCN 2008. 2008 IUCN Red List of Threatened Species
- Brooke M (2004) *Albatrosses and petrels across the world*. Oxford University Press, Oxford
- Forero MG, Hobson KA (2003) Using stable isotopes of nitrogen and carbon to study seabird ecology: applications in the Mediterranean seabird community. *Sci Mar* 67:23–32
- Furness RW, Tasker ML (2000) Seabird-fishery interactions: quantifying the sensitivity of seabirds to reductions in sandeel abundance, and identification of key areas for sensitive seabirds in the North Sea. *Mar Ecol Prog Ser* 202:253–264
- Grémillet D, Pichegru L, Kuntz G, Woakes AG, Wilkinson S, Crawford RJM, Ryan PG (2008) A junk-food hypothesis for gannets feeding on fishery waste. *Proc R Soc B* 275:1149–1156
- Lewis S, Grémillet D, Daunt F, Ryan P, Crawford RJM, Wanless S (2006) Using behavioural variables to identify proximate causes of population change in a seabird. *Oecologia* 147:606–614
- Lloret J, Palomera I, Salat J, Sole I (2004) Impact of freshwater input and wind on landings of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) in shelf waters surrounding the Ebro (Ebro) River delta (north-western Mediterranean). *Fish Oceanogr* 13:102–110
- Louzao M, Hyrenbach KD, Arcos JM, Abelló P, de Sola LG, Oro D (2006a) Oceanographic habitat of an endangered Mediterranean procellariiform: implications for marine protected areas. *Ecol Appl* 16:1683–1695
- Louzao M, Igual JM, McMinn M, Aguilar JS, Triay R, Oro D (2006b) Small pelagic fish, trawling discards and breeding performance of the critically endangered Balearic shearwater: improving conservation diagnosis. *Mar Ecol Prog Ser* 318:247–254
- Louzao M, Igual JM, Genovart M, Forero MG, Hobson KA, Oro D (2008) Spatial variation in egg size of a top predator: interplay of body size and environmental factors? *Acta Oecol* 34:186–193
- Montevecchi WA (2002) Interactions between fisheries and seabirds. In: Schreiber EA, Burger J (eds) *Biology of marine birds*. CRC Press, Boca Raton, pp 527–557
- Myers RA, Worm B (2003) Rapid worldwide depletion of predatory fish communities. *Nature* 423:280–283
- Navarro J, González-Solís J, Viscor G (2007) Nutritional and feeding ecology in the Cory's shearwater (*Calonectris diomedea*) during breeding. *Mar Ecol Prog Ser* 351:261–271
- Oro D (1999) Trawler discards: a threat or a resource for opportunistic seabirds? In: Adams NJ, Slotow RH (eds) *Proceedings 22 International Ornithological Congress Durban*. BirdLife South Africa, Johannesburg, pp 717–730
- Oro D, Aguilar JS, Igual JM, Louzao M (2004a) Modelling demography and extinction risk in the endangered Balearic shearwater. *Biol Conserv* 116:93–102
- Oro D, Cam E, Pradel R, Martínez-Abraín A (2004b) Influence of food availability on demography and local population dynamics in a long-lived seabird. *Proc R Soc B* 271:387–396
- Palomera I, Olivar MP, Salat J, Sabatés A, Coll M, Gracia A, Morales-Nin B (2007) Small pelagic fish in the NW Mediterranean Sea: an ecological review. *Prog Ocean* 74:37–396
- Pearson SF, Levey DJ, Greenberg CH, Martínez del Río C (2003) Effects of elemental composition on the incorporation of dietary nitrogen and carbon isotopic signatures in an omnivorous songbird. *Oecologia* 135:516–523
- Phillips DL, Gregg JW (2001) Uncertainty in source partitioning using stable isotopes. *Oecologia* 127:171–179
- Phillips RA, Hamer KC (1999) Lipid reserves, fasting capability and the evolution of nestling obesity in procellariiform seabirds. *Proc R Soc B* 266:1329–1334
- Post DM (2002) Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83:703–718
- Ramos R, Ramírez F, Sanpera C, Jover L, Ruiz X (2009) Feeding ecology of yellow-legged gulls (*Larus michahellis*) in the western Mediterranean: a comparative assessment using conventional and isotopic methods. *Mar Ecol Prog Ser* 377:289–297
- Schlechtriem CH, Focken U, Becker K (2003) Effect of different lipid extraction methods on $\delta^{13}\text{C}$ of lipid and lipid-free fractions of fish and different fish feeds. *Isot Environ Health Stud* 39:135–140
- Sholto-Douglas AD, Field JG, James AG, Merwe NJ (1991) $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ isotope ratios in the Southern Benguela Ecosystem: indicators of food web relationships among different size-classes of plankton and pelagic fish; differences between fish muscle and bone collagen tissues. *Mar Ecol Prog Ser* 78:23–31
- Smedes F (1999) Determination of total lipid using non-chlorinated solvents. *Analyst* 124:1711–1718
- Tasker ML, Camphuysen CJ, Cooper J, Garthe S, Montevecchi WA, Blaber SJM (2000) The impacts of fishing on marine birds. *ICES J Mar Sci* 57:531–547

-
- Thompson DR, Phillips RA, Stewart FM, Waldron S (2000) Low $\delta^{13}\text{C}$ signatures in pelagic seabirds: lipid ingestion as a potential source of ^{13}C -depleted carbon in the Procellariiformes. *Mar Ecol Prog Ser* 208:265–271
- Vanderklift MA, Ponsard S (2003) Sources of variation in a consumer-diet $\delta^{15}\text{N}$ enrichment: a meta-analysis. *Oecologia* 136:169–182
- Votier SC, Furness RW, Bearhop S, Crane JE, Caldow RWG, Catry P, Ensor K, Hamer KC, Hudson AV, Kalmbach E, Klomp NI, Pfeiffer S, Phillips RA, Prieto I, Thompson DR (2004) Changes in fisheries discard rates and seabird communities. *Nature* 427:727–730
- Williams TD (2005) Mechanisms underlying the costs of egg production. *Bioscience* 55:39–48
- Williams CT, Buck CL, Sears J, Kitaysky AS (2007) Effects of nutritional restriction on nitrogen and carbon stable isotopes in growing seabirds. *Oecologia* 153:11–18