1 **Year-round reproduction in a seasonal sea:**
2 **Biological cycle of the introduced ascidian**
3 ***Styela plicata* in the Western Mediterranean**

4

5

6 M. Carmen Pineda¹, Susanna López-Legentil¹ and Xavier Turon^{2*}

7

8 ¹*Department of Animal Biology (Invertebrates), University of Barcelona, Diagonal*

9 *Avenue 643, 08028 Barcelona, Spain*

10 ²*Centre for Advanced Studies of Blanes (CEAB-CSIC), Accés Cala S. Francesc 14, 17300*

11 *Blanes, Girona, Spain*

12

13 ***Corresponding author:** Xavier Turon. E-mail: xturon@ceab.csic.es. Tel.: +34 972 336

14 101, fax: +34 972 337 806

15

16

Abstract

The widely introduced ascidian *Styela plicata* is very common in the Western Mediterranean, an area that can act as a source for secondary introductions due to its high shipping activity. In order to understand the potential of this species to colonize new habitats, its reproductive features were assessed in the Western Mediterranean by means of monthly monitoring of two populations (Vilanova i la Geltrú 41°12'53''N, 1°44'11''E; Blanes 41°40'29"N, 2°47'56"E) from January 2009 to December 2010. The reproductive activity of this species was assessed through gonad histology and a gonad index. Population size structure was measured monthly in order to study recruitment dynamics. No clear seasonal pattern was observed, and mature gametes and recruits were present all year long. Spawning was potentially continuous, although it seemed punctuated with pulses of gamete release, particularly in spring. A prolonged reproductive period is likely to confer a competitive advantage on *S. plicata* in temperate seas, where most species reproduce seasonally, and may promote recurrent introductions as larvae are available for settlement on transport vectors over much of the year.

Introduction

Countless marine species travel daily from their source of origin to new locations, either attached to ship's hulls or sea chests, in ballast waters, or being co-translocated with organisms associated with aquaculture activities (Allen 1953; Carlton and Geller 1993; Ruiz et al. 2000; Floerl and Inglis 2005; Blakeslee et al. 2010). Accordingly, introductions have increased notably during the last century, favored by maritime traffic and aquaculture (Carlton 1989; Vermeij 1996; Ruiz et al. 1997; Mack and D'Antonio 1998; Crooks 2002; Grosholz 2002). Although newly introduced species are often restricted to marginal habitats, such as harbors and aquaculture facilities, some may eventually spread to open habitats (Lambert 2002). Available information indicates that only one out of ten introduced species is able to survive and spread away from the introduced habitat, thus becoming invasive and causing serious alterations to native populations, communities and ecosystems (Williamson and Fitter 1996). Invasive species are considered, after habitat loss and fragmentation, the second most important cause of species extinction (Zibrowius 1991; Mack et al. 2000; Clavero and García-Berthou 2005).

Successful colonization of a new environment depends on the occurrence of adequate physical and biological conditions, both for adults and larvae (Blackburn and Duncan 2001; Stachowicz et al. 2002; Verween et al. 2007; Fowler et al. 2011; Zerebecki and Sorte 2011). Invasive species are often opportunistic, exploiting temporal windows of tolerable conditions to proliferate and occupy new habitats (McKinney 2002). Other characters that make introduced species prone to become invasive are adaptation to

disturbance (Hobbs 1992; Altman and Whitlatch 2007), wide environmental tolerances (McMahon 1996; Marchetti et al. 2004), the ability to overcome local control by resident species (Osman and Whitlatch 1998; Stachowicz et al. 2002), and high growth rates and reproductive output (McMahon 1996; Marchetti et al. 2004). Thus, in order to determine the invasive potential of a new introduced species and develop efficient management tools, it is necessary to acquire a better knowledge of their biological strategies and especially of their reproductive cycle (e.g., Grosholz and Ruiz 1996; Fine et al. 2001; Thornber et al. 2004).

Ascidians, or sea squirts, are conspicuous components of epibenthic marine communities all over the globe (e.g., Glasby 2001; Voultsiadou et al. 2007) and are among the most important marine invaders worldwide (Lambert 2002, 2007; Whitlatch and Bullard 2007). This group has the ability to severely modify the structure of coastal habitats by forming large aggregates that out-compete other organisms for resources (Zajac et al. 1989; Nandakumar et al. 1993; Lambert and Lambert 2003; Castilla et al. 2004; Agius 2007; Turon et al. 2007). *Styela plicata* is a solitary ascidian commonly found inhabiting harbors and salt marsh habitats of warm and temperate waters (i.e., Pacific, Indian, and Atlantic oceans, including the Mediterranean Sea), usually at high densities (Barros et al. 2009; David et al. 2010; Pineda et al. 2011, 2012). A recent genetic study has confirmed that this species has been present in these oceans for a long time, and that recurrent colonization events and shuffling among populations are determining its current genetic signature (Pineda et al. 2011). It is presently unclear where this species is endemic. Available evidence, albeit non conclusive, points to an origin in the NW Pacific (Barros et al 2009; Carlton 2009; Pineda et al. 2011). The

introduction success of *S. plicata* has been attributed to its high tolerance of polluted waters (Naranjo et al. 1996) and to moderately wide changes in temperature and salinity (Sims 1984; Thiagarajan and Qian 2003; Pineda et al. 2012). The high genetic variability reported in *S. plicata* (Pineda et al. 2011) may also enhance this species' ability to adapt to new environments (Sakai et al. 2001) and displace indigenous species (Rius et al. 2009b).

As is usual in solitary ascidians, *S. plicata* is hermaphroditic and oviparous, with development involving a non-feeding, short-lived larval stage (Svane and Young 1989). *Styela plicata* is reported to become sexually mature at about 40-mm body length (Tucker 1942; Yamaguchi 1975). The reproductive cycle or the settlement patterns of *S. plicata* have been determined for populations in Japan (Yamaguchi 1975) and the Eastern Mediterranean (Sabbadin 1957; Sciscioli et al. 1978; Tursi and Materrese 1981). These reports point to an arrest of reproductive activity during winter months and to either a prolonged settlement during the summer (June-September, Sciscioli et al. 1978), a restricted settlement period in summer (August-September, Tursi and Matarrese 1981), or a variable number of discrete generations per year (Sabbadin 1957; Yamaguchi 1975). These differences in the reproductive cycle could be due to a number of factors, including temperature, location and the genetic structure of the investigated populations. To date, no life-cycle data are available for populations in Western Mediterranean, where *S. plicata* is abundant in most harbors and marinas. Considering the high maritime traffic in some of those harbors (e.g., Barcelona, Alicante, Marseilles) and the existence of smaller marinas all along the coast, this area could act as a source for recurrent introductions in the Mediterranean and in other oceans, and thus deserves further investigation.

The goal of this study was to assess the reproductive features and population dynamics of the introduced ascidian *S. plicata* in the Western Mediterranean to assist in predicting the spreading potential of this species. To achieve this goal, the reproductive cycle was determined for two populations from the NE coast of Spain through examination of the gonad histology and calculation of a gonad index (GI) over a two-year period. We also took monthly measurements of the population size structure at one of the sites, in order to determine population dynamics and recruitment patterns. We hypothesized that *S. plicata* has a seasonal cycle of reproduction coupled to the strong seasonality of environmental parameters in the Mediterranean.

Material and methods

Study site, sampling and size-structure

The study was undertaken in two harbors of the NW Mediterranean coast: Vilanova i la Geltrú (41° 12' 53'' N, 1° 44' 11'' E), the larger and more polluted harbor, and Blanes (41° 40' 29"N, 2° 47' 56"E) located ca. 100 km NE from the former. Both harbors sustain a variety of marine-related activities including recreational boating and commercial fishing. From January 2009 to December 2010, ten adult *Styela plicata* (> 40 mm, Yamaguchi 1975) were collected monthly at each site from depths of 0–1 m by pulling up harbor ropes or removing individuals from submersed docks. The ascidians were immediately fixed in 4% formaldehyde and stored at room temperature until used for histology and morphometric analyses. Water temperature was automatically registered at

126 1-m depth every hour with HOBO® loggers (0.001°C precision), Onset Computer
127 Corporation, Massachusetts. The size-structure of this species was determined from
128 November 2009 to December 2010 in Vilanova i la Geltrú, the population where *S.*
129 *plicata* was more abundant. Every month, multiple ropes located in the same area of the
130 harbor were pulled from the water and the height of 150 randomly selected individuals
131 was measured directly on the ropes with a calliper. Clumps of individuals and ropes were
132 scrutinized to detect small recruits (<10 mm). After measurement, the ascidians were left
133 on the ropes and put back into the water. We used different ropes on sampling date.
134 Individual lengths recorded monthly were used to plot size frequency histograms using 2-
135 mm size classes.

136

137 Morphometric variables

138

139 Once in the laboratory, the individuals kept in 4% formaldehyde were carefully cleaned
140 to remove as many epibionts as possible from their tunics. For each individual, we
141 measured the height (maximal distance from base to tip) as a proxy for size using a
142 calliper. As we worked with fixed, contracted organisms, the length measured was an
143 underestimation of the lengths of live, relaxed individuals. However, the tough tunic in
144 this species only allows for slight contraction of the siphons, and we assume that
145 individuals contracted to a similar degree and, thus, that measurements were comparable.
146 The tunic was cut open to separate the mantle, and dissect it along the ventral side to
147 remove the branchial sac and to expose the gonads. Each individual possessed 2 to 11
148 gonads attached to the right side of the body wall, and 1 to 3 gonads to the left side. Each

gonad has a central, elongated ovary covered with testis follicles (Tucker 1942). From each specimen, a small piece (< 1 cm long) of one gonad from the right side was cut, weighed and kept in 4% formaldehyde for histology; the remaining gonadal tissue was dissected, weighed and placed in an oven at 60°C for 48 h to obtain its dry weight. To obtain the total dry weight (DW) of the gonads, we estimated the DW of the removed piece for histological purposes using the observed wet/dry weight ratio obtained for the other gonads. Wet and dry weights were also obtained for the tunic and mantle. A gonadosomatic index (hereafter referred to as gonad index, GI) was calculated as the gonad DW divided by the mantle DW.

The mean gonad index over time was correlated between sites and with temperature variations using monthly means and cross correlation analyses (Pearson coefficient). In these analyses, values of one variable were correlated with values of the other at different time lags (months). Correlation at time lag 0 corresponds to the usual Pearson correlation, positive lags correlate values in the first series to values in the second series that number of lags afterwards, and negative lags relate values in the first series to previous values in the second one. All analyses were performed using the software SYSTAT v. 12 (©SYSTAT Software, Inc. 2007), and SigmaStat v. 3.11 (©SYSTAT Software, Inc. 2004).

Histological analysis

Among the individuals used for morphometric analysis, each month, a piece of gonad from at least 5 individuals per population was dehydrated, embedded in paraffin,

sectioned, and stained. Methylene blue staining was used in preliminary sections to assess the gross morphology of the gonads and to guide further sectioning. For detailed histological observation, hematoxylin-eosin staining was used following standard procedures. The gonad sections were observed under a microscope equipped with a micrometer. The diameter of 100 oocytes sectioned through the nucleolus was measured per individual following Bingham (1997). In some individuals fewer than 100 oocytes could be found with nucleolus visible, but in all cases we measured a minimum of 200 oocytes mo^{-1} . In total, 7,504 oocytes were measured for individuals from Vilanova i la Geltrú, and 6,602 from Blanes. We calculated oocyte size-frequencies each month using 50- μm size classes as follows: $<50 \mu\text{m}$ (pre-vitellogenic), $50\text{--}100 \mu\text{m}$ and $100\text{--}150 \mu\text{m}$ (vitellogenic), and $>150 \mu\text{m}$ (mature, Sciscioli et al. 1978). For the testes, a categorical maturity index was established, according to the development of the male follicles (i.e. 1 = immature, only spermatogonia, 2 = mature sperm, 3 = spawning, empty spaces within the lumen).

Results

Population structure

The introduced ascidian *S. plicata* was present all year round in both harbors, Vilanova i la Geltrú and Blanes, on hanging ropes and virtually any hard substrate available. This species was often found forming aggregates of individuals attached to each other's tunics. During the study period, this species was more abundant and formed larger aggregates in

Vilanova i la Geltrú than in Blanes. Height was a good descriptor for size, as it showed a high correlation with total dry weight in the individuals collected (Pearson $r=0.735$, $p<0.001$).

The population at Vilanova i la Geltrú was characterized by the presence of individuals of all sizes most of the year. Heights ranged from a minimum of 2 mm in October to 68 mm in an individual collected in August. Based on occasional observations of settled individuals on newly exposed ropes, we considered as recruits those individuals that were < 15 mm high, which corresponded to an approximate age of one month (authors' pers. obs.). Recruits were especially abundant after mid-summer, and absent only in May. Adults (> 40 mm) were found all year round, although the largest sizes classes (> 50 mm) were almost absent in December 2009 and in February and March 2010 (Fig. 1). Thus, size-frequency plots suggested higher recruitment in fall and winter coupled with slower growth and loss of the oldest individuals (Fig. 1). The presence of juveniles and adults all year long masked any evidence of a cohort structure in the size-frequency histograms. Instead, the population structure seemed to result from overlapping, successive generations (Fig. 1).

Reproductive cycle

The mean gonad index (GI) and seawater temperature over the study period for Vilanova i la Geltrú and Blanes are shown in Figure 2. Seawater temperature showed a clear seasonal cycle at both sites, with the lowest values at the end of winter (10.2 and 12.4°C in February 2010 for Vilanova i la Geltrú and Blanes, respectively) and the highest in

September 2009 (29.1 and 26°C, respectively). Thus, temperatures in Blanes oscillated less (13.6°C from lowest to highest) than in Vilanova i la Geltrú (18.9°C). The mean GI did not show a clear seasonal pattern for either of the two populations, although a sharp decrease in the mean GI was observed in April 2009 in both (Fig. 2). Decreases in GI values were also observed in September 2009, and January, May, August and October 2010 for Vilanova i la Geltrú (Fig. 2a), and September and November 2009, and January to May, July and December 2010 in Blanes (Fig. 2b). A noticeable peak was also observed in January 2010 for the population in Blanes (Fig. 2b).

Cross correlation analyses were not significant for most time lags. Nevertheless, there was a clear wave-like pattern of positive correlations between gonad index and temperature in the previous months for both populations, while correlations were negative between GI and the temperature measured in subsequent months (Figs. 3a, 3b). This positive correlation indicated that GI tended to increase some months (peak of correlation at 2–3 mo) after temperature increases. Correlations at time lag 0 were small and negative. Mean GIs followed similar patterns in both localities, as indicated by a significant correlation coefficient (cross-correlation analysis at time lag 0, Fig 3c). The relationship was also positive and significant between the mean GI in Vilanova i la Geltrú and the values in Blanes the two subsequent months (time lags +1 and +2, Fig. 3c).

Gonad histology showed, in cross section, the presence of a central core of female follicles with oocytes surrounded by peripheral male follicles (Fig. 4a). Mature oocytes had test cells inside the chorion and were surrounded by two layers of follicle cells (Fig 4b). Male follicles were characterized by a thick wall of germinative epithelium and a lumen occupied by developing spermatozoa (Fig 4a,c,d) or by empty spaces after

spawning (Fig. 4e). We found considerable variability in the maturation state of the male gonads within the same month and population, while mature oocytes were always present in at least some individuals. These observations confirmed the year-round reproductive cycle previously found with the GI (Fig. 5). A sharp decrease in the mean oocyte diameter was also observed in spring 2009 and 2010 in both populations (Fig. 5). Minimum values for the mean oocyte diameter were also recorded in August 2009 and December 2010 for the Vilanova i la Geltrú population, while for the Blanes population minima were recorded in July and September 2009, and January, April, August and December 2010 (Fig. 5). Mature male follicles and follicles with partially empty lumens, due to the release of sperm, were also found over most of the year except for some of the coolest months (January 2009 in Vilanova i la Geltrú and December 2010 in Blanes), when gonads had mostly immature male follicles (Fig. 5).

The oocyte size-frequencies showed that pre-vitellogenic oocytes ($< 50 \mu\text{m}$) were present all year round, but increased in proportion concomitantly with the decreases in mean oocyte diameter (Fig. 6). Mature oocytes ($>150 \mu\text{m}$) were also present all year, but were especially abundant during winter. In the individuals collected from Vilanova i la Geltrú, the proportion of mature oocytes peaked in winter-early spring followed by a sharp decrease in April (Fig. 6a). In Blanes, the proportion of mature oocytes was more variable, with additional minimum values in May, June, and September 2009 and April, August and December 2010 (Fig. 6b).

Discussion

The introduced ascidian *Styela plicata* was present year round in the two studied populations from the Western Mediterranean. Individuals were 2 mm–68 mm in height. There was no clear seasonal cycle for reproduction in Vilanova i la Geltrú or Blanes in 2009–2010. In fact, this species had mature oocytes and male follicles all year round. The mean gonad index and the mean oocyte diameter showed marked decreases in spring 2009 and, to a lesser degree, in spring 2010. Gonad index decreases alone may not correspond to spawning events, as shown by a lack of correlation between GI and oocyte diameter in some ascidian species (e.g., Bingham et al. 1997; Wong et al. 2011). Mobilization of reserves stored in the gonads in periods of shortage, as noted also in other groups (e.g., echinoderms King et al. 1994; Lozano et al. 1995), can occur in ascidians (Bingham et al. 1997). However, the coincidence of several parameters (GI drop, oocyte diameter drop, and minima in proportion of mature oocytes) suggests that pulses of spawning occur in spring in *S. plicata*. Less confidence can be placed on secondary peaks of GI as indicative of spawning, as they are often not coincident with changes in oocyte diameter. A combination of several approaches (histology and gonad indices) seems the best way to obtain a complete picture of the reproductive cycle in ascidians (Becerro and Turon 1992; Sahade et al. 2004). In our case, results indicate that *S. plicata* reproduces throughout the year in the Western Mediterranean, although more intensely so in spring. This pattern of pulses of spawning against a background of continuous reproduction has been described in other temperate ascidians (Svane and Lundälv 1981, 1982; Bingham et al. 1997).

Prolonged reproductive activity of *S. plicata* from spring to autumn was previously reported in the Eastern Mediterranean (Sabbadin 1957; Sciscioli et al. 1978).

287 These authors considered that this species did not actively reproduce during the coldest
288 months, as no recruitment was detected in winter on artificial panels (Sciscioli et al.
289 1978; Tursi and Matarrese 1981). Sabbadin (1957) set the low temperature threshold for
290 *S. plicata* reproduction at 10°C, a seawater temperature that was easily reached during the
291 winter months in the Lagoon of Venice (Italy). Below 10°C this species not only was
292 unable to reproduce but could also disappear (Sabbadin 1957). Such low temperatures are
293 seldom reached in the Western Mediterranean (Margalef 1985; Coma et al. 2000) and,
294 although there was a short period in which male follicles were not mature in the coolest
295 months, mature individuals with large oocytes were found in winter in both of our study
296 populations. Moreover, small recruits were observed in Vilanova i la Geltrú during the
297 winter months. Thus, taken together, our results indicate that *S. plicata* is also
298 reproductively active in the winter months in the Western Mediterranean. These results
299 are in agreement with Panagiotou et al. (2007), who reported the presence of *S. plicata*
300 recruits all year in Thessaloniki Bay (Greece). In Tokyo Bay (Japan), where the
301 temperature regime is more similar to the Western Mediterranean, Yamaguchi (1975)
302 observed that individuals of *S. plicata* were ripe in winter, although they did not spawn
303 during the coldest months of the year. Finally, previous studies have reported the
304 existence of several generations per year for this species in central Japan (Yamaguchi
305 1975), and Eastern Mediterranean (Sabbadin 1957). Our results did not show any clear-
306 cut succession of generations for populations in Western Mediterranean harbors. Rather,
307 individuals of a wide range of sizes were present all year round, suggesting that there
308 were several overlapping cohorts of recruits each year. Interannual differences in the
309 number and intensity of GI peaks and mean oocyte diameter changes were also observed

in the two monitored harbors. Temperature has been suggested as the main factor triggering not only ascidian reproduction (e.g., Millar 1971; Turon 1988) but also the reproductive cycles of many marine invertebrates in temperate seas (Orton 1920). Accordingly, temperature differences between years and sites could explain the similarities and differences found. For instance, there was an unusual and sharp increase in seawater temperature in February 2009 (6°C). This anomaly could in turn explain the spawning event observed for both populations two months later (April 2009). Similarly, the smaller spawning episode recorded in spring 2010 could be due to a later and more gradual increase in seawater temperature (March).

In the Mediterranean, most marine invertebrates present seasonal biological cycles (Coma et al. 2000), including colonial (Turon 1988; Caralt et al. 2002; López-Legentil et al. 2005) and solitary ascidians (Becerro and Turon 1992; Panagiotou et al. 2008; Vafidis et al. 2008; Rius et al. 2009a). Summer is an unfavorable season for many sessile invertebrates in the Mediterranean, due to food shortage (Coma et al. 2000) and high densities of macroalgae competing for space (Ballesteros 1989). Therefore, the continuous presence of this species and the ability to reproduce all year round may confer a competitive advantage on *S. plicata* compared to seasonally reproducing invertebrate species. Extended reproductive cycles and fast growth to maturity have been reported for several invasive ascidians (Bourque et al. 2007; Shenkar and Loya 2008; Wong et al. 2008). Thus, based on our current results, this species could become a threat to local biota if it spreads to natural habitats. However, although the species has been found outside harbors, to date its abundance has always been low and it never monopolizes the substrate as it does inside harbors, marinas or on artificial structures (authors' pers. obs.).

Other factors controlling the spread of the species to natural substrates, such as predation (Sutherland 1974), competition, or the effects of hydrology, should be investigated.

Besides conferring a competitive advantage, a continual reproductive period could also allow *S. plicata* to exploit temporal windows of favorable conditions. It has been demonstrated that adults of this species can respond to changes in temperature and salinity by increasing the production of stress proteins (Pineda et al. 2012). However, the embryos and larvae of *S. plicata*, although relatively resistant to pollutants such as copper, are very sensitive to changes in temperature or salinity (Pineda 2012). Based on these observations, it is likely that some reproductive episodes do not result in local recruitment and spreading them over time would increase the probability of some larval recruitment. In addition, the existence of multiple batches of larvae over the year ensures a plentiful supply of juveniles on ships' hulls, ready to spread to other marinas, and facilitating recurrent introduction of the species. This fact, coupled with a large genetic pool (coherent with the genetic structure described for this species, Pineda et al. 2011) would ensure the persistence of *S. plicata* populations living under harsh conditions such as those usually encountered in enclosed man-made habitats.

Acknowledgements

We are grateful to E. Arias, C. Torres, A. Pineda and L. Gelabert for assistance in the field. A. Muntadas and R. Espluga helped in the laboratory. Dr. M. Durfort provided useful comments on the histology. Dr. A. Tursi kindly provided some essential bibliography. Dr. P. Erwin reviewed the manuscript for English grammar. "Ports de la

Generalitat” granted collecting permits and access to the harbors. This research was supported by the Marie Curie International Reintegration Grant FP7-PEOPLE-2010-RG 277038 within the 7th European Community Framework Program, by the Spanish Government projects CTM2010-22218 and CTM2010-17755, the Catalan Government grant 2009SGR-484 for Consolidated Research Groups, and by a University of Barcelona APIF fellowship to MCP.

References

- Agius BP (2007) Spatial and temporal effects of pre-seeding plates with invasive ascidians: Growth, recruitment and community composition. *J Exp Mar Biol Ecol* 342:30–39
- Allen F (1953) Distribution of marine invertebrates by ships. *Aust J Mar Freshw Res* 4:307–316
- Altman S, Whitlatch RB (2007) Effects of small-scale disturbance on invasion success in marine communities. *J Exp Mar Biol Ecol* 342:15–29
- Ballesteros E (1989) Production of seaweeds in Northwestern Mediterranean marine communities: Its relation with environmental factors. *Sci Mar* 53:357–364
- Barros R, Rocha R, Pie M (2009) Human-mediated global dispersion of *Styela plicata* (Tunicata, Ascidiacea). *Aquatic Inv* 4: 45–57
- Becerro MA, Turon X (1992) Reproductive cycles of the ascidians *Microcosmus sabatieri* and *Halocynthia papillosa* in the Northwestern Mediterranean. *Mar Ecol* 13:363–373
- Bingham BL (1997) Light cycles and gametogenesis in three temperate ascidian species. *Invertebr Biol* 116:61–70
- Blackburn TM, Duncan RP (2001) Determinants of establishment success in introduced birds. *Nature* 414:195–197
- Blakeslee AMH, McKenzie CH, Darling JA, Byers JE, Pringle JM, Roman J (2010) A hitchhiker's guide to the Maritimes: anthropogenic transport facilitates long-distance dispersal of an invasive marine crab to Newfoundland. *Divers Distrib* 16:879–891

386 Bourque D, Davidson J, MacNair NG, Arsenault G, LeBlanc AR, Landry T, Miron G
387 (2007) Reproduction and early life history of an invasive ascidian *Styela clava* Herdman
388 in Prince Edward Island, Canada. J Exp Mar Biol Ecol 342:78–84

389 Caralt S, López-Legentil S, Tarjuelo I, Uriz MJ, Turon X (2002) Contrasting biological
390 traits of *Clavelina lepadiformis* (Ascidiacea) populations from inside and outside
391 harbours in the Western Mediterranean. Mar Ecol-Prog Ser 244:125–137

392 Carlton JT (1989) Mans role in changing the face of the Ocean - Biological invasions and
393 implications for conservation of near-shore environments. Conserv Biol 3:265–273

394 Carlton JT, Geller JB (1993) Ecological roulette - the global transport of nonindigenous
395 marine organisms. Science 261:78–82

396 Carlton JT (2009) Deep invasion ecology and the assembly of communities in
397 historical time. In: Rilov G, Crooks JA (eds) Biological invasions in marine ecosystems.
398 Springer, Berlin, pp 13–56

399 Castilla JC, Guinez R, Caro AU, Ortiz V (2004) Invasion of a rocky intertidal shore by
400 the tunicate *Pyura praeputialis* in the Bay of Antofagasta, Chile. Proc Natl Acad Sci
401 USA 101:8517–8524

402 Clavero M, Garcia-Berthou E (2005) Invasive species are a leading cause of animal
403 extinctions. Trends Ecol Evol 20:110

404 Coma R, Ribes M, Gili JM, Zabala M (2000) Seasonality in coastal benthic ecosystems.
405 Trends Ecol Evol 15:448–453

406 Crooks JA (2002) Characterizing ecosystem-level consequences of biological invasions:
407 the role of ecosystem engineers. Oikos 97:153–166

408 David GK, Marshall DJ, Riginos C (2010) Latitudinal variability in spatial genetic
409 structure in the invasive ascidian, *Styela plicata*. Mar Biol 157: 1955–1965

410 Fine M, Zibrowius H, Loya Y (2001) *Oculina patagonica*: a non-lessepsian scleractinian
411 coral invading the Mediterranean Sea. Mar Biol 138:1195–1203

412 Floerl O, Inglis GJ (2005) Starting the invasion pathway: the interaction between source
413 populations and human transport vectors. Biol Invasions 7:589–606

414 Fowler AE, Gerner NV, Sewell MA (2011) Temperature and salinity tolerances of Stage
415 1 zoeae predict possible range expansion of an introduced portunid crab, *Charybdis*
416 *japonica*, in New Zealand. Biol Invasions 13:691–699

417 Glasby TM (2001) Development of sessile marine assemblages on fixed versus moving
418 substrata. Mar Ecol Prog Ser 215:37–47

- 419 Grosholz E (2002) Ecological and evolutionary consequences of coastal invasions.
420 Trends Ecol Evol 17:22–27
- 421 Grosholz ED, Ruiz GM (1996) Predicting the impact of introduced marine species:
422 Lessons from the multiple invasions of the European green crab *Carcinus maenas*. Biol
423 Conserv 78:59–66
- 424 Hobbs RJ, Huenneke LF (1992) Disturbance, diversity, and invasion - Implications for
425 conservation. Conserv Biol 6:324–337
- 426 King CK, Hoegh-Guldberg O, Byrne M (1994) Reproductive cycle of *Centrostephanus*
427 *rodgersii* (Echinoidea), with recommendations for the establishment of a sea urchin
428 fishery in New South Wales. Mar Biol 120:95–106
- 429 Lambert CC, Lambert G (2003) Persistence and differential distribution of nonindigenous
430 ascidians in harbors of the Southern California Bight. Mar Ecol Prog Ser 259:145–161
- 431 Lambert G (2002) Nonindigenous ascidians in tropical waters. Pac Sci 56:291–298
- 432 Lambert G (2007) Invasive sea squirts: A growing global problem. J Exp Mar Biol Ecol
433 342:3–4
- 434 López-Legentil S, Ruchty M, Domenech A, Turon X (2005) Life cycles and growth rates
435 of two morphotypes of *Cystodytes* (Ascidacea) in the Western Mediterranean. Mar Ecol
436 Prog Ser 296:219–228
- 437 Lozano J, Galera J, López S, Turon X, Palacín C, Morera G (1995) Biological cycles and
438 recruitment of *Paracentrotus lividus* (Echinodermata: Echinoidea) in two contrasting
439 habitats. Mar Ecol-Prog Ser 122:179–191
- 440 Mack MC, D'Antonio CM (1998) Impacts of biological invasions on disturbance
441 regimes. Trends Ecol Evol 13:195–198
- 442 Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzaz FA (2000) Biotic
443 invasions: causes, epidemiology, global consequences, and control. Ecol Appl
444 10:689–710
- 445 Marchetti MP, Moyle PB, Levine R (2004) Alien fishes in California watersheds:
446 Characteristics of successful and failed invaders. Ecol Appl 14:587–596
- 447 Margalef R (1985) Key environments: Western Mediterranean. Pergamon Press, New
448 York
- 449 McKinney ML (2002) Urbanization, biodiversity, and conservation. BioScience
450 52:883–890
- 451 McMahon RF (1996) The physiological ecology of the zebra mussel, *Dreissena*
452 *polymorpha*, in North America and Europe. Am Zool 36:339–363

453 Nandakumar K, Tanaka M, Kikuchi T (1993) Interspecific competition among fouling
454 organisms in Tomioka Bay, Japan. *Mar Ecol-Prog Ser* 94:43–50

455 Naranjo SA, Carballo JL, García-Gómez JC (1996) Effects of environmental stress on
456 ascidian populations in Algeciras Bay (southern Spain) Possible marine bioindicators?
457 *Mar Ecol-Prog Ser* 144:119–131

458 Orton JH (1920) Sea-temperature, breeding and distribution in marine animals. *J Mar*
459 *Biol Assoc UK* 12:339–366

460 Osman RW, Whitlatch RB (1998) Local control of recruitment in an epifaunal
461 community and the consequences to colonization processes. *Hydrobiologia*
462 375/376:113–123

463 Panagiotou M, Antoniadou C, Chintiroglou C (2008) Population dynamics and
464 reproductive status of *Microcosmus savignyi* Monniot, 1962 (Thermaikos Gulf, Eastern
465 Mediterranean): a preliminary assessment. *J Nat Hist* 42:545–558

466 Panagiotou M, Antoniadou C, Krestenitis Y, Chintiroglou C (2007) Stock assessment of
467 the dominant ascidians: *Microcosms savignyi*, *Styela plicata* and *Phallusia mammillata*,
468 in Thessaloniki bay (Thermaikos Gulf). *Fresen Environ Bull* 16:1012–1019

469 Pineda, MC (2012) A global wanderer: Biology, phylogeography and resilience of the
470 introduced ascidian *Styela plicata*. PhD thesis, Universitat de Barcelona, Barcelona

471 Pineda MC, López-Legentil S, Turon X (2011) The whereabouts of an ancient wanderer:
472 Global phylogeography of the solitary ascidian *Styela plicata*. *PLoS One* 6:e25495

473 Pineda MC, López-Legentil S, Turon X (2012) Stress levels over time in the introduced
474 ascidian *Styela plicata*: The effects of temperature and salinity variations on *hsp70* gene
475 expression. *Cell Stress Chaperon* 17: 435–444

476 Reynolds ES (1963) The use of lead citrate at high pH as an electron-opaque stain in
477 electron microscopy. *J Cell Biol* 17: 208–212

478 Rius M, Pineda MC, Turon X (2009a) Population dynamics and life cycle of the
479 introduced ascidian *Microcosmus squamiger* in the Mediterranean Sea. *Biol Invasions*
480 11:2181–2194

481 Rius M, Turon X, Marshall DJ (2009b) Non-lethal effects of an invasive species in the
482 marine environment: the importance of early life-history stages. *Oecologia* 159:873–882

483 Ruiz GM, Carlton JT, Grosholz ED, Hines AH (1997) Global invasions of marine and
484 estuarine habitats by non-indigenous species: Mechanisms, extent, and consequences.
485 *Am Zool* 37:621–632

- 486 Ruiz GM, Fofonoff PW, Carlton JT, Wonham MJ, Hines AH (2000) Invasion of coastal
487 marine communities in North America: Apparent patterns, processes, and biases. *Annu*
488 *Rev Ecol Syst* 31:481–531
- 489 Sabbadin A (1957) Il ciclo biologico di *Ciona intestinalis* (L.), *Molgula manhattensis* (de
490 Kay) e *Styela plicata* (Lesueur) nella laguna veneta. *Arch Oceanog Limnol* XI:1–28
- 491 Sahade R, Tatian M, Esnal GB (2004) Reproductive ecology of the ascidian
492 *Cnemidocarpa verrucosa* at Potter Cove, South Shetland Islands, Antarctica. *Mar Ecol-*
493 *Prog Ser* 272:131–140
- 494 Sakai AK, Allendorf FW, Holt JS, Lodge DM, Moilofsky J et al. (2001) The population
495 biology of invasive species. *Annu Rev Ecol Syst* 32:305–332
- 496 Sciscioli M, Lepaore E, Tursi A (1978) Relationship between *Styela plicata* (Les.)
497 (Tunicata) settlement and spawning. *Memorie di Biologia Marina e di Oceanografia*
498 8:65–75
- 499 Shenkar N, Loya Y (2008) The solitary ascidian *Herdmania momus*: native (Red Sea)
500 versus non-indigenous (Mediterranean) populations. *Biol Invasions* 10:1431–1439
- 501 Sims LL (1984) Osmoregulatory capabilities of 3 macrosympatric Stolidobranch
502 ascidians, *Styela clava* Herdman, *Styela plicata* (Lesueur), and *Styela montereyensis*
503 (Dall). *J Exp Mar Biol Ecol* 82:117–129
- 504 Stachowicz JJ, Fried H, Osman RW, Whitlatch RB (2002) Biodiversity, invasion
505 resistance, and marine ecosystem function: Reconciling pattern and process. *Ecology*
506 83:2575–2590
- 507 Sutherland JP (1974) Multiple stable points in natural communities. *Am Nat*
508 108:859–873
- 509 Svane I, Lundälv T (1981) Reproductive patterns and population dynamics of *Ascidia*
510 *mentula* O.F. Müller on the Swedish west coast. *J Exp Mar Biol Ecol* 50: 163–182
- 511 Svane I, Lundälv T (1982) Population dynamics and reproductive patterns of *Boltenia*
512 *echinata* (Acidiacea) on the Swedish west coast. *Neth J Sea Res* 16: 105–118
- 513 Svane I, Young C (1989) The biology and behaviour of ascidian larvae. *Oceanogr Mar*
514 *Biol Annu Rev* 27:45–90
- 515 Thiagarajan V, Qian PY (2003) Effect of temperature, salinity and delayed attachment
516 on development of the solitary ascidian *Styela plicata* (Lesueur). *J Exp Mar Biol Ecol*
517 290:133–146
- 518 Thornber CS, Kinlan BP, Graham MH, Stachowicz JJ (2004) Population ecology of the
519 invasive kelp *Undaria pinnatifida* in California: environmental and biological controls on
520 demography. *Mar Ecol-Prog Ser* 268:69–80

- 521 Tucker GH (1942) The histology of the gonads and development of the egg envelopes of
522 an ascidian (*Styela plicata* Lesueur). J Morphol 70:81–113
- 523 Turon X (1988) The ascidians of Tossa de Mar (NE Spain). II.-Biological cycles of the
524 colonial species. Cah Biol Mar 29:407–418
- 525 Turon X, Nishikawa T, Rius M (2007) Spread of *Microcosmus squamiger* (Ascidacea :
526 Pyuridae) in the Mediterranean Sea and adjacent waters. J Exp Mar Biol Ecol
527 342:185–188
- 528 Tursi A, Matarrese A (1981) Phenomena of settling in *Styela plicata* (Les.) (Tunicata).
529 Mem Biol Mar Oceanogr 11:117–130
- 530 Vafidis D, Antoniadou C, Chintiroglou C (2008) Population dynamics, allometric
531 relationships and reproductive status of *Microcosmus sabatieri* (Tunicata : Ascidacea) in
532 the Aegean Sea. J Mar Biol Assoc UK 88:1043–1051
- 533 Vermeij GJ (1996) An agenda for invasion biology. Biol Conserv 78:3–9
- 534 Verween A, Vincx M, Degraer S (2007) The effect of temperature and salinity on the
535 survival of *Mytilopsis leucophaeata* larvae (Mollusca, Bivalvia): The search for
536 environmental limits. J Exp Mar Biol Ecol 348:111–120
- 537 Voultsiadou E, Pyrounaki MM, Chintiroglou C (2007) The habitat engineering tunicate
538 *Microcosmus sabatieri* Roule, 1885 and its associated peracarid epifauna. Estuar Coast
539 Shelf Sci 74:197–204
- 540 Whitlatch RB, Bullard SG (2007) Introduction to the Proceedings of the 1st International
541 Invasive Sea Squirt Conference. J Exp Mar Biol Ecol 342:1–2
- 542 Williamson M, Fitter A (1996) The varying success of invaders. Ecology 77:1661–1666
- 543 Wong NA, McClary D, Sewell MA (2011) The reproductive ecology of the invasive
544 ascidian, *Styela clava*, in Auckland Harbour, New Zealand. Mar Biol 158:2775–2785
- 545 Yamaguchi M (1975) Growth and reproductive cycles of marine fouling Ascidians *Ciona*
546 *intestinalis*, *Styela plicata*, *Botrylloides violaceus*, and *Leptoclinum mitsukurii* at
547 Aburatsubo-Moroiso Inlet (Central Japan). Mar Biol 29:253–259
- 548 Zajac RN, Whitlatch RB, Osman RW (1989) Effects of inter-specific density and food-
549 supply on survivorship and growth of newly settled benthos. Mar Ecol-Prog Ser
550 56:127–132
- 551 Zerebecki RA, Sorte CJ (2011) Temperature tolerance and stress proteins as mechanisms
552 of invasive species success. PLoS One 6:e14806
- 553 Zibrowius H (1991) Ongoing modification of the Mediterranean marine fauna and flora
554 by the establishment of exotic species. Mésogée 51:83–107
555

Figure Legends

Fig. 1 Size-frequency histograms (height, mm) of *S. plicata* for each sampled month in 2009–2010 in Vilanova i la Geltrú

Fig. 2 Mean Gonad Index ($\pm$ SE) and seawater temperature ($^{\circ}$ C) in 2009–2010 in a) Vilanova i la Geltrú and b) Blanes

Fig. 3 Cross-correlation analyses of *S. plicata* a) Gonad Index vs. temperature in Vilanova i la Geltrú, b) Gonad Index vs. temperature in Blanes, c) Gonad Index in Vilanova i la Geltrú vs. Gonad Index in Blanes. Curved lines represent 95% confidence intervals of correlation coefficient

Fig. 4 Light micrographs of **a)** a gonad histological section stained with haematoxylin-eosin (MMF: mature male follicle, DO: developing oocyte, MO: mature oocyte. **b)** female gonad stained with methylene blue (TC: test cells, FC: follicle cells, N: nucleolus). **c–e)** male follicles stained with haematoxylin-eosine (c: immature, d: mature, e: mature and spawning). Scale bars = 200 μ m (**a**), 50 μ m (**b**), 100 μ m (**c–e**)

Fig. 5 Mean oocyte diameter ($\pm$ SE) in *S. plicata* and seawater temperature in a) Vilanova i la Geltrú and b) Blanes. Horizontal bars show male maturation state over study period

579 **Fig. 6** Percentage of each oocyte size-class (previtellogenic, 0-50 μm ; mature, >150 μm)
580 in *S. plicata* in 2009–2010 in a) Vilanova i la Geltrú and b) Blanes
581

Figure1
[Click here to download high resolution image](#)

Figure2

[Click here to download high resolution image](#)

Figure3

[Click here to download high resolution image](#)

Figure4
[Click here to download high resolution image](#)

Figure5

[Click here to download high resolution image](#)

Figure6

[Click here to download high resolution image](#)