**Bioactive compounds, myrosinase activity and antioxidant capacity of
white cabbages grown in different locations of Spain**

Elena Peñas, Juana Frias, Cristina Martínez-Villaluenga and Concepción
Vidal-Valverde*

*Institute of Food Science, Technology and Nutrition (ICTAN-CSIC), Juan
de la Cierva 3, 28006 Madrid, Spain*

*Corresponding author

Telephone number: +34 91 5622900

Fax number: +34 91 5644853

E-mail: cvidal@ifi.csic.es

ABSTRACT

The influence of two Spanish growing locations with well differentiated climatic conditions (North and East areas) on the main bioactive compounds, glucosinolates (GLS), total phenolic compounds (TPC), vitamin C, as well as myrosinase activity and antioxidant capacity in five white cabbage (*Brassica oleracea* L. var. *capitata*) cultivars were investigated. Cabbages with the highest concentration of total GLS presented the highest vitamin C level ($r=0.75$, $P\leq 0.05$) and the lowest antioxidant capacity ($r=-0.76$, $P\leq 0.05$). The cultivars with the highest vitamin C content had the lowest myrosinase activity ($r=-0.89$, $P\leq 0.05$) and antioxidant capacity ($r=-0.86$, $P\leq 0.05$), while those with the largest TPC amount showed the highest antioxidant capacity ($r=0.71$, $P\leq 0.05$). Cabbage cultivars grown in North area of Spain with low temperatures and radiation led to higher mean values of myrosinase activity (29.25 U/g d.m.), TPC (10.0 GAE mg/g d.m.) and antioxidant capacity (81.6 μmol Trolox/g d.m.), while cultivars grown in the East area with high temperature and radiation led to larger mean values of GLS (14.3 μmol /g d.m.) and vitamin C (5.3 mg/g d.m.). The results of this investigation provide information regarding the most suitable Spanish growing location to produce white cabbage with an optimized content of health promoting compounds.

Keywords: White cabbage, glucosinolates, myrosinase, vitamin C, total phenolics, antioxidant capacity, growing location, genotype

INTRODUCTION

Epidemiological studies supported by extensive research in human volunteers, animal models and cell culture systems have established the protective role of *Brassica* vegetables in several types of cancer.¹⁻³ This protective effect is associated with the health-promoting phytochemical content in *Brassica* vegetables, which includes glucosinolates (GLS) and their breakdown products (isothiocyanates -ITC- and indoles) as well as antioxidants such as vitamin C and phenolic compounds.^{4,5}

GLS are β -thioglycoside N-hydroxysulfates that upon degradation by either myrosinase (β -thioglucoside glucohydrolase) within the plant or by enzymatic decomposition within the gastrointestinal tract yield biologically active hydrolysed products.⁵ There is a large body of evidence showing the chemopreventive action of GLS hydrolysis products by modulating detoxification enzymes⁶, which protects from DNA damage and proliferation of cancer cells^{1,2}. Ascorbigen is the major GLS breakdown product found in fermented white cabbage (sauerkraut) which results from the hydrolysis of glucobrassicin by myrosinase enzyme and the further reaction with L-ascorbic acid at low pH. The anticarcinogenic, antioxidant and free radical scavenging properties of ascorbigen have been reviewed recently.⁷

Vitamin C and phenolic compounds are potent antioxidants which may exert their action directly by scavenging free radical species, by metabolizing peroxides to non-radical products and by chelating metal ions to prevent generation of oxidizing species.⁸⁻¹⁰ In addition, some phenolic compounds inhibit pro-oxidant enzymes and modulate proinflammatory gene expression, enzyme activities and proinflammatory molecules such as cytokines,

prostaglandins and reactive oxygen species, thus they could contribute to prevention of cancer and cardiovascular diseases.¹¹

Cultivar, location and growing conditions play important roles in the production of bioactive compounds in *Brassica* vegetables.¹² The concentration and composition of GLS, phenolics and vitamin C in *Brassica* vegetables is genotype dependent.^{13,14} Moreover, climatic factors such as temperature, irradiation and water supply also have an important influence on the phytochemical content in *Brassica* vegetables.^{15,16} GLS breakdown product levels are due to the combination of GLS content in the plant and myrosinase activity.¹⁷ The activity of this enzyme depends also on the genetic variation¹⁸ and on some intrinsic (metal ions, ascorbic acid, pH) and extrinsic (temperature) factors.^{17,19} Therefore, cultivar selection should be tailored to specific environmental factors at each location to achieve the optimization in phytochemical content of *Brassica* vegetables. In addition, selected white cabbages with an optimized bioactive compound content could be used as raw material for sauerkraut production, enhancing the human dietary intake in health promoting compounds.

No information has been found about the effect of climatic growing conditions in different Spanish locations on the content of health promoting compounds of diverse white cabbage cultivars. Therefore, the objective of the present work was to determine the content of individual and total glucosinolates (GLS), myrosinase activity, vitamin C, total phenolics compounds (TPC), and antioxidant capacity (ORAC) in five white cabbage (*Brassica oleracea* L. var. *capitata*) cultivars grown in the North and East area of Spain.

MATERIALS AND METHODS

Plant material. White cabbage (*Brassica oleracea* L. var. *capitata*) cv.

Hinova, cv. Megaton, cv. Alfredo, cv. Candela and cv. Bronco were provided by Bejo Iberica S. L. (Madrid, Spain), planted in September 2008 and harvested in December 2008 in the fields of two different geographical Spanish locations: the North (Calahorra, La Rioja) and the East (Alboraya, Valencia) areas of the country. Three cabbage heads for each cultivar were randomly selected from the field at each location. Cabbage heads were trimmed of their outer leaves and their central cores were removed. White cabbages were frozen upon reception in liquid nitrogen, freeze-dried, milled and stored at -20°C for further analysis.

Individual and total glucosinolate (GLS) content. Total GLS were

extracted from freeze-dried samples followed by an enzymatic desulfatation, according to the method reported in the Official Journal of European Communities.²⁰ Briefly, 200 mg of freeze-dried samples were extracted twice in 3 mL of 70% (v/v) methanol at 70 °C and held at that temperature for 2 min. Samples were homogenised for 1 min using an Ultra-Turrax T25 Digital (IKA-Werke GmbH & Co, Staufen, Germany) at 20,000 rpm and centrifuged at 1089 x g for 10 min at 5 °C. Desulfatation of GLS and sinigrin (Sigma-Aldrich, Steinheim, Germany) used as standard were carried out in Sephadex A25 columns (Sigma-Aldrich) using 75 µL of purified sulfatase from *Helix pomatia* Type H-1 (E.C 2312.772.1 10kU/g solid from Sigma-Aldrich) purified according to the Official Journal of European Communities.²⁰ For separation of desulfo-

GLS, an Alliance Waters 2695 HPLC (Waters, Milford, USA) with a photodiode array detector and a Spherisorb ODS2 column (150 x 4.6 mm, 3 μ m) from Waters was used. Separation, detection and identification were performed as described elsewhere.¹⁵ GLS content was quantified by using desulfated sinigrin as external standard (concentration range 0-0.25 mM) and response factors relative to desulfated sinigrin.²⁰ Samples were independently analyzed in triplicate and results were reported as μ mol GLS per gram of dry matter (μ mol/g d.m.).

Myrosinase activity. Myrosinase activity was determined as described by Travers-Martin et al.²¹ with some modifications as it is described below. Crude extracts were prepared by homogenizing 350 mg of a freeze-dried sample in 10 mL Tris-EDTA buffer (200 mM Tris, 10 mM EDTA, pH 5.5) for 1 min in an ice bath using an Ultra-Turrax homogenizer T25 Digital. The homogenate was centrifuged at 23,708 x g for 15 min at 4 °C. To remove endogeneous glucosinolates and glucose, the crude extract was applied onto an Amicon ultrafiltration cell (Millipore, Billireca, MA, USA) of 10,000 Da molecular weight cut off and washed several times at 4 °C using Tris-EDTA buffer pH 5.5. The myrosinase activity was measured as the release of glucose using the GOD-PAP method.²² Briefly, 100 μ L of purified extracts and 25 μ L of 2 mM sinigrin in 0.2 M phosphate buffer pH 6.5 as substrate were mixed with 50 μ L freshly prepared color reagent (pH 7.0) containing 57 U/mL glucose oxidase from *Aspergillus niger* (E.C. 1.1.3.4., Sigma-Aldrich), 5.6 U/mL horseradish peroxidase (E.C. 1.11.1.7, Sigma-Aldrich), 2.8 mM 4-aminoantipyrine (Sigma-Aldrich), 30.7 mM phenol (Sigma-Aldrich) and 0.136 M

imidazole (Sigma-Aldrich). The release of glucose was determined by measuring the absorbance of the colored product N-(4-antipyril)-p-benzoquinone imine at 492 nm at room temperature in a 96-well clear-bottomed polystyrene plate (Sterilin, London, UK) using a microplate reader (Biotek Instruments, Winooksi, USA). Absorbance was read every minute for 60 min and plate was shaken between measurements. A linear part of at least 25 time points of the reaction kinetic was selected to determine myrosinase activity. Means of three replicate absorbance measurements were calculated after subtraction of the means of the background controls (100 μ L of extraction buffer, 25 μ L of 0.2 M phosphate buffer pH 6.5 and 50 μ L color reagent). Glucose concentrations were calculated using a linear standard curve. Samples were independently analyzed in triplicate and myrosinase activity was expressed as μ mol of glucose formed per minute and gram of dry matter (U/g d.m.) as well as per mg of protein (specific activity, U/mg soluble protein). The soluble protein concentration of cabbage extracts were determined using the DC Protein Assay (Bio-Rad Laboratories, Hercules, CA) following the manufacturer's instructions and bovine serum albumin was used as the standard.

Vitamin C content. Determination of vitamin C was performed in freeze-dried samples by capillary electrophoresis (CE) using a P/ACE system 2050 (Beckman Instruments, Fullerton, CA, USA) and UV detection at 254 nm as described earlier.²³ Briefly, 300 mg of freeze-dried cabbage were extracted with 20 mL of 3% (w/v) metaphosphoric acid (Sigma-Aldrich) and homogenised using an Ultra-Turrax homogenizer T25 Digital for 2 min. Final volume was

adjusted to 25 mL with 3% metaphosphoric acid. The resultant slurry was filtered through a Whatman No. 1 filter paper. A volume of 100 μ L isoascorbic acid (Fluka, Steinheim, Germany) at a concentration of 0.6 mg/mL containing 0.2% (w/v) D,L-dithiothreitol (Sigma-Aldrich) was added as internal standard into 1.5 mL of filtrate. Final volume was adjusted to 2 mL with 0.2% (w/v) D,L-dithiothreitol, mixed thoroughly and filtered through a 0.45 μ m membrane. D,L-dithiothreitol was added to prevent the oxidation of ascorbic acid to dehydroascorbic acid. Vitamin C content was quantified by external calibration using L-ascorbic acid (Sigma-Aldrich) (concentration range 0-50 μ g/mL) and using a response factor relative to the internal standard. Samples were independently analyzed in triplicate and results have been reported as mg L-ascorbic acid per gram of dry matter (mg/g d.m.).

Total phenolic compounds (TPC) content. The TPC content of white cabbage was determined in methanolic extracts using the Folin-Ciocalteu colorimetric method²⁴. Briefly, 1g of freeze-dried sample was suspended in 10 mL of 70% methanol and stirred for 1 h at room temperature. Extracts were filtered using Whatman No.1 filter paper. An aliquot of 400 μ L of a 20-fold dilution of each extract was mixed with 2.5 mL of distilled water, 1 mL 7.5% Na₂CO₃ (w/v), and 100 μ L of 2 N Folin-Ciocalteu reagent (Sigma-Aldrich). Samples were vortexed and incubated for 30 min at room temperature. The absorbance was measured at 736 nm using a microplate reader (Biotek Instruments). Total phenolics were quantified by external calibration using gallic acid (Sigma-Aldrich) as standard. Samples were independently analyzed in

triplicate and results were expressed as mg of gallic acid equivalents per gram of dry matter (mg GAE/g d.m.).

Antioxidant capacity. Oxygen Radical Antioxidant Capacity (ORAC) was determined in methanolic extracts by suspension of 1 g of freeze-dried sample in 10 mL of 70% methanol, which was stirred for 1 h at room temperature. Extracts were filtered using Whatman No.1 filter paper. The ORAC value was determined as described by Davalos et al.²⁵ Briefly, the reaction was carried out at 37 °C in 75 mM phosphate buffer (pH 7.4), and the final assay mixture (200 µL) contained 70 nM fluorescein (Sigma-Aldrich), 12 mM 2,2'-azobis(2-methylpropionamidine) dihydrochloride (Sigma-Aldrich), and antioxidant [1–8 µM Trolox (Sigma-Aldrich) or sample at different concentrations]. 2,2'-Azobis(2-methylpropionamidine) dihydrochloride and Trolox solutions were prepared daily, and fluorescein was diluted from a stock solution (1.17 mM) in 75 mM phosphate buffer (pH 7.4). Fluorescence measurements were carried out on a Polarstar Galaxy microplate reader (BMG Labtechnologies GmbH, Offenburg, Germany) equipped with a fluorescent filter (excitation 485 nm and emission 520 nm) using a black 96-F Microwell (Nunc A/S, Roskilde, Denmark). The plate was automatically shaken before the first reading, and the fluorescence was recorded every minute for 98 min. The equipment was controlled by Fluostar Galaxy software version (4.11–0) for fluorescence measurement. All reaction mixtures were prepared in triplicate, and at least two independent analyses were performed for each sample. The areas under the fluorescence decay curve (AUC), based on relative fluorescence values to the initial reading were recorded and the AUC of blanks

subtracted. Results were expressed as μmol Trolox equivalents (TE) per gram of dry matter ($\mu\text{mol TE/g d.m.}$).

Statistical analysis. Data were expressed as means $\pm$ standard deviation of three independent determinations. The statistical methods used were: one-way analysis of variance (ANOVA) using the least significant difference test to determine whether there were significant ($P \leq 0.05$) differences between genotypes within growing locations; a second one-way ANOVA to determine differences between the same genotype grown in different locations; principal component analysis to examine the relationships among variables; and stepwise discriminant analysis to select the variables most useful in differentiating the groups. The STATISTICA 7.0 (Statsoft Inc. Tulsa, OK, USA) and STATGRAPHICS 5.0 (Statistical Graphics Corp, Rockville, MD, USA) softwares for Windows were used.

RESULTS

Figure 1 shows the climatic conditions including global radiation ($\text{MJ/m}^2/\text{d}$), precipitation (L/m^2) and temperature (max, min, and mean, $^{\circ}\text{C}$) during the cabbage growing periods in the North and East areas of Spain. Global radiation was higher in the East than in the North only in November and December. Precipitation was higher in this geographical area during September and October and lower in November and December compared to the North area. Temperature was higher in the Eastern than in the Northern area of Spain.

240 Cabbages grown in the Northern region showed a content of total GLS
241 (Table 1) between 7.2 $\mu\text{mol/g d.m.}$ (cv. Candela) and 10.3 $\mu\text{mol/g d.m.}$ (cv.
242 Megaton) while those harvested in the East contained amounts between 10.7 -
243 20.0 $\mu\text{mol/g d.m.}$ Total GLS content was significantly ($P \leq 0.05$) higher in
244 cabbage cultivars grown in the East than in the North. Aliphatic GLS were
245 predominant and represented 75-90% and 84-89% in those cultivars grown in
246 Northern and Eastern areas, respectively. Cabbages cultivated in the North
247 contained sinigrin between 2.8-6.0 $\mu\text{mol/g d.m.}$, and glucobrassicin between 1.3-
248 5.4 $\mu\text{mol/g d.m.}$ while those cultivated in the East showed values of 3.4-6.6 and
249 1.6-8.2 $\mu\text{mol/g d.m.}$, respectively. Progoitrin, glucoraphanin and gluconapin
250 were present in lower amounts ($P \leq 0.05$), showing Eastern cabbages higher
251 content than those from the North ($P \leq 0.05$). Regarding indole GLS,
252 glucobrassicin was predominant and its ratio within the total GLS content
253 ranged from 8% in cv. Alfredo cultivated in the East to 23% in cv. Megaton
254 cultivated in the North. 4-Methoxy glucobrassicin and neoglucobrassicin were
255 present in negligible amounts and their contribution to total GLS content was
256 below 3%.

257 The soluble protein content in cultivars grown in the North (Table 2)
258 tended to be higher than those cultivated in the East. Significant differences
259 ($P \leq 0.05$) were found between growing locations with the exception of cv.
260 Bronco. Similar tendency was observed in myrosinase activity (Table 2).
261 Cabbages cultivated in the North of Spain shown significant differences
262 ($P \leq 0.05$) among genotypes. Cultivar Hinova exhibited the highest myrosinase
263 activity, followed by cv. Candela and cv. Megaton (36, 33 and 31 U/g d.m.,
264 respectively), while cv. Bronco and cv. Alfredo showed the lowest values (25

and 21 U/g d.m., respectively). The lowest ($P \leq 0.05$) specific myrosinase activity was observed in the Northern white cabbages cv. Megaton and cv. Alfredo (0.83 and 0.86 U/mg soluble protein, respectively). Among the white cabbages cultivated in the East, cv. Megaton and cv. Bronco presented significantly ($P \leq 0.05$) higher myrosinase activity (12-13 mg/g d.m.) than cv. Hinova, cv. Alfredo and cv. Candela (10-11 mg/g d.m.). Regarding myrosinase specific activity, cv. Candela showed the highest ($P \leq 0.05$) value (0.7 U/mg soluble protein) and no significant differences ($P \leq 0.05$) among the rest of cultivars were found (0.6 U/mg soluble protein) (Table 2).

The vitamin C content of cabbages (Table 3) was dependent on the cultivar and Spanish growing location. Among the cabbages cultivated in the North, cv. Alfredo exhibited the highest vitamin C content (3.6 mg/g d.m.) followed in descending order by cv. Hinova and cv. Bronco (2.9 mg/g d.m.), cv. Candela (2.7 mg/g d.m.) and cv. Megaton (2.4 mg/g d.m.). A significantly larger ($P \leq 0.05$) vitamin C content was found in cabbages grown in the East, irrespective of the cultivar; their values ranged from 4.2 to 6.0 mg/g d.m. In this Spanish geographical region, cv. Candela contained the highest vitamin C content (6.0 mg/g d.m.) whereas cv. Hinova, cv. Megaton, cv. Alfredo and cv. Bronco showed significantly lower ($P \leq 0.05$) values (5.0, 4.2, 5.5 and 5.6 mg/g d.m., respectively).

Northern white cabbage cv. Alfredo was characterized by the highest TPC content (12.3 mg/g d.m.) followed by cv. Bronco, while cv. Hinova, cv. Megaton and cv. Candela showed significantly ($P \leq 0.05$) lower values (9.0 to 9.2 mg/g d.m.) (Table 3). TPC of Eastern white cabbage cultivars were significantly ($P \leq 0.05$) lower than those from the North and ranged from 7.7 to 8.5 mg/g d.m.,

with the exception of cv. Megaton which showed the lowest TPC content (6.1 mg/g d.m.).

Antioxidant capacity of white cabbage (Table 3) was also influenced by cultivar and growing location. Across growing location, the antioxidant capacity ranged from 45.2 to 92 $\mu\text{mol TE/g d.m.}$; the cv. Alfredo grown in the North contained the highest level. In general, the influence of geographical growing location on the antioxidant capacity, irrespective of the cultivar, was similar to those observed for TPC content. Thus, cultivars grown in the Northern area of Spain exhibited significantly ($P \leq 0.05$) higher antioxidant capacity (ORAC) levels than those grown in the Eastern area.

In order to examine the relationship among the variables (GLS, vitamin C, myrosinase activity, TPC, and antioxidant capacity) of white cabbage cultivars and among cabbages grown in different geographical areas, principal component analysis from the correlation matrix was used in the present work. This analysis suggested that there are two main components which explained 87.5% of the total variance. The first principal component, which explained 74.1% of the total variance, was highly correlated with vitamin C content ($r=0.94$), myrosinase activity ($r=-0.87$), antioxidant capacity ($r=-0.95$), total GLS content ($r=0.83$) and TPC ($r=-0.70$). The second principal component explained 13.4% of the total variance. Figure 2 shows the graphical representation of the contribution of these quantitative variables of cabbages in the plane defined by the first two factors. Figure 3 shows the cabbage samples on the plane defined by the first two principal components and it can be observed that the Northern cabbages had clearly different values from the Eastern ones, and among these,

Bronco and Candela cultivars were completely separated from the rest of the Eastern cabbages.

From the matrix correlation carried out with the data, it was observed that the cabbages with the highest level of total GLS presented the highest level of vitamin C ($r=0.75$, $P\leq 0.05$) and the lowest level of antioxidant capacity ($r=-0.76$, $P\leq 0.05$). The cultivars with the highest amount of vitamin C had the lowest level of myrosinase activity ($r=-0.89$, $P\leq 0.05$) and antioxidant capacity ($r=-0.86$, $P\leq 0.05$), while the cabbages with the highest amount of TPC also presented the highest levels in antioxidant capacity ($r=0.71$, $P\leq 0.05$).

Table 4 shows the means and standard deviation of selected quantitative variables (GLS, vitamin C, TPC, myrosinase and antioxidant capacity) taking into account the cabbages cultivated in different areas of Spain. Cabbages grown in the North of Spain were characterised by the highest TPC content, myrosinase activity and antioxidant capacity ($P\leq 0.05$) and the lowest amounts of total GLS and vitamin C content ($P\leq 0.05$).

DISCUSSION

The content of total GLS found in the present work for cabbages are within the range reported in the literature.^{15,26-29} However, GLS values found in the present work are rather lower than those observed in turkish cabbages (50 and 70 $\mu\text{mol/g d.m.}$)¹⁴ and higher than those presented in cabbages from different European regions (3.3 to 7.7 $\mu\text{mol/g d.m.}$).³⁰

Regarding the profile of individual GLS observed in the current work, our results agree with several studies^{15,26,30} showing sinigrin as the predominant GLS in cabbage. The presence of the indolic glucobrassicin in the edible parts

of white cabbages should also be emphasized since this GLS is the precursor of indol-3-carbinol (I3C), a potent chemopreventive agent.³

Our results showed higher GLS content in the Eastern area of Spain where higher temperature and lower precipitations were registered. This could be due to increased synthesis of GLS precursors such as amino acids and sugars as consequence of higher temperatures.³⁰ These results are in agreement with previous studies. Martínez-Villaluenga et al.¹⁵ showed that white cabbage cv. Taler cultivated in summer led to larger total GLS content than cabbage cultivated in winter. Similarly, Cartea et al.²⁹ showed that total GLS concentration in cabbages harvested in the spring season was higher than those in autumn. On the other hand, in the present work glucoraphanin was present in larger amounts in those cabbages cultivated in the Eastern area (0.2-1.6 $\mu\text{mol/g d.m.}$) where higher temperatures were recorded. A reason for the different GLS profile in cabbages cultivated in different geographical locations could be due to enzymes involved in each GLS synthesis are affected differently by temperature and radiation.¹ Our results agree with those of Cartea et al.,²⁹ Martínez-Villaluenga et al.¹⁵ and Sarikamis et al.¹⁴ who showed that aliphatic GLS were the most abundant during autumn/winter season cabbages, and glucobrassicin was the predominant indolic GLS.

Regarding myrosinase activity determination, methods described by other authors^{31,32} were successful to determine myrosinase activity using a commercial enzyme but they did not work in cabbage extracts. These problems could be due to sample turbidity, as reported by Travers-Martin et al.²¹ In the present work, a modification to the Travers-Martin method has been introduced to increase myrosinase concentration that consisted of ultrafiltration (10 kDa

molecular weight cut off) of cabbage extracts and, hence, allowed the removal of both endogenous glucose and GLS, which could interfere in the analysis.

A large intraspecific variation for myrosinase activity among white cabbage cultivars has also been observed by Singh et al.³³ although it was rather lower than those for the Spanish cabbages found in the present work. Furthermore, such activity also seems to depend on the part of the cabbage analyzed, as has been shown by Charron and Sams.²⁶ Regarding climatic conditions, the growing season and year of cultivation have been reported to affect the myrosinase activity of *Brassica oleracea* cv. Early Round Dutch.³²

As observed in the present study, the vitamin C content in white cabbages was affected by the cultivar, which is in agreement with previous reports.^{13,34} Vitamin C contents found in white cabbage cultivars of the present study were comparable to those found by other researchers who obtained values ranging from 0.56 to 4.70 mg/g d.m.^{13,15,34,35} Growing location also had an impact on vitamin C content of white cabbages since cultivars grown in the East area of Spain presented higher vitamin C content than those grown in the North. This could be due to the different environmental conditions between both geographical locations. Mean temperature was higher in the East than in the North during the entire growing period (Figure 1). Additionally, global radiation was higher in the East than in the North of Spain during November and December (Figure 1). A higher light quantity and temperature during growth and development of plant tissues promote biosynthesis and accumulation of ascorbate because this antioxidant protects against environmental stress.³⁶ In particular, high concentrations of ascorbic acid in plant chloroplasts have been reported to protect against damaging oxygen-derived species that are produced

in the presence of light.³⁷ Our results are consistent with those of Vallejo et al.³⁸ and Martinez-Villaluenga et al.¹⁵ who observed higher vitamin C content in broccoli and white cabbage cv. Taler grown in the spring season than those grown in the winter season.

Total phenolic compounds (TPC) in white cabbage were influenced also by cultivar. These results are in accordance with those previously reported by other researchers.^{13,34,35} In addition, climatic factors such as temperature and radiation may have an impact in the biosynthesis of phenolic compounds. Our results showed that Northern white cabbage cultivars contained higher concentration of TPC than Eastern ones. Lower temperatures and lower global radiation (registered during November and December) in the North of Spain could be responsible for this effect. Low temperature enhances the formation of reactive oxygen species (ROS) which leads to gene expression of enzymes involved in the biosynthesis of phenolic compounds such as phenylalanine ammonia lyase and chalcone synthase as has been found in maize and *Arabidopsis thaliana*.³⁹ Besides low temperature, global radiation also induces the synthesis and accumulation of phenolic compounds such as flavonoids and hydroxycinnamic acids that function as shielding components against UV radiation.⁴⁰ A significant increase in flavonoids content during the cultivation of kale (*Brassica oleracea* var. *sabellica*) at low temperatures parallel to low radiation has been also reported¹⁶.

In accordance with results observed for antioxidants such as vitamin C and phenolic compounds, antioxidant capacity was also influenced by cultivar and growing location. Zietz et al.⁴¹ showed that antioxidant capacity was strongly positively correlated with total polyphenol content in four kale (*Brassica*

414 *oleracea* var. *sabellica*) cultivars grown in four different autumn/winter months.
415 The antioxidant capacity determined as ORAC values in the white cabbages
416 studied in the present work are within the range reported in 111 white cabbages
417 (23-146 μ moles TE/g d.m.).⁴²

418 Taking into consideration the results obtained, it may be concluded that
419 Spanish geographical locations experiencing low temperatures and radiation
420 during the growing season (Northern area) might be more appropriate for the
421 enhanced accumulation of myrosinase activity, total phenolic compounds and
422 antioxidant capacity of white cabbages, while the cabbages cultivated with high
423 temperature and radiation (Eastern area) would present a high amount of total
424 GLS and vitamin C. This knowledge provides information regarding the climatic
425 conditions in which white cabbages with the highest bioactive compounds
426 content can be obtained.

427

428 **ACKNOWLEDGEMENT**

429 E. P. is indebted to a JAE-doc grant funded by CSIC.

430

LITERATURE CITED

- (1) Verkerk, R.; Schreiner, M.; Krumbein, A.; Ciska, E.; Holst, B.; Rowland, I.; De Schrijver, R.; Hansen, M.; Gerhäuser, C.; Mithen, R.; Dekker, M. Glucosinolates in Brassica vegetables: the influence of the food supply chain on intake, bioavailability and human health. *Mol. Nutr. Food Res.* **2009**, *53*, S219-S265
- (2) Bonnesen, C.; Eggleston, I. M.; Hayes, J. D. Dietary indoles and isothiocyanates that are generated from cruciferous vegetables can both stimulate apoptosis and confer protection against DNA damage in human colon cell lines. *Cancer Res.* **2001**, *61*, 6120-6130.
- (3) Talalay, P.; Fahey, J. W. Phytochemicals from cruciferous plants protect against cancer by modulating carcinogen metabolisms. *J. Nutr.* **2001**, *31* (11 Suppl.), 3027S-3033S.
- (4) Podsedek, A. Natural antioxidant and antioxidant capacity of *Brassica* vegetables: A review. *LWT.* **2007**, *40*, 1-11.
- (5) Fenwick, G. R.; Heaney, R. K.; Mullin, W. J. Glucosinolates and their breakdown products in food and food plants. *Crit. Rev. Food Sci. Nutr.* **1983**, *18*, 123–201.
- (6) Das, S.; Tuagi, A. K.; Haur, H. Cancer modulation by glucosinolates: A review. *Curr. Sci.* **2000**, *79*, 1665-1671.
- (7) Wagner, A. E.; Rimbach, G. Ascorbigen: chemistry, occurrence and biological properties. *Clin. Dermatol.* **2009**, *27*, 217-224.
- (8) Duthie, S. J.; Ma, A.; Ross, M. A.; Collins, A. R. Antioxidant supplementation decreases oxidative DNA damage in human lymphocytes. *Cancer Res.* **1996**, *56*, 1291–1295.

- 456 (9) Wagner, A. E.; Huebbe, P.; Konishi, T.; Rahman, M. M.; Nakahara, M.,
457 Marsugo, S.; Rimbach, G. Free readical scavenging and antioxidant activity
458 of ascorbigen versus ascorbic acid: Studies in vitro and in cultures human
459 keratinocytes. *J. Agric. Food Chem.* **2008**, *56*, 11694-11699.
- 460 (10) Kelse, N. A.; Wilkins, H.M.; Linseman, D. A. Nutraceutical antioxidants as
461 novel neuroprotective agents. *Molecules*, **2010**, *15*, 7792-7814.
- 462 (11) Garcia-Lafuente, A.; Guillamon, E.; Villares, A.; Rostagno, M. A.;
463 Martinez, J. A. Flavonoids as anti-inflammatory agents: implications in
464 cancer and cardiovascular disease. *Inflam. Res.* **2009**, *58*, 537-552.
- 465 (12) Rosa, E. A. S., Heaney, R. K., Fenwick, G. R., Portas, C. A. M.
466 Glucosinolates in crop plants. *Hort. Rev.* **1997**, *19*, 99-215.
- 467 (13) Singh, J.; Upadhyay, A. K.; Prasad, K.; Bahadur, A.; Rai, M.; Variability of
468 carotenes, vitamin C, E and phenolics in *Brassica* vegetables. *J. Food*
469 *Comp. Anal.* **2007**, *20*, 106-112.
- 470 (14) Sarikamis, G.; Balkaya, A.; Yanmaz, R. Glucosinolates within a collection
471 of white head cabbages (*Brassica oleracea* var. *capitata* sub. var. *alba*)
472 from Turkey. *Afr. J. Biotechnol.* **2009**, *8*, 5046-5052.
- 473 (15) Martinez-Villaluenga, C.; Peñas, E.; Frias, J.; Ciska, E.; Honke, J.; Piskula,
474 M. K.; Kozłowska, H.; Vidal-Valverde, C. Influence of fermentation
475 conditions on glucosinolates, ascorbigen, and ascorbic acid content in
476 white cabbage (*Brassica oleracea* var. *capitata* cv. Taler) cultivated in
477 different seasons. *J. Food Sci.* **2009**, *74*, C62-C67.
- 478 (16) Schmidt, S.; Zietz, M.; Schreiner, M.; Rohn, S.; Kroh, L. W. Genotypic and
479 climatic influences on the concentration and composition of flavonoids in
480 kale (*Brassica oleracea* var. *sabellica*). *Food Chem.* **2010**, *119*, 1293-1299.

- 481 (17) Ludikhuyze, L.; Rodrigo, L.; Hendrickx, M. The activity of myrosinase from
482 broccoli (*Brassica oleracea* L. cv. *Italica*): influence of intrinsic and extrinsic
483 factors. *J. Food Prot.* **2000**, 63, 400-403.
- 484 (18) Rask, L.; Andreasson, E.; Ekbom, B.; Eriksson, S.; Pontoppidan, B.; Meijer,
485 J. Myrosinase: Gene family evolution and herbivore defense in
486 *Brassicaceae*. *Plant Mol. Biol.* **2000**, 42, 93-113.
- 487 (19) Bones, A. M.; Rossiter, J. T. The myrosinase-glucosinolate system, its
488 organisation and biochemistry. *Physiol. Plant.* **1996**, 97, 194-208.
- 489 (20) Official Journal of the European Communities. Oil seeds – Determination of
490 glucosinolates by high performance liquid chromatography. No. L 170, vol.
491 33 (3 July 1990).
- 492 (21) Travers-Martin, N.; Kuhlmann, F.; Müller, C. Revised determination of free
493 and complexed myrosinase activities in plant extracts. *Plant Physiol.*
494 *Biochem.* **2008**, 46, 506-516.
- 495 (22) Trinder, P. Determination of blood glucose using an oxidase-peroxidase
496 system with a non-carcinogenic chromogen. *J. Clin. Pathol.* **1969**, 22, 158-
497 161.
- 498 (23) Peñas, E.; Frias, J.; Sidro, B.; Vidal-Valverde, C. Chemical evaluation and
499 sensory quality of sauerkrauts obtained by natural and induced
500 fermentations at different NaCl levels from *Brassica oleracea* Var. *capitata*
501 cv. Bronco grown in Eastern Spain. Effect of storage. *J. Agric. Food Chem.*
502 **2010**, 58, 3549-3557.
- 503 (24) Singleton, V.; Orthofer, R.; Lamuela-Raventos, R. M. Analysis of total
504 phenols and other oxidation substrates and antioxidants by means of Folin-
505 Ciocalteu reagent. *Meth. Enzymol.* **1999**, 299, 152–175.

- 506 (25) Davalos, A.; Gomez-Cordoves, C.; Bartolome, B. Extending applicability of
507 the oxygen radical absorbance capacity (ORAC-fluorescein) assay. *J.*
508 *Agric. Food Chem.* **2004**, *52*, 48-54.
- 509 (26) Charron, C. S.; Sams, C. E. Glucosinolate content and myrosinase activity
510 in rapid-cycling *Brassica oleracea* grown in a controlled environment. *J.*
511 *Amer. Soc. Hort. Sci.* **2004**, *129*, 321-330.
- 512 (27) Ciska, E.; Pathak, D. Glucosinolate derivatives in stored fermented
513 cabbage. *J. Agric. Food Chem.* **2004**, *52*, 7938-7943.
- 514 (28) Radovich, T. J. K., Kleinhenz, D., Streeter, J. G., Miller, A. R., Scheerens,
515 J. C. Planting date affects total glucosinolate concentrations in six
516 commercial cabbage cultivars. *Hortic. Sci.* **2005**, *40*, 106-110.
- 517 (29) Cartea, M. E.; Velasco, P.; Obregón, S.; Padilla, G.; de Haro, A. Seasonal
518 variation in glucosinolate content in *Brassica oleracea* crops grown in
519 northwestern Spain. *Phytochem.* **2008**, *69*, 403-410.
- 520 (30) Kusznierevich, B.; Bartoszek, A.; Wolska, L.; Drzewiecki, J.; Gorinstein, S.,
521 Namiesnik, J. Partial characterization of white cabbage (*Brassica oleracea*
522 var. capitata f. alba) from different regions by glucosinolates, bioactive
523 compounds, total antioxidant activities and proteins. *LWT.* **2008**, *41*, 1-9.
- 524 (31) Li, X.; Kushad, M. M. Correlation of glucosinolate content to myrosinase
525 activity in horseradish (*Armoracia rusticana*). *J. Agric. Food Chem.* **2004**,
526 *52*, 6950-6955.
- 527 (32) Charron, C. S.; Saxton, A. M.; Carl, E. S. Relationship of climate and
528 genotype to seasonal variation in the glucosinolate–myrosinase system II.
529 Myrosinase activity in ten cultivars of *Brassica oleracea* grown in fall and
530 spring seasons. *J. Sci. Food Agric.* **2005**, *85*, 682–690.

- 531 (33) Singh, J.; Rai, M.; Upadhyay, A. K.; Prasad, K. Sinigrin (2-propenyl
532 glucosinolate) content and myrosinase activity in *Brassica* vegetables. *Int.*
533 *J. Veg. Sci.* **2007**, *13*, 21-31.
- 534 (34) Podsedek, A.; Sosnowska, D.; Redzynia, M.; Anders, B. Antioxidant
535 capacity of *Brassica oleracea* dietary antioxidants. *Int. J. Food Sci.*
536 *Technol.* **2006**, *41*, 49-58.
- 537 (35) Martinez, S.; Olmos, I.; Carballo, J.; Franco, I. Quality parameters of
538 *Brassica* spp. grown in northwest Spain. *Int. J. Food Sci. Technol.* **2010**,
539 *45*, 776-783.
- 540 (36) Ishikawa, T.; Shigeoka, S. Recent advances in ascorbate biosynthesis the
541 physiological significance of ascorbate peroxidase in photosynthesizing
542 organisms. *Biosci. Biotechnol. Biochem.* **2008**, *72*, 1143-1154.
- 543 (37) Smirnoff, N. Ascorbate biosynthesis and function in photoprotection. *Philos.*
544 *Trans. R. Soc. Lond. B Biol. Sci.* **2000**, *355*, 1455–1464.
- 545 (38) Vallejo, F.; Tomas-Barberan F. A.; Gonzalez Benavente-García, A.; Garcia-
546 Viguera, C. Total and individual glucosinolate contents in inflorescences of
547 eight broccoli cultivars grown under various climatic and fertilization
548 conditions. *J. Sci. Food Agric.* **2004**, *83*, 307-313.
- 549 (39) Leyva, A.; Jarillo, J. A.; Salinas, J.; Martinezzapater, J. M. Low temperature
550 induces the accumulation of phenylalanine ammonialyase and chalcone
551 synthase messenger-RNAs of *Arabidopsis thaliana* in a light-dependent
552 manner. *Plant Physiol.* **1995**, *108*, 39-46.
- 553 (40) Edreva, A. The importance of non-photosynthetic pigments and cinnamic
554 acid derivatives in the photoprotection. *Agric. Ecosys. Environ.* **2005**, *106*,
555 135-146.

(41) Zietz, M.; Weckmuller, A.; Schmidt, S; Rohn, S.; Schreiner, M.; Krumbein, A.; Kroh, W. Genotypic and climatic influence on the antioxidant activity of flavonoids in kale (*Brassica oleracea* var. *sabellica*). *J. Agric. Food Chem.* **2010**, *58*, 2123-2130.

(42) USDA Database for the oxygen radical absorbance capacity (ORAC) of selected foods, Release 2. 2010. U.S. Department of Agriculture, ML, USA.

Funding Sources

This work was supported by the Spanish Commission of Science and Technology (CICYT), Project number AGL2007-62044.

568 **FIGURE CAPTIONS**

569 Figure 1. Climatic conditions during cabbage growing period in the Northern
570 and Eastern areas of Spain.

571

572 Figure 2. Plot of the contribution of variables (total GLS, vitamin C, TPC,
573 antioxidant capacity and mirosinase activity) of cabbages in the plane defined
574 by the first two factors

575

576 Figure 3. Plot of cabbage cultivars growing in Northern and Eastern areas of
577 Spain in the plane defined by the first two principal components and ellipses for
578 95% confidence.

Table 1. Individual and total GLS content ($\mu\text{mol/g d.m.}$) in five white cabbage (*Brassica oleracea* var. capitata) cultivars grown in two geographical regions of Spain*.

Growing location	Cabbage cultivars	Aliphatic GLS					Indole			Total GLS
		IB	PROG	SIN	GRAP	NAP	GB	4-met-GB	neo-GB	
Northern area	Hinova	2.06 $\pm$ 0.07 ^a	0.22 $\pm$ 0.07	6.02 $\pm$ 0.17	ND ^a	0.13 $\pm$ 0.05	1.00 $\pm$ 0.12 ^a _A	0.08 $\pm$ 0.02	0.08 $\pm$ 0.01 ^a	9.61 ^a
	Megaton	1.33 $\pm$ 0.33	2.25 $\pm$ 0.03	4.06 $\pm$ 0.01 ^b	0.08 $\pm$ 0.02	ND ^a	2.41 $\pm$ 0.34	0.12 $\pm$ 0.03 ^b	0.09 $\pm$ 0.02 ^a	10.33 ^b
	Alfredo	5.42 $\pm$ 0.21	0.14 $\pm$ 0.03 ^a	3.44 $\pm$ 0.43 ^a	0.11 $\pm$ 0.03	ND ^a	0.89 $\pm$ 0.09 ^a _B	0.05 $\pm$ 0.00 ^a	0.09 $\pm$ 0.00 ^a	10.13 ^{ab}
	Candela	1.91 $\pm$ 0.05 ^a	0.85 $\pm$ 0.04	2.77 $\pm$ 0.06 ^a	ND ^a	ND ^a	1.39 $\pm$ 0.05	0.03 $\pm$ 0.00 ^a	0.25 $\pm$ 0.01	7.20
	Bronco	2.67 $\pm$ 0.24	0.15 $\pm$ 0.02 ^a	4.46 $\pm$ 0.25 ^b	ND ^a	ND ^a	1.02 $\pm$ 0.18 ^a	0.11 $\pm$ 0.04 ^b	0.03 $\pm$ 0.01 _A	8.44
Eastern area	Hinova	4.18 $\pm$ 0.37	0.31 $\pm$ 0.03	4.18 $\pm$ 0.33	0.17 $\pm$ 0.01	0.57 $\pm$ 0.06	1.11 $\pm$ 0.11 ^b _A	0.16 $\pm$ 0.01	0.05 $\pm$ 0.01	10.68 ^c
	Megaton	1.65 $\pm$ 0.03	3.19 $\pm$ 0.03	3.40 $\pm$ 0.03	1.61 $\pm$ 0.07	0.28 $\pm$ 0.07	1.12 $\pm$ 0.16 ^b	0.26 $\pm$ 0.05	0.02 $\pm$ 0.01 ^b	11.53 ^{cd}
	Alfredo	2.88 $\pm$ 0.73	1.10 $\pm$ 0.27 ^b	5.82 $\pm$ 0.55 ^c	0.74 $\pm$ 0.10	0.24 $\pm$ 0.07	0.87 $\pm$ 0.07 _B	0.36 $\pm$ 0.03	0.03 $\pm$ 0.03 ^b	12.05 ^d
	Candela	4.19 $\pm$ 0.42	2.29 $\pm$ 0.17	6.25 $\pm$ 0.40 ^{cd}	0.50 $\pm$ 0.05	0.68 $\pm$ 0.07	1.53 $\pm$ 0.11	0.13 $\pm$ 0.01	0.07 $\pm$ 0.01	15.64
	Bronco	8.19 $\pm$ 0.71	1.23 $\pm$ 0.08 ^b	6.62 $\pm$ 0.45 ^d	0.25 $\pm$ 0.02	0.41 $\pm$ 0.03	3.06 $\pm$ 0.21	0.21 $\pm$ 0.02	0.03 $\pm$ 0.00 ^b _A	20.00

* Mean values $\pm$ standard deviation of three independent experiments.

The same subscripts in the same column mean not significant differences between growing locations for each genotype ($P \leq 0.05$).

The same superscripts in the same column mean not significant differences among genotypes for each growing location ($P \leq 0.05$).

IB=Glucobrassicin; PROG=Progoitrin; SIN=Sinigrin; GRAP=Glucoraphanin; NAP=Gluconapin; GB=Glucobrassicin, 4-met-GB=4-Methoxy-glucobrassicin, neo-GB=Neo-glucobrassicin; Total GLS=total glucosinolates. ND = not detected.

Table 2. Myrosinase activity and soluble protein content in five white cabbage (*Brassica oleracea* var. capitata) cultivars grown in two geographical regions of Spain*.

Growing location	Cabbage cultivars	Soluble protein (mg /g d.m.)	Myrosinase activity (U/g d.m.)	Myrosinase activity (U/mg soluble protein)
Northern area	Hinova	20.00±1.29 ^a	36.28±1.25	1.82±0.06
	Megaton	37.20±2.69	30.61±1.60	0.83±0.02 ^a
	Alfredo	24.45±0.72 ^b	21.06±0.50	0.86±0.03 ^a
	Candela	23.85±2.21 ^b	33.10±1.49	1.39±0.07
	Bronco	20.70±0.93 ^a _A	25.20±0.66	1.22±0.06
Eastern area	Hinova	17.13±0.50	10.72±1.20 ^a	0.63±0.08 ^a
	Megaton	19.69±0.53	12.30±0.38 ^b	0.63±0.03 ^a
	Alfredo	15.73±0.44	9.89±0.29 ^a	0.63±0.01 ^a
	Candela	13.68±0.92	10.00±0.97 ^a	0.73±0.03
	Bronco	20.95±0.61 _A	12.89±0.56 ^b	0.62±0.04 ^a

* Mean values ± standard deviation of three independent experiments.

The same subscripts in the same column mean not significant differences between growing locations for each genotype ($P \leq 0.05$).

The same superscripts in the same column mean not significant differences among genotypes for each growing location ($P \leq 0.05$).

Table 3. Vitamin C, total phenolic compounds (TPC), antioxidant capacity (ORAC) and water content in five white cabbage (*Brassica oleracea* var. *capitata*) cultivars grown in two geographical regions of Spain*.

Growing location	Cabbage cultivars	Vitamin C (mg/g d.m.)	TPC (mg GAE/g) d.m.)	ORAC (μmoles TE/g d.m.)	Water (%)
Northern Area	Hinova	2.92 ± 0.19 ^a	9.21±0.16 ^a	83.80±3.29 ^a	92.1
	Megaton	2.41 ± 0.17	9.15±1.43 ^a	70.48±1.85	91.7
	Alfredo	3.58 ± 0.12	12.26±1.53	91.97±3.60	91.9
	Candela	2.68 ± 0.12	9.06±0.44 ^a	77.37±5.63	89.9
	Bronco	2.89 ± 0.11 ^a	10.41±0.02	84.44±3.86 ^a	92.2
Eastern area	Hinova	5.04 ± 0.13	8.48±0.19 ^b	57.39±3.29	91.7
	Megaton	4.20 ± 0.22	6.09±1.12	63.80±2.02	91.7
	Alfredo	5.52 ± 0.09 ^a	8.56±0.58 ^b	51.32±0.64	91.3
	Candela	5.99 ± 0.08	7.76±0.05 ^a	47.81±0.59	88.9
	Bronco	5.62 ± 0.23 ^a	7.73±0.13 ^a	45.17±1.53	91.2

* Mean values ± standard deviation of three independent experiments.

The same subscripts in the same column mean not significant differences between growing locations for each genotype (P≤0.05).

The same superscripts in the same column mean not significant differences among genotypes for each growing location (P≤0.05).

Table 4. Mean value $\pm$ SD of discriminant variables of growing location of cabbage cultivars*

Growing location	Total Glucosinolates ($\mu\text{mol/g d.m.}$)	Vitamin C (mg/g d.m.)	Total Polyphenols (mg GAE/g d.m.)	Myrosinase activity (U/g d.m.)	Antioxidant capacity ($\mu\text{moles TE/g d.m.}$)
Northern area	9.14 $\pm$ 1.23 ^a	2.91 $\pm$ 0.42 ^a	10.02 $\pm$ 1.30 ^b	29.25 $\pm$ 5.69 ^b	81.61 $\pm$ 7.86 ^b
Eastern area	14.27 $\pm$ 3.87 ^b	5.28 $\pm$ 0.64 ^b	7.72 $\pm$ 0.90 ^a	11.28 $\pm$ 1.43 ^a	52.83 $\pm$ 6.98 ^a

* The same superscripts in the same column mean not significant difference ($P \leq 0.05$)

Figure 1.

Figure 2.

Projection of the variables on the factor-plane (1 x 2)

Figure 3.