A Synthron for a 14-Electron Ir(III) Species: Catalyst for Highly Selective β -(Z) Hydrosilylation of Terminal Alkynes

Manuel Iglesias,*^a María Pérez-Nicolás,^a Pablo J. Sanz Miguel,^a Victor Polo,^b Francisco J. Fernández-Alvarez,*^a Jesús J. Pérez-Torrente,^a and Luis A. Oro*^a

Abstract: A synthron for a 14-electron Ir(III) species is described. The geometrical control exerted by the ligand system over the Ir-alkenyl intermediate in hydrosilylation of terminal alkynes precludes formation of the more thermodynamically stable β -(E)-vinylsilane, thus affording the β -(Z) isomer in excellent yields.

Bis-N-heterocyclic carbenes (bis-NHCs) are a fascinating class of ligands. The high σ -donor ability of NHCs and the strength of the resulting C–M bond, together with the inherent stability arising from the chelating nature of such ligands, render extremely robust electron rich transition metal species. In particular, $[M(\text{bis-NHC})(\text{OAc})_2]$ complexes ($M = \text{Rh}$ [1] and Ir[2]) have been demonstrated to be efficient air stable catalysts for transfer hydrogenation.

Transition-metal complexes containing latent coordination sites[3] have attracted much attention due to their central role as transient intermediates in many homogeneous catalytic processes.[4] Coordinating interactions with the solvent or fragments of the ligand system[5] generally stabilise coordinatively unsaturated complexes in solution. Hence, fine tuning of steric and electronic parameters of ligands with hemilabile character in order to achieve a truly reversible coordination—i.e. relatively small energy differences between open and closed conformations—is a matter of great interest.[6]

Hoping to shed some light on the effect of heterotopic ligands of hemilabile character in the generation of latent coordination sites, we envisaged a 14-electron bis-NHC iridium(III) fragment stabilised by two hemilabile side arms (Scheme 1), which would allow ready accessible coordination sites and concomitant protection of the active catalyst. Herein, we wish to report the synthetic route to such species and its remarkable β -(Z)-selectivity in catalysed hydrosilylation of terminal alkynes under mild conditions.

The aforementioned 14-electron metal fragment $[\text{Ir}(\text{bis-NHC})\text{I}_2]^+$ was generated upon treatment of $[\text{Ir}(\text{L-C,C})(\text{AcO})\text{I}_2]$ ($\text{L} = \text{methylenebis}[(\text{N-2-methoxyethyl})\text{imidazole-2-ylidene}]$, $\text{AcO} = \text{acetate}$) (2) with one equivalent of $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ in dry dichloromethane at 0 °C. Stabilisation by coordination of the oxygen atoms in the wingtip groups to the metal centre permits isolation of $[\text{Ir}(\text{L-C,C,O,O})\text{I}_2]\text{BF}_4$ (3), as a crystalline orange solid in 74% yield (Scheme 1 and Supporting Information).

Scheme 1 Synthetic route for the preparation of complex 3. Reaction conditions: i) NaI, KOAc, [Ir(cod)Cl]₂ in acetonitrile, 2 days at 80 °C; ii) HBF₄·Et₂O, in CH₂Cl₂ at 0 °C.

Successful removal of the acetato ligand in 2 was indicated by loss of the peak corresponding to the methyl group of the acetate in ¹H and ¹³C{¹H} NMR spectra at δ 1.85 and 25.9 ppm, respectively. Moreover, the A₂X₂ spin system (3J_{HH} = 4.8 Hz) corresponding to the protons of the methylene groups which appear at δ 3.75 (OCH₂) and 4.57 (NCH₂) ppm in 2 becomes an AA'XX' spin system with resonances centred at δ 4.51 (OCH₂) and 4.74 (NCH₂) ppm in 3. The singlet of the methoxy group undergoes a downfield shift similar to that observed for the OCH₂ groups upon coordination of the ethers (from δ 3.31 in 2 to 4.18 ppm in 3).

Fig. 1 View of 3 with two apical iodide ligands and a “tetradentate” bis-NHC ligand. Hydrogen atoms and BF₄[−] counterion are omitted for clarity.

Complex 3 was isolated by slow diffusion of diethyl ether into a saturated acetone solution. Its global connectivity pattern was confirmed by single crystal X-ray diffraction.[7] The molecular structure of cation 3 (Figure 1) shows the expected octahedral geometry for the central Ir^{III} metal ion. The two iodo ligands are located at the axial positions (Ir1–I1, 2.6750(3) Å; Ir1–I2, 2.6614(3) Å; I2–Ir1–I1, 179.396(18)°), while the equatorial sites are occupied by the “tetradentate” bis-NHC ligand. Both cis-positioned NHC-rings (Ir1–C12, 1.955(4) Å; Ir1–C22, 1.946(5) Å) extend their side arms toward the other two available positions, resulting in long distances between Ir1 and the oxygen atoms (Ir1–O43, 2.200(3) Å; Ir1–O53, 2.204(3) Å). Dihedral angles between both NHC-rings and the equatorial coordination plane of Ir1 are 16.2(2)° and 16.9(2)°. We are aware of only three additional examples in which a transition metal binds simultaneously an NHC-ring and its ether side arm.[8]

Complex 3 was tested in the hydrosilylation of terminal alkynes in order to explore the influence of the postulated latent coordination sites and the hypothetical generation of a cavity, formed by dissociation of the oxygen atoms, which would preclude formation of β-(E) isomers

due to steric interactions with the ligands. The major issue concerning this reaction is that arising from its selectivity, which very much depends on the nature of the catalyst and the reaction conditions—the selective formation of one of the reaction products being not easily achievable in most cases. Among the three possible isomers obtainable from hydrosilylation of terminal alkynes, the β -(E) is the most thermodynamically stable and usually the major reaction product. Regarded as much more challenging targets, β -(Z)-vinylsilanes, are highly valuable building blocks in organic synthesis, e.g. synthesis of natural products such as cis-retinoids via Hiyama cross-coupling.[9] However, examples of their selective metal-catalysed synthesis are scarce.[10]

Table 1. Hydrosilylation of Terminal Alkynes.

entry	Alkyne (R)	Silane	(Z)/(E) ^a	yield(%) ^a
1	Ph	Ph ₂ MeSiH	91/9	94
2	Ph	PhMe ₂ SiH	90/10	95
3	Ph	Et ₃ SiH	92/8	93
4	4- ^t Bu-Ph	Ph ₂ MeSiH	91/9	92
5	4- ^t Bu-Ph	PhMe ₂ SiH	94/6	96
6	4- ^t Bu-Ph	Et ₃ SiH	91/9	91
7	4-MeO-Ph	Ph ₂ MeSiH	100/0	94
8	4-MeO-Ph	PhMe ₂ SiH	92/8	91
9	4-MeO-Ph	Et ₃ SiH	95/5	98
10	4-CF ₃ -Ph	Ph ₂ MeSiH	100/0	98
11	4-CF ₃ -Ph	PhMe ₂ SiH	89/11	97
12	4-CF ₃ -Ph	Et ₃ SiH	95/5	98
13	4-Me-Ph	Ph ₂ MeSiH	98/2	94
14	4-Me-Ph	PhMe ₂ SiH	95/5	96
15	4-Me-Ph	Et ₃ SiH	92/8	99
16	<i>n</i> -Bu	Ph ₂ MeSiH	87/13	86
17	<i>n</i> -Bu	PhMe ₂ SiH	73/27	82
18	<i>n</i> -Bu	Et ₃ SiH	85/15	86

General conditions: Alkyne (0.15 mmol), Silane (0.16 mmol), **3** (5 mol%) in 0.5 mL of dry acetone-d₆ at 50 °C for 5 h. ^a Yields and selectivities were calculated by ¹H NMR.

Using **3** as catalyst, a range of terminal alkynes were converted in excellent yields and selectivities to their corresponding β -(Z)-vinylsilanes with Ph₂MeSiH, PhMe₂SiH or Et₃SiH (1.1 equivalents) at 50 °C (Table 1). The corresponding alkyne was, in all cases, completely consumed within 5 h and no traces of catalyst decomposition were observed. The remarkably high activities and selectivities toward the β -(Z)-vinylsilanes presented in Table 1 are among the best hitherto reported for iridium catalysts.^{10f} The steric hindrance of the silane seems to play a significant role in the selectivity of the reaction, the best selectivities being those obtained with Ph₂MeSiH (Table 1). Regarding the alkyne, no major effects on the selectivity of the reaction arise from electronic effects of para substitution in phenylacetylenes. Moreover, the β -(Z) selectivity was maintained for aliphatic alkynes, namely 1-hexyne (entry 16). Additionally, it is worth to note that the hydrosilylation reaction can be carried out on a preparative scale retaining high conversions and selectivities as demonstrated for the synthesis of (Z)-dimethyl(4-methylstyryl)(phenyl)silane.

Kinetic studies performed for representative alkynes (R $\text{C}\equiv\text{CH}$) with Ph₂MeSiH revealed conversions above 70% after 1.5 h, and TOF_{1/2} (h⁻¹) values of 55 (R = 4-CF₃-Ph), 25 (R = 4-MeO-Ph) and 21 (R = Ph).

Under catalytic conditions ¹H-NMR spectroscopy shows that the percentage of **3**, the only metal species present in solution, remains constant throughout the reaction. This observation suggests an equilibrium between **3** and **4** (Scheme 2) caused by random dissociation of the oxygen atoms as the only conceivable option to generate an active catalyst which, at the end of the catalytic cycle, would convert back to **3**. This premise, in addition to the fact that no reaction between **3** and R $\text{C}\equiv\text{CH}$ was observed, strongly points toward initial formation of Ir(σ -Si-H) complex [11] **5** followed by oxidative addition to give Ir(V)10e,^[12] intermediate **6** as first step of the catalytic cycle (Scheme 2). Therefore, in the overall process complex **3** could be regarded as a synthon for the 14-electron fragment [Ir(L-C,C)I₂]⁺ due to reversible coordination of the wingtip groups.^[13]

We have observed that, in the ¹H DOSY spectrum,^[14] the peak of residual acetone shows a higher diffusion coefficient for a sample of acetone-d₆ ($D_c = 4.323 \times 10^{-9} \text{ m}^2\text{s}^{-1}$) than for a solution of **3** in acetone-d₆ (0.03 mmol in 0.5 mL of acetone-d₆; $D_c = 3.875 \times 10^{-9} \text{ m}^2\text{s}^{-1}$). This behaviour can be ascribed to an interaction between **3** and the acetone molecules that results in a reduction of their mobility, which could be interpreted as consequence of a certain degree of dissociation of the ether functions, essential requisite to generate the coordination vacancies needed for the catalytic reaction to happen.

The second step of the catalytic cycle requires coordination of the alkyne to give **7**. For this to occur, we postulate substitution of an iodide by the corresponding alkyne, which we expect to be the rate limiting step. This assumption is supported by the fact that a significant reduction of reaction rate was observed upon addition of excess NaI to the reaction mixture. The TOF_{1/2} (h⁻¹) decreases from 1250 to 706 in the case of the hydrosilylation of phenylacetylene with HSiPh₂Me.

In order to explain the β -(Z) selectivity, a 2,1-insertion of the alkyne into the Ir-Si bond according to the modified Chalk-Harrod mechanism¹⁵ must take place. Therefore, invoking a metal-assisted isomerisation,^[10d,16] the cavity about the two coordination vacancies

stabilises **8b** against **8a** through geometrical control exerted by the ligand system over the Ir-alkenyl intermediate (Scheme 2). This assumption is supported by the fact that the relative free energies of the postulated intermediates **8a** and **8b**, calculated by DFT (B3LYP) methods based on the model $[\text{Ir}(\text{L-C,C})(\text{CPh=CHSiMe}_3)(\text{I})(\text{H})]\text{BF}_4$, show that the latter is more stable ($\Delta G = 4.9 \text{ kcal mol}^{-1}$).^[17] Finally, the last step of the proposed Ir(III)/Ir(V) catalytic cycle would be the reductive elimination of the β -(Z)-vinylsilane from **8b** and coordination of the iodido ligand. In this context, it seems to be interesting to mention that there is increasing support for the existence of Ir(III)/Ir(V) cycles in iridium catalytic chemistry.^[18]

The proposed catalytic cycle rests on the following experimental observations and DFT calculations: (i) presence of **3** throughout the reaction and hints of an equilibrium between **3** and **4**, (ii) reduction of the reaction rate upon addition of NaI, and (iii) the unusual selectivity of the reaction, only explainable by a 2,1-insertion of the alkyne into the M–Si bond followed by reductive elimination of the vinylsilane from **8b**, which lies $4.9 \text{ kcal mol}^{-1}$ below **8a**.

Scheme 2 Proposed catalytic cycle for the hydrosilylation of terminal alkynes.

In summary, we have developed a straightforward synthetic route to a cationic bis-NHC iridium(III) complex (**3**) with two latent coordination sites accessible under catalytic conditions. As a first application, we have shown that the cavity created by dissociation of the side arms leads to unusually high β -(Z) selectivities in hydrosilylation of terminal alkynes under mild conditions. Remarkably, the X-ray structure of **3** represents the first example of a heterotopic Ir(NHC) complex featuring coordinated ether functions. Steric and electronic tuning of the

ether functions in order to optimise the catalytic activity and a deeper mechanism insight are currently on-going.

This work was supported by the Spanish Ministry of Economy and Competitiveness (MINECO/FEDER) (CTQ2010-15221, CSD2009-00050 CONSOLIDER INGENIO-2010, and CTQ2011-27593 projects, and “Ramón y Cajal” (P.J.S.M.) and “Juan de la Cierva” (M.I.) programmes) and the DGA/FSE (project E07).

Notes and references

aDepartamento de Química Inorgánica, Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), Universidad de Zaragoza–CSIC, 50009-Zaragoza, Spain. bDepartamento de Química Física, Universidad de Zaragoza, 50009, Zaragoza, Spain. E-mail: miglesia@unizar.es; paco@unizar.es; oro@unizar.es.

† Electronic Supplementary Information (ESI) available: Synthesis, characterization, and experimental details. See DOI: 10.1039/b000000x/

1. M. Albrecht, R. H. Crabtree, J. Mata, E. Peris, *Chem. Commun.*, 2002, 32.
2. M. Albrecht, J. R. Miecznikowski, A. Samuel, J. W. Faller, R. H. Crabtree, *Organometallics*, 2002, 21, 3596.
3. S. H. Strauss, *Chemtracts: Inorg. Chem.* 1994, 6, 1.
4. (a) J. P. Collman, L. S. Hegedus, J. R. Norton, R. G. Finke, In *Principles and Applications of Organotransition Metal Chemistry*; University Science: Mill Valley, CA, 1987. (b) A. Binobaid, M. Iglesias, D. Beetstra, A. Dervisi, I. Fallis, K. J. Cavell, *Eur. J. Inorg. Chem.*, 2010, 5426.
5. N. M. Scott, R. Dorta, E. D. Stevens, A. Correa, L. Cavallo, S. P. Nolan, *J. Am. Chem. Soc.*, 2005, 127, 3516.
6. (a) P. Braunstein, F. Naud, *Angew. Chem. Int. Ed.*, 2001, 40, 680. (b) D. S. McGuinness, K. J. Cavell, *Organometallics*, 2000, 19, 741.
7. Crystal data of 3: [C₁₆H₂₆BF₄IrN₄O₃], orthorhombic, Pna2(1), a = 19.5550(14) Å, b = 10.6594(8) Å, c = 11.3757(8) Å, Z = 4, Mr = 855.22 g mol⁻¹, V = 2371.2(3) Å³, D_{calcd} = 2.396 g cm⁻³, λ(Mo Kα) = 0.71073 Å, T = 100 K, μ = 8.291 mm⁻¹, 24968 reflections collected, 6159 observed (R_{int} = 0.0567), R₁(Fo) = 0.0252 [I > 2σ(I)], wR₂ (Fo²) = 0.0650 (all data), GOF = 1.023. CCDC 883047.
8. (a) C. L. Lund, M. J. Sgro, D. W. Stephan, *Organometallics* 2012, 31, 580. (b) I. Özdemir, S. Demir, B. Çetinkaya, L. Toupet, R. Castarlenas, C. Fischmeister, P. H. Dixneuf, *Eur. J. Inorg. Chem.*, 2007, 2862. (c) C. L. Lund, M. J. Sgro, D. W. Stephan, *Organometallics* 2012, 31, 802.
9. a) E. Langkopf, D. Schinzer, *Chem. Rev.*, 1995, 95, 1375. (b) J. Bergueiro, J. Montenegro, F. Cambeiro, C. Saá, S. López, *Chem. Eur. J.*, 2012, 18, 4401. (c) J. Montenegro, J. Bergueiro, C. Saá, S. López, *Org. Lett.*, 2009, 11, 141.

10. (a) E. Mas-Marza, M. Poyatos, M. Sanau, E. Peris, *Inorg. Chem.*, 2004, 43, 2213. (b) Y. Na, S. Chang, *Org. Lett.*, 2000, 2, 1887. (c) M. Nagao, K. Asano, K. Umeda, H. Katayama, F. Ozawa, *J. Org. Chem.*, 2005, 70, 10511. (d) R. S. Tanke, R. H. Crabtree, *J. Am. Chem. Soc.* 1990, 112, 7984. (e) V. S. Sridevi, W. Y. Fan, W. K. Leong, *Organometallics*, 2007, 26, 1157. (f) B. Marciniak, I. Kownacki, In *Iridium Complexes in Organic Synthesis*; L. A. Oro, and C. Claver, 2009 Wiley-VCH: Weinheim, 2009, Chapter 14; pp 349-352.
11. J. Y. Corey, *Chem. Rev.*, 2011, 111, 873.
12. For examples of Ir(V) silyl complexes see: (a) M.-J. Fernandez, P. M. Maitlis, *Organometallics*, 1983, 2, 164. (b) T. M. Gilbert, F. J. Hollander, R. G. Bergman, *J. Am. Chem. Soc.*, 1985, 107, 3508. (c) M. A. Esteruelas, F. J. Lahoz, E. Oñate, L. A. Oro, L. Rodríguez, *Organometallics*, 1996, 15, 823; (d) M. A. Esteruelas, F. J. Fernández-Alvarez, A. M. López, E. Oñate, P. Ruiz-Sánchez, *Organometallics*, 2006, 25, 5131.
13. D. T. Tilley, B. V. Mork, *J. Am. Chem. Soc.*, 2004, 126, 4375.
14. S. Cozzolino, M. G. Sanna, M. Valentini, *Magn. Reson. Chem.* 2008, 46, S16–S23.
15. A. J. Chalk, J. F. Harrod, *J. Am. Chem. Soc.*, 1965, 87, 16.
16. (a) I. Ojima, N. Clos, R. J. Donovan, P. Ingallina, *Organometallics*, 1990, 9, 3127. (b) M. V. Jiménez, J. J. Pérez-Torrente, M. I. Bartolomé, V. Gierz, F. J. Lahoz, L. A. Oro, *Organometallics*, 2008, 27, 224.
17. Coordination of the outcoming iodido ligand to 8a or 8b has been discarded based on DFT (B3LYP) calculations.
18. For examples of Ir(III)/Ir(V) catalytic cycles see: (a) J. Mazuela, P. O. Norby, P. G. Andersson, O. Pàmies, M. Diéguez, *J. Am. Chem. Soc.*, 2011, 133, 13634 (b) K. H. Hopmann, A. Bayer, *Organometallics*, 2011, 30, 2483 (c) T. L. Church, T. Rasmussen, P. G. Andersson, *Organometallics*, 2010, 29, 6769.