A diet rich in cocoa attenuates *N*-nitrosodiethylamine-induced liver injury in rats.

Ana Belén Granado-Serrano, María Angeles Martín, Laura Bravo, Luis Goya and Sonia Ramos*

Department of Metabolism and Nutrition

Instituto del Frío, Consejo Superior de Investigaciones Científicas (CSIC)

José Antonio Novais 10, 28040 Madrid (Spain)

Phone: +34.91.544.56.07

Fax: +34.91.549.36.27

Running title: Protective effects of cocoa in liver damage.

Keywords: Cocoa, liver, antioxidant defences, cell death, survival/proliferation pathways.

* Corresponding author: e-mail: s.ramos@if.csic.es.

Abbreviations: AKT/PKB, protein kinase B; ALP, alkaline phosphatase; ALT, alanine transaminase; AST, aspartate transaminase; DEN, *N*-nitrosodiethylamine; ERK, extracellular regulated kinase; FBS, fetal bovine serum; GPx, glutathione peroxidase; GR, glutathione reductase; GSH, glutathione; GST, glutathione-S-transferase; JNK, *c-jun* amino-terminal kinase; LDH, lactate dehydrogenase; MAPK, mitogen-activated protein kinase; OPT, *o*-phthalaldehyde.

Abstract

The effects of cocoa feeding against *N*-nitrosodiethylamine (DEN)-induced liver injury were studied in rats. Animals were divided into five groups. Groups 1 and 2 were fed with standard and cocoa diet, respectively. Groups 3 and 4 were injected with DEN at 2 and 4 weeks, and fed with standard and cocoa diet, respectively. Group 5 was treated with DEN, received the standard diet for 4 weeks and then it was replaced by the cocoa-diet. DEN-induced hepatic damage caused a significant increase in damage markers, as well as a decrease in the hepatic glutathione, diminished levels of p-ERK and enhanced protein carbonyl content, caspase-3 activity and values of p-AKT and p-JNK. The cocoa-rich diet prevented the reduction of hepatic glutathione concentration and catalase and GPx activities in DEN-injected rats, as well as diminished protein carbonyl content, caspase-3 activity, p-AKT and p-JNK levels, and increased GST activity. However, cocoa administration did not abrogate the DEN-induced body weight loss and the increased levels of hepatic-specific enzymes and LDH. These results suggested that cocoa-rich diet attenuates the DEN-induced liver injury.

Introduction

Chronic liver damage is a worldwide common pathology characterized by inflammation and fibrosis that can lead to chronic hepatitis, cirrhosis and cancer (Köhle, C. et al., 2008, Tessitore, L. and Bollito, E., 2006, Thorgeirsson, S. and Grisham, J., 2002). It has been well established that oxidative stress plays a causative role at the initiation, promotion and progression of hepatic diseases (Ramos, S., 2008, Vitaglione, P. et al., 2004) and that the liver is the main target for several toxic agents that can provoke free radical-mediated apoptosis (Jaeschke, H. et al., 2002).

Cocoa and its derived products are widely consumed in Europe and the United States (Vinson, J. et al., 2006) and they have increasingly attracted researchers, food manufacturers, as well as consumers due to their biological properties, which have been mostly related to its phenolic compounds (Lamuela-Raventós, R. et al., 2005). Cocoa flavonoids can influence several important biological functions by their free-radical scavenging ability or through the regulation of signal transduction pathways to stimulate apoptosis and/or to inhibit inflammation and proliferation in different human cancer cell lines (Lamuela-Raventós, R. et al., 2005, Martin, M. et al., 2009, Martin, M. et al., 2008, Ramos, S., 2007, Ramos, S., 2008). Therefore, cocoa phenolic compounds have been suggested not only as potential protective and even therapeutic agents to reduce liver damage (Ramos, S., 2008, Vitaglione, P. et al., 2004).

The modulation of phase I [glutathione reductase (GR), glutathione peroxidase (GPx) and catalase (CAT)] and phase II [glutathione-S-transferase (GST)] enzymes, and glutathione (GSH) levels play a primary role in the balance of the redox status through the reduction of reactive oxygen species (ROS) and peroxides produced in the organism, as well as in the detoxification of xenobiotics (Ramiro Puig, E. et al., 2007, Ramos, S., 2008). Cocoa has been

shown to enhance the activity of phase I enzymes in hepatic and thymocyte cells, respectively (Martin, M. et al., 2009, Ramiro Puig, E. et al., 2007), and it may also induce cell survival and proliferation, as well as protective effects, through the modulation of different transduction pathways, such as AKT and MAPKs (Martin, M. et al., 2009, Ramiro-Puig, E. et al., 2009). Additionally, upregulation of phase I and II enzymes mediated by MAPKs and/or AKT has been reported for different polyphenols (Masella, R. et al., 2005, Na, H. and Surh, Y., 2008). In this regard, induction of GPx and GR by cocoa has been shown to occur via ERK in hepatic cells (Martin, M. et al., 2009). Moreover, GST expression has been implicated in the regulation of cell proliferation and protection from apoptosis through JNK-mediated mechanisms (Holley, S.L. et al., 2007, Mates, J.M. et al., 2008).

N-nitrosodiethylamine (DEN) is a potent hepatotoxic, carcinogen and mutagen (Chuang, S. et al., 2000, Köhle, C. et al., 2008, Sreepriya, M. and Bali, G., 2006, Tessitore, L. and Bollito, E., 2006). Human exposure could occur through the diet (meat, whiskey, etc.) (Hecht, S., 1997, Sen, N. et al., 1980), in certain occupational settings, smoking or through the use of cosmetics, pharmaceutical products and agricultural chemicals (Hecht, S., 1997). DEN has been extensively used as an initiating carcinogen in experimental animal models (Chuang, S. et al., 2000, Köhle, C. et al., 2008, Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2005, Sreepriya, M. and Bali, G., 2006, Sundaresan, S. and Subramanian, P., 2008, Tessitore, L. and Bollito, E., 2006) and induces hepatic necrosis through metabolic activation by CYP2E1 in experimental animals (Kang, J. et al., 2007). Activation of DEN, which takes place mainly in liver microsomes (Kang, J. et al., 2007), has been shown to stimulate Kupfer cells leading to generate high levels of ROS, capable of damaging liver cells and participating in the induction of hepatocarcinogenesis (Kang, J. et al., 2007). It has been shown that a cocoa bean preparation protects against different types of cancers (Yamagishi, M. et al., 2002, Yamagishi, M. et al., 2003), as well as against a number

of hepatotoxic agents including nitrosamines (Amin, I. et al., 2004, Yamagishi, M. et al., 2000). However, the mechanisms by which cocoa elicits such hepatoprotective and chemopreventive effects related to antioxidant defences, as well as survival/proliferation pathways, remain poorly understood.

Thus, the aim of this study was to evaluate the protective effect of a cocoa-rich diet against DEN-induced liver toxicity. Here, we report that cocoa partly attenuates DEN-induced liver injury in rats via induction of antioxidant defence (GSH, CAT, GPx and GST) and modulation of signals related to cell death (caspase-3 and JNK) and survival/proliferation (AKT).

Materials and Methods

Materials and chemical

N-nitrosodiethylamine (DEN) was purchased from Sigma Chemical (Madrid, Spain). Anti-AKT and antiphospho-Ser473-AKT (p-AKT), anti-ERK1/2 and antiphospho-ERK1/2 (p-ERKs) recognizing ERK1/2 phosphorylated Thr202/Thy204, anti-JNK1/2 and antiphospho-JNK1/2 (p-JNKs) recognizing JNK1/2 phosphorylated Thr183/Tyr185 and anti- β -actin were obtained from Cell Signaling Technology (9271, 9272, 9101, 9102, 9251, 9252, and 4697, respectively; Izasa, Madrid, Spain). Caspase-3 substrate (Ac-DEVD-AMC) was purchased from Pharmingen (San Diego, CA). Materials and chemicals for electrophoresis and the Bradford reagent were from BioRad (BioRad Laboratories S.A., Madrid, Spain).

Cocoa

Natural Forastero cocoa powder (Nutrexpa, Barcelona, Spain) was used for this study. It contains epicatechin (383.5 mg/100g), catechin (116 mg/100g) and procyanidins (254.5 mg/100g) and non-flavonoid compounds such as theobromine. A detailed description of this cocoa is given elsewhere (Martin, M. et al., 2008).

Animal treatments

Forty five male Sprague-Dawley rats (6 weeks old) were obtained from the School of Medicine, Universidad Autónoma (Madrid, Spain). Animals were placed individually in stainless steel wire-bottomed metabolic cages housed in a room under controlled conditions (19-23 °C, 50-60 % humidity and 12 h light/darkness cycles).

Diets were prepared from a Fibre Free AIN-93M Purified Rodent Diet (Panlab S.L., Barcelona, Spain) providing all nutrients required by adult rats. Rats were randomly assorted into five different experimental groups (eight to ten animals per group) (Figure 1) and were

provided with food and water *ad libitum*. Initially, three groups received a standard diet (C, groups 1, 3 and 5) and two groups were fed with cocoa-rich diet (Ca, groups 2 and 4), containing the basal diet supplemented with 16 % powdered cocoa (providing 54.4 g of dietary fibre per Kg of diet), added as a source of polyphenols. In line with our previous research, this powdered supplement was formulated to provide 7.5 g of polyphenols per Kg of diet, since this percentage of cocoa supplementation improve the antioxidant defence system, the lipid profile and reduced the lipid peroxidation. (Lecumberri, E. et al., 2007). The composition of the diets is given in Table 1 and cocoa powder contains 26% proteins and 12% lipids.

To induce hepatic damage, three groups of rats (groups 3-5) were injected intraperitoneally with 200 mg/Kg DEN diluted in saline on days 14 and 28. Rats in the non-DEN treated groups were injected with saline the same day. In addition, on day 28, group 5 was changed from control diet to cocoa diet after the second DEN injection (Figure 1) to test the potential therapeutic activity of the cocoa-rich diet.

Animal weight and food intake were monitored daily for 6 weeks. Blood was harvested from the trunk after decapitation, and serum was separated by centrifugation at 1000 xg, 10 min, 4°C for further biochemical analysis. Livers were collected, weighted and frozen in liquid N₂ and stored at -80°C.

Animals were treated according to the Institutional Care Instructions (Bioethical Commission from Consejo Superior de Investigaciones Cientificas).

Analysis of hepatic enzymes (AST, ALT, ALP)

As markers for liver function in serum, the activity of aspartate transaminase (AST), alanine transaminase (ALT) and alkaline phosphatase (ALP) [expressed as Units (U)/L serum] were analysed. Serum AST was measured spectrophotometrically at 340 nm in the presence of α -

ketoglutarate, aspartate, NADH and malate dehydrogenase, following the method of Rej and Horder (Rej, R. and Horder, M., 1984). Serum ALT was assayed at 340 nm in the presence of α -ketoglutarate, pyruvate, NADH and lactate dehydrogenase as described by Horder and Rej (Horder, M. and Rej, R., 1984). Serum ALP was analysed spectrophotometrically at 405 nm using the formation of *p*-nitrophenol from *p*-nitrophenylphosphate as a substrate (Principato, G.B. et al., 1985). Serum protein concentration was measured by the Bradford reagent.

Determination of GSH

The content of GSH was quantitated by the fluorometric assay of Hissin and Hilf (Hissin, P. and Hilf, R., 1976). The method takes advantage of the reaction of GSH with *o*-phthalaldehyde (OPT) at pH 8.0, which generates fluorescence.

Livers were homogenized (1:20 w/v) in 50 mM phosphate buffer pH 7.0, proteins precipitated with 5% trichloroacetic acid and then centrifuged for 30 min at 10.000 xg. Following, 50 μ L of the clear supernatant were transferred to a 96 multiwell plate for the assay. Fluorescence was measured at an excitation wavelength of 340 nm and emission wavelength of 460 nm. The results were interpolated in a glutathione standard curve (5 ng-1 μ g) and expressed as nmol GSH per milligram of protein, which was determined by the Bradford reagent. The precise protocol has been described elsewhere (Alia, M. et al., 2003, Alía, M. et al., 2006).

Determination of GPx, GR, CAT and GST activities

The activity of antioxidant enzymes (GPx, GR and catalase) was determined in liver homogenates. Livers (0.3 g) were homogenized (1:5 w/v) in 0.25 M Tris, 0.2 M sucrose and 5 mM 1,4-dithiothreitol (DTT) buffer pH 7.4 and centrifuged at 3.000 xg for 15 min. Determination of GPx activity is based on the oxidation of GSH by GPx, using t-BOOH as a substrate, coupled to the disappearance of NADPH by GR (Gunzler, W. et al., 1974),

expressing the results as mU/mg protein. GR activity was determined by following the decrease in absorbance due to the oxidation of NADPH utilized in the reduction of oxidized glutathione (Goldberg, D. and Spooner, R.). The GR activity was expressed as $\mu\text{U}/\text{mg}$ protein. Catalase (CAT) activity was determined by following the decomposition of H_2O_2 measured as a decrease in absorbance at 240 nm (Aebi, H., 1987) and expressed as UI/mg protein. The methods have been previously described (Alia, M. et al., 2003, Alía, M. et al., 2006).

Analysis of GST was carried out in liver and serum by the Biovision Commercial kit GST fluorometric activity assay (Biovision, Mountain View, CA) and the activities were expressed as U/mg protein and mU/mg protein, respectively. Protein was measured by the Bradford reagent.

Protein carbonyl determination in liver

Protein oxidation of liver homogenates was measured as carbonyl groups content according to the method of Richert *et al.* (Richert, S. et al., 2002). Absorbance was measured at 360 nm and carbonyl content was expressed as nmol/mg protein using an extinction coefficient of $22000 \text{ nmol L}^{-1} \text{ cm}^{-1}$. Protein in liver homogenates was determined by the Bradford reagent.

Lactate Dehydrogenase (LDH) Leakage Assay

Livers were homogenized (1:5 w/v) in 50 mM phosphate buffer (PBS) pH 7.5, centrifuged for 30 min at 1000 xg and the supernatants collected. LDH was spectrophotometrically assayed in the samples by measuring the disappearance of NADH at 340 nm, as previously described (Alía, M. et al., 2006, Bergmeyer, H. and Bernt, E., 1974). The reaction mixture contained 5 mM pyruvate, 0.35 mM NADH, 84 mM Tris and 50 μg of protein per condition. Enzyme activity was calculated by using an extinction coefficient of $6.22 \text{ mM}^{-1} \text{ cm}^{-1}$ and expressed in

units using mg^{-1} protein. One unit of the enzyme activity is defined as the amount of enzyme required to oxidize $1 \mu\text{mol NADH min}^{-1}$.

Liver caspase-3 activity

Activation of caspase-3 was determined as previously described (Granado-Serrano, A. et al., 2007). Briefly, livers were lysed (1:5 w/v) in a buffer containing 5 mM Tris (pH 8), 20 mM EDTA, and 0.5 % Triton X-100. The reaction mixture contained 20 mM HEPES (pH 7), 10 % glycerol, 2 mM DTT, 50 μg of protein per condition and 20 μM Ac-DEVDAMC (*N*-acetyl-Asp-Glu-Val-Asp-7-amino-4-methylcoumarin) as substrate. Enzymatic activity was determined by measuring fluorescence at an excitation wavelength of 380 nm and an emission wavelength of 440 nm (Bio-Tek, Winooski, VT, USA).

Preparation of cell lysates for Western blotting

Samples of frozen liver were homogenized 1:10 (w:v) in extraction buffer [50 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM MgCl_2 , 1 mM CaCl_2 , 10 % glycerol, 10 mM $\text{Na}_4\text{P}_2\text{O}_7$, 10 mM NaF, 2 mM EDTA, 1 % Nonidet P-40, 2 mM Na_3VO_4 , 5 $\mu\text{g/mL}$ leupeptin, 20 $\mu\text{g/mL}$ aprotinin, 2 mM benzamidin and 2 mM phenylmethylsulphonyl fluoride (PMSF)] to detect AKT, p-AKT, ERK1/2, p-ERKs, JNK1/2 and p-JNKs (Gavete, L. et al., 2005). Homogenates were centrifuged at 14000 $\times g$ for 60min and the supernatants were collected, assayed for protein concentration by using the Bradford reagent, aliquoted and stored at -80°C until use for Western blot analyses.

Protein determination by Western Blotting

Equal amounts of protein (100 μg) were separated by SDS-PAGE and transferred to polyvinylidene difluoride (PVDF) filters (Protein Sequencing Membrane, BioRad). Membranes were probed with the corresponding primary antibody followed by incubation

with peroxide-conjugated antirabbit Ig (GE Healthcare, Madrid, Spain). Blots were developed with the ECL system (GE Healthcare, Madrid, Spain). Normalization of Western blot was ensured by β -actin and band quantification was carried out with a scanner and the Scion Image software.

Statistics

Prior to statistical analysis, data were tested for homogeneity of variances by the test of Levene. For multiple comparisons, one-way ANOVA was followed by the Bonferroni test when variances were homogeneous or by the Tamhane test when variances were not homogeneous. $P < 0.05$ was considered significant. A SPSS version 15.0 program was used.

Results

Body and liver weights and food consumption

Final body weights and food consumption decreased after DEN administration in all groups of animals (groups 3-5) (Table 2). Consistent with these results, body weight gains of groups 1 and 2 were higher than those in DEN-injected groups. The lowest values were observed in group 5 corresponding to DEN-treated animals that received the cocoa-rich diet at day 28, after being fed with standard diet (Table 2). Therefore, the food efficiency diminished in groups 3-5 when compared to DEN-untreated animals (groups 1 and 2), and the smallest alimentary efficiencies were observed in DEN-rats receiving the cocoa-rich diet (groups 4 and 5).

Liver weights of group 5 were diminished when compared to all other groups of animals. However, no significant differences in liver-to-body weight ratio were observed among the experimental groups (Table 2).

Levels of markers of liver function in serum

As shown in Table 3, the serum total protein levels were reduced in DEN-treated rats, and were not normalized by the cocoa supplementation; on the contrary, group 5, in which diet was changed from standard to cocoa-rich one, showed the lowest values (Table 3).

The serum activity of ALT exhibited an elevation in DEN-treated groups and remained increased in animals receiving cocoa-diet in comparison with untreated rats (Table 3).

Addition of cocoa to DEN-injected animals (groups 4 and 5) showed an enhanced activity of AST when compared to untreated (groups 1 and 2) and DEN-treated (group 3) rats. Similarly, ALP activity was increased because of the administration of DEN in all experimental groups

(groups 3-5). However, only in cocoa-fed rats treated with DEN (group 4), an increase in GST activity was observed (Table 3).

Levels of liver GSH and phase I and II enzymes

Liver GSH concentration decreased by DEN administration in group 3, which showed the lowest levels (Figure 2). However, liver GSH content was elevated by cocoa in control and DEN-treated rats (groups 2 and 4, Figure 2), whereas animals fed with cocoa-diet for 2 weeks (group 5) displayed similar values to control DEN-untreated rats (group 1).

Similar results were obtained for GPx activity, with increased values in the animals consuming cocoa for 6 weeks (groups 2 and 4), decreased GPx activity after DEN treatment in rats fed with standard diet (group 3), and recovering of control values in group 5. GR showed an increased activity in untreated animals fed cocoa (group 2), while all the DEN-treated groups showed similar values to controls (group 1) (Figure 2). Similarly to GPx, catalase activity decreased in animals fed with standard diet and treated with DEN (group 3), whereas the other groups of rats displayed similar levels to controls (Figure 2). GST activity in liver remained unaltered in all animals except in group 4, where an increase in its activity was detected (Figure 2). Therefore, cocoa induces the antioxidant defence in the liver of the control and DEN-injected rats.

Protein carbonyl content and LDH levels and caspase-3 activity in liver

Levels of protein carbonyl content were increased in group 3 due to DEN administration when compared to no-injected animals (Figure 3). However, cocoa-fed rats treated with DEN (groups 4 and 5) showed a decreased protein carbonyl levels when compared to group 3, values were reverted to control levels in group 4 (cocoa-fed during 4 weeks) or partly diminished in group 5 (cocoa-fed for 2 weeks).

Lactate dehydrogenase levels were increased by DEN administration in all groups (groups 3-5), pointing at a DEN-induced liver injury that was not prevented by the cocoa-rich diet (Figure 3). DEN also induced caspase-3 activity (group 3), but cocoa administration was capable of returning caspase-3 activity to control levels in animals consuming the cocoa-rich diet throughout the experiment (group 4), or partly decreased the DEN-induced activation of caspase-3 in animals receiving the cocoa-rich diet only for two weeks (group 5). Animals fed with the cocoa-rich diet (group 2) showed LDH levels similar to control rats and decreased values of caspase-3 activity (Figure 3). These results indicate that cocoa diminished the oxidative (carbonyl protein content) and apoptotic damages (caspase-3 activation), but not the necrotic DEN-induced liver injury (LDH activity).

Liver AKT, ERK and JNK levels

Figures 4 and 5 illustrate that DEN-treated animals fed with the standard diet (group 3) showed a significant increase in the levels of phosphorylated AKT and JNK proteins, whereas no changes in these two proteins were found in the other groups of animals. Phosphorylated ERK levels were increased by cocoa administration in control animals (group 2), but they were dramatically decreased by DEN treatment in all injected rats (groups 3-5) (Figure 5). There was no difference in the total levels of AKT, ERKs and JNKs among the groups. Thus, cocoa prevented the activation of AKT and JNK, although it did not modify the decreased values of active ERK induced by DEN.

Discussion

DEN toxicity is primarily associated to an excessive production of free radicals in the liver. As a consequence, reactive electrophilic intermediates are formed, which overwhelms the antioxidant defences and ultimately proceeds to oxidative stress paving way to liver damage (Kang, J. et al., 2007). DEN induces a postnecrotic hepatocellular proliferation that contributes to enhance the number of initiated cells (Cascales, M., 2001), and it is accepted as a model to study the relations among liver necrosis, cancer initiation and replication (Cascales, M., 2001, Chuang, S. et al., 2000, Köhle, C. et al., 2008, Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2005, Sreepriya, M. and Bali, G., 2006, Sundaresan, S. and Subramanian, P., 2008, Tessitore, L. and Bollito, E., 2006).

Cocoa and its derived products have been demonstrated to contain important antioxidant polyphenols that inhibit different tumoral processes and exhibit antioxidant and anti-inflammatory properties (Amin, I. et al., 2004, Lamuela-Raventós, R. et al., 2005, Martin, M. et al., 2009, Martin, M. et al., 2008, Yamagishi, M. et al., 2000, Yamagishi, M. et al., 2002, Yamagishi, M. et al., 2003). In this study, we show that a cocoa-rich diet partly protect the liver against an oxidative-induced damage by decreasing the protein carbonyl content and modulating the activities of antioxidant enzymes and by regulating key proteins of cell signalling cascades, although cocoa-rich diet was unable to abrogate the DEN-induced enhanced values of hepatospecific enzymes and LDH. The cocoa dose was finally selected based on our own previous in vivo studies, where it is showed that it contributed to reduce the cardiovascular risk acting as an effective antioxidant (Lecumberri, E. et al., 2007).

Reduction of food intake and consequently, the reduction of body weight gain observed in DEN-treated animals, could be largely due to losses from skeletal muscle and adipose tissue

as previously shown (Sreepriya, M. and Bali, G., 2005), and it could be considered as an indirect indication of the declining hepatic function following exposure to DEN (Amin, I. et al., 2004, Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2005). In addition, assessment of the liver/body weights ratio was used to investigate potential changes in the liver size, but no differences were found (Ramakrishnan, G. et al., 2006, Sreepriya, M. and Bali, G., 2005, Yamagishi, M. et al., 2003).

Low levels of albumin have been reported in the serum of patients and animals with early hepatocellular cancer (Sreepriya, M. and Bali, G., 2005, Yoshida, H. et al., 2005). Contrary to other antioxidants such as curcumin and embelin that reverted the diminished serum proteins values induced by DEN (Sreepriya, M. and Bali, G., 2005), our results indicate that cocoa did not exhibit its protective effects during DEN-induced liver damage by enhancing the total protein levels.

GSH is an important non-enzymatic antioxidant defence required to maintain the normal redox state of cells and to counteract deleterious effects of oxidative stress. GSH depletion ultimately promotes oxidative stress, with a cascade of effects thereby affecting functional and structural integrity of cell and organelle membrane (Masella, R. et al., 2005). Our results showed that the significant decrease in GSH levels in animals exposed to DEN was restored or enhanced by cocoa feeding, which indicated the potentiality of cocoa to counteract the oxidative damage induced by DEN and to reinforce the antioxidant defence in normal conditions, as previously reported for other natural antioxidants (Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2006).

Chemical induction of liver damage by DEN administration is associated with changes in the oxygen radical metabolism in this organ, which were demonstrated by measuring protein carbonyl content and the activity of the antioxidant enzymes. Generation of protein, DNA and lipid oxidation products by DEN administration at the initiation stage can be prevented by

natural antioxidants (Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2006). Thus, DEN-treated rats fed with cocoa (groups 4 and 5) displayed lower levels of protein carbonyl content when compared with rats injected with DEN receiving regular diet (group 3). This reveals the protective role of cocoa against the induced-oxidative stress (Lecumberri, E. et al., 2007, Martin, M. et al., 2008, Ramiro-Puig, E. et al., 2009).

Catalase converts H_2O_2 to H_2O and GPx catalyses the transformation of H_2O_2 to harmless byproducts. During H_2O_2 scavenging, GSH is oxidized to GSSG by GPx. The reduction of GSSG to GSH is catalysed by GR using NADPH as reducing potential. GPx and GR activities were increased and catalase activity was unchanged in control animals fed cocoa in agreement with our previous results, where the induction of both enzymatic activities by a cocoa polyphenolic extract (Martin, M. et al., 2009) and a cocoa fibre byproduct (Lecumberri, E. et al., 2007) was demonstrated. The decrease of GSH levels and GPx and catalase activities indicates the severity of the oxidative stress induced during the exposure to DEN. Interestingly, cocoa counteracted the hepatic oxidative damage by preventing the reduction of these parameters provoked by DEN. In line with this, decreases in GSH levels, as well as in activities of catalase, GPx and GR, have been reported in hepatic tumours (Ramakrishnan, G. et al., 2006, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2006).

GST catalyses the reaction of endogenous GSH with numerous electrophiles to yield less toxic conjugates that are easily eliminated (Masella, R. et al., 2005). Although increased and decreased liver and serum GST activities have been reported after DEN-induced damage (Amin, I. et al., 2004, Sivaramakrishnan, V. et al., 2008, Sreepriya, M. and Bali, G., 2006, Sundaresan, S. and Subramanian, P., 2008, Yadav, A. and Bhatnagar, D., 2007), we found that liver and serum GST activities increased in rats fed with cocoa and injected with DEN (group 4), suggesting that cocoa administration contributes to the liver protection against the

oxidative induced-injury. In this regard, it has been demonstrated that different natural compounds and polyphenols, as well a cocoa bean product, induce GST as one of the principal anticarcinogenic mechanisms (Amin, I. et al., 2004, Masella, R. et al., 2005, Sundaresan, S. and Subramanian, P., 2008). Moreover, some studies point to the induction of GST as a mechanism to protect against chemically induced cancer and oxidative stress by increasing the metabolism of electrophilic intermediates and ROS (Masella, R. et al., 2005). Enhancement of liver LDH activity, a sensitive marker of hepatocyte injury, indicates a nonspecific alteration in the plasma membrane integrity and permeability and/or may be due to its overproduction by tumour cells (Kamaraj, S. et al., 2007, Manso, M. et al., 2007, Sivaramakrishnan, V. et al., 2008). The significant DEN-induced increase in serum hepatic marker enzymes and liver LDH activities that indicated the hepatocellular dysfunction and the severity of the liver necrotic damage (Amin, I. et al., 2004, Kamaraj, S. et al., 2007, Sivaramakrishnan, V. et al., 2008), were not avoided by the administration of a cocoa-rich diet.

Activation of caspase-3, the most important enzyme responsible for apoptosis, has been considered as a sensitive method of detecting liver damage and has been associated with progressive liver fibrosis (Bantel, H. et al., 2004). Moreover, high rates of apoptosis have been demonstrated in liver cancer (Kang, J. et al., 2007, Schimtz, K.J. et al., 2007). In addition, it should be mentioned that administration of DEN (200 mg/Kg) causes fibrosis and necrosis in the liver (Tessitore, L. and Bollito, E., 2006). It is noticeable that DEN-injected rats fed with standard diet (group 3) showed levels of AST and ALT that were below those of the other DEN-treated groups. This finding could be a consequence of a reduction in the number of viable hepatocytes due to enhanced cell death in liver, as these animals also showed the highest levels of LDH and caspase-3 (Manso, M. et al., 2007). Thus, the decreased caspase-3 activity in animals not injected with DEN and fed with cocoa (group 2),

as well as in DEN-treated rats receiving cocoa (groups 4 and 5), suggested a potential protective effect of cocoa against cell death in normal and under oxidative conditions. Therefore, although additional studies are needed, it could be suggested that cocoa could partly protected hepatocytes against the progression of the fibrotic damage induced by DEN. The signalling mechanisms associated with liver damage induced by oxidative-stress are not completely known. In our study, the increase in p-AKT levels in DEN-treated rats fed the standard diet (group 3) agrees with that reported in animals receiving DEN in drinking water (Parekh, P. and Rao, K.V.K., 2007). These data support the important role of AKT in controlling the balance between survival and apoptosis (Ramos, S., 2008), which has been considered a critical factor in the aggressiveness of hepatocellular cancer (Parekh, P. and Rao, K.V.K., 2007, Schimtz, K.J. et al., 2007). Moreover, the value of p-AKT was restored in rats injected with DEN but fed cocoa (groups 4 and 5) suggests that cocoa prevents the activation of this main protein related to cell survival/proliferation; this could result in a potential attenuation of the postnecrotic proliferation induced by DEN and in a reduction of the number of initiated cells (Cascales, M., 2001).

ERK plays a critical role in controlling the balance between cell survival and proliferation, and cell cycle progression (Schimtz, K.J. et al., 2007). Although data on p-ERK values remain controversial (Ito, Y. et al., 1998, Parekh, P. and Rao, K.V.K., 2007), increased p-ERK levels have recently been related to an advanced, but not early, tumour stage (Huynh, H. et al., 2003). This finding supports our results, since decreased levels of p-ERK were found in all DEN-injected animals, a model to study the relations among necrosis and cancer initiation. Moreover, p-ERK was increased in rats fed a cocoa-rich diet indicating that cocoa components can activate cellular kinases (Ramos, S., 2008) in concert with our previous results (Granado-Serrano, A. et al., 2007, Martin, M. et al., 2009, Ramiro-Puig, E. et al., 2009). Similarly, the enhanced levels of p-ERK in cocoa fed rats (group 2) could be related to

their increased GR and GPx activities, since MAPKs pathway has recently been implicated in the up-regulation of several antioxidant enzymes activities in liver cancer cells (Martin, M. et al., 2009).

The JNK family belongs to the MAPKs superfamily (same as ERKs) and its activation has been associated to apoptosis in liver (Czaja, M., 2003, Parekh, P. and Rao, K.V.K., 2007). DEN administration enhanced the p-JNK/JNK ratio in animals fed with the standard diet (group 3) and returned to control values in DEN-groups receiving cocoa (groups 4 and 5). In line with this, a protective effect of cocoa flavonoids has been described on a model of oxidative stress in neurons, showing a down-regulation of p-JNK levels (Ramiro-Puig, E. et al., 2009). Moreover, DEN-treated rats fed with regular diet showed the highest LDH and caspase-3 activities and p-JNK/JNK values, pointing out to an enhanced cell death in the liver of these animals, although this group also exhibited enhanced p-AKT values suggestive of a promotion of cellular survival/proliferation. At the moment we cannot explain this finding and further studies are needed to elucidate this result, but similar effects have been reported in patients with liver cancer (Kang, J. et al., 2007, Schimtz, K.J. et al., 2007).

Finally, it is important to mention that in vitro studies have attributed the protective effect of cocoa to the polyphenols, since the contribution of theobromine was considered negligible (Martin, M. et al., 2008). Additionally, a similar protective effect has been recently described after treating cells with a cocoa phenolic extract or (-)-epicatechin (Ramiro-Puig, E. et al., 2009). Therefore, it could be suggested that the beneficial effects of cacao in the present study might be ascribed to the polyphenolic compounds, which even may develop synergism among them, but further studies are needed.

In summary, a cocoa-rich diet protects cells against DEN-induced oxidative stress by activating the antioxidant defence system. In addition, cocoa appears to exert an antiapoptotic effect in the liver of DEN-treated animals that could be associated with the prevention of JNK and

caspase-3 activation. Cocoa also prevented the DEN-induced increase of the survival/proliferation signal AKT, associated with poor prognosis in liver cancer. All these effects contribute to attenuate the liver DEN-induced oxidative damage. However, cocoa was not completely capable of avoiding the liver necrotic damage induced by the hepatotoxic as shown by the increased activities of the hepatospecific enzymes and LDH as well as decreased p-ERK values. These results provide new insights into the antioxidative mechanisms of cocoa flavonoids and point towards their antiapoptotic and modulatory effects on cell death and survival/proliferation pathways as additional mechanisms of action of these compounds. Therefore, it could be suggested that cocoa or cocoa products enriched in flavonoids, which lack of toxicity in humans, may contribute to the protection against liver oxidative stress-related diseases.

475 **Acknowledgements**

476

477 This work was supported by the grants 200870I198, AGL2004-302, AGL2007-64042 and
478 CSD2007-00063 from the Spanish Ministry of Science and Innovation. A.B. Granado-
479 Serrano is a predoctoral fellow of the Spanish Ministry of Science and Education.

480 The authors declare that there are no conflicts of interest.

481

References

- Aebi, H., 1987. Catalase. In *Methods of Enzymatic Analysis*; Bergmeyer, H. V. Ed.; Verlag-Chemie: Weinheim, Germany, 273-282.
- Alia, M., Horcajo, C., Bravo, L. and Goya, L., 2003. Effect of grape antioxidant dietary fiber on total antioxidant capacity and the activity of liver antioxidant enzymes in rats. *Nutr Res.* 23, 1251-1267.
- Alía, M., Ramos, S., Mateos, R., Granado-Serrano, A., Bravo, L. and Goya, L., 2006. Quercetin protects human hepatoma cell line (HepG2) against oxidative stress induced by tertbutyl hydroperoxide. *Toxicol Appl Pharmacol.* 212, 110-118.
- Amin, I., Koh, B. and Asmah, R., 2004. Effect of cacao liquor on tumor marker enzymes during chemical hepatocarcinogenesis in rats. *J Med Food.* 7, 7-12.
- Bantel, H., Lügering, A., Heideman, J., Volkmann, X., Poremba, C., Strassburg, C., Manns, M. and Schulze-Osthoff, K., 2004. Detection of apoptotic caspase activation in sera from patients with chronic HCV infection is associated with fibrotic liver injury. *Hepatology.* 40, 1078-1087.
- Bergmeyer, H. and Bernt, E., 1974. Lactic dehydrogenase. *Meth Enzym Anal.* 2 574-579.
- Cascales, M., 2001. *Mecanismos de hepatotoxicidad*. Madrid: Real Academia de Doctores.
- Czaja, M., 2003. The future of GI and liver research: Editorial perspectives III. JNK/AP-1 regulation of hepatocyte death. *Am J Physiol Gastrointest Liver Physiol.* 284, G875-G879.
- Chuang, S., Cheng, A., Lin, J. and Kuo, M., 2000. Inhibition by curcumin of diethylnitrosamine-induced hepatic hyperplasia, inflammation, cellular gene products and cell-cycle-related proteins in rats. *Food Chem Toxicol.* 38, 991-995.

- 506 Gavete, L., Martin, M., Alvarez, C. and Escriva, F., 2005. Maternal food restriction enhances
507 insulin-induced GLUT-4 translocation and insulin signaling pathway in skeletal muscle
508 from suckling rats. *Endocrinology*. 146, 3368-3378.
- 509 Goldberg, D. and Spooner, R., 1987. Glutathione reductase. In *Methods of Enzymatic*
510 *Analysis*; Bergmeyer, H. V. Ed.; Verlag-Chemie: Weinheim, Germany, 258-265.
- 511 Granado-Serrano, A., Martín, M., Izquierdo-Pulido, M., Goya, L., Bravo, L. and Ramos, S.,
512 2007. Molecular mechanisms of (–)-epicatechin and chlorogenic acid on the regulation of
513 the apoptotic and survival/proliferation pathways in a human hepatoma cell line. *J Agric*
514 *Food Chem*. 55, 2020-2027.
- 515 Gunzler, W., Kremers, H. and Flohe, L., 1974. An improved coupled test procedure for
516 glutathione peroxidase. *Klin Chem Klin Biochem*. 12, 444-448.
- 517 Hecht, S., 1997. Approaches to cancer prevention based on an understanding N-nitrosamine
518 carcinogenesis. *Proc Soc Exp Biol Med*. 216, 181-191.
- 519 Hissin, P. and Hilf, R., 1976. A fluorometric method for determination of oxidised and reduced
520 glutathione in tissues. *Anal Biochem*. 74, 214-26.
- 521 Holley, S.L., Fryer, A.A., Haycock, J.W., Grubb, S.E.W., Strange, R.C. and Hoban, P.R.,
522 2007. Differential effects of glutathione S-transferase pi (GSTP1) haplotypes on cell
523 proliferation and apoptosis. *Carcinogenesis*. 28, 2268-2273.
- 524 Horder, M. and Rej, R., 1984. Alanine aminotransferase. L-alanine: 2-oxoglutarate
525 aminotransferase, EC 2.6.1.2. routine UV method. In *Methods of Enzymatic Analysis*;
526 Bergmeyer, H. V. Ed.; Verlag-Chemie: Weinheim, Germany, 444-450.
- 527 Huynh, H., Nguyen, T.T.T., Chow, K.H.P., Tan, P.H., Soo, K.C. and Tran, E., 2003. Over-
528 expression of the mitogen-activated protein kinase (MAPK) kinase (MEK)-MAPK in
529 hepatocellular carcinoma: its role in tumor progression and apoptosis. *BMC*
530 *Gastroenterol*. 3, 19.

- 531 Ito, Y., Sasaki, Y., Horimoto, M., Wada, S., Tanaka, Y., Kasahara, A., Ueki, T., Hirano, T.,
532 Yamamoto, H., Fujimoto, J., Okamoto, E., Hayashi, N. and Hori, M., 1998. Activation of
533 mitogen-activated protein kinases/extracellular signal-regulated kinases in human
534 hepatocellular carcinoma. *Hepatology*. 27, 951-958.
- 535 Jaeschke, H., Gores, G., Cederbaum, A., Hinson, J., Pessayre, D. and Lemasters, J., 2002.
536 Forum mechanisms of hepatotoxicity. *Toxicol Sci*. 65, 166-176.
- 537 Kamaraj, S., Vinodhkumar, R., Anandakumar, P., Jagan, S., Ramakrishnan, G. and Devaki, T.,
538 2007. The effects of quercetin on antioxidant status and tumor markers in the lung and
539 serum of mice treated with benzo(a)pyrene. *Biol Pharm Bull*. 30, 2268-2273.
- 540 Kang, J., Wanibuchi, H., Morimura, K., Gonzalez, F. and Fukushima, S., 2007. Role of
541 CYP2E1 in diethylnitrosamine-induced hepatocarcinogenesis *in vivo*. *Cancer Res*. 67,
542 11141-11146.
- 543 Köhle, C., Schwarz, M. and Bock, K., 2008. Promotion of hepatocarcinogenesis in humans and
544 animal model. *Arch Toxicol*. 82, 623-631.
- 545 Lamuela-Raventós, R., Romero-Pérez, A., Andrés-Lacueva, C. and Tornero, A., 2005. Review:
546 health effects of cocoa flavonoids. *Food Sci Technol Int*. 11, 159-176.
- 547 Lecumberri, E., Goya, L., Mateos, R., Alía, M., Ramos, S., Izquierdo-Pulido, M. and Bravo,
548 L., 2007. A diet rich in dietary fiber from cocoa improves lipid profile and reduces
549 malondialdehyde in hypercholesterolemic rats. *Nutrition*. 23, 332-341.
- 550 Manso, M., Ramudo, L. and De Dios, I., 2007. Extrapaneatic organ impairment during acute
551 pancreatitis induced by bile-pancreatic duct obstruction. Effect of N-acetylcysteine. *Int J*
552 *Exp Path*. 88, 343-349.
- 553 Martin, M., Granado-Serrano, A., Ramos, S., Izquierdo-Pulido, M., Bravo, L. and Goya, L.,
554 2009. Cocoa flavonoids up-regulate antioxidant enzyme activity via the ERK1/2 pathway

- 555 to protect against oxidative stress-induced apoptosis in HepG2 cells. J Nutr Biochem.
556 doi:10.1016/j.jnutbio.2008.10.009,
- 557 Martin, M., Ramos, S., Mateos, R., Granado-Serrano, A., Izquierdo-Pulido, M., Bravo, L. and
558 Goya, L., 2008. Protection of human HepG2 cells against oxidative stress by cocoa
559 phenolic extract. J Agric Food Chem. 56, 7765-7772.
- 560 Masella, R., Di Benedetto, R., Vari, R., Filesi, C. and Giovannini, C., 2005. Novel mechanisms
561 of natural antioxidant compounds in biological systems: involvement of glutathione-
562 related enzymes. J Nutr Biochem. 16, 577-586.
- 563 Mates, J.M., Segura, J.A., Alfonso, J.F. and Marquez, J., 2008. Intracellular redox status and
564 oxidative stress: implications for cell proliferation, apoptosis and carcinogenesis. Arch
565 Toxicol. 82, 273-299.
- 566 Na, H. and Surh, Y., 2008. Modulation of Nrf2-mediated antioxidant and detoxifying enzyme
567 induction by the green tea polyphenol EGCG. Food Chem Toxicol. 46, 1271-1278.
- 568 Parekh, P. and Rao, K.V.K., 2007. Overexpression of cyclin D1 is associated with elevated
569 levels of MAP kinases, Akt and Pak1 during diethylnitrosamine-induced progressive
570 liver carcinogenesis. Cell Biol Int. 31, 35-43.
- 571 Principato, G.B., Aisa, M.C., Talesa, V., Rosi, G. and Giovanini, E., 1985. Characterization of
572 the soluble alkaline phosphatase from hepatopancreas of *Squilla mantis* L. Comp
573 Biochem Physiol. 80B, 801-804.
- 574 Ramakrishnan, G., Rao, H., Raghavendran, B., Vinodhkumar, R. and Devaki, T., 2006.
575 Suppression of *N*-nitrosodiethylamine induced hepatocarcinogenesis by silymarin in rats.
576 Chem Biol Interact. 161, 104-114.
- 577 Ramiro-Puig, E., Casadesus, G., Lee, H., Zhu, X., McShea, A., Perry, G., Perez-Cano, F.,
578 Smith, M. and Castell, M., 2009. Neuroprotective effect of cocoa flavonoids on in vitro
579 oxidative stress. Eur J Nutr. 48, 54-61.

- 580 Ramiro Puig, E., Urpí-Sardá, M., Pérez-Cano, F., Franch, A., Castellote, C., Andrés-Lacueva,
581 C., Izquierdo-Pulido, M. and Castell, M., 2007. Cocoa-enriched diet enhances antioxidant
582 enzyme activity and modulates lymphocyte composition in thymus from young rats. J
583 Agric Food Chem. 55, 6431-6438.
- 584 Ramos, S., 2007. Effects of dietary flavonoids on apoptotic pathways related to cancer
585 chemoprevention. J Nutr Biochem. 18, 427-442.
- 586 Ramos, S., 2008. Cancer chemoprevention and chemotherapy: Dietary polyphenols and
587 signalling pathways. Mol Nutr Food Res. 52, 507-526.
- 588 Rej, R. and Horder, M., 1984. Aspartate aminotransferase. L-aspartate: 2-oxoglutarate
589 aminotransferase, EC 2.6.2.1. routine UV method. In Methods of Enzymatic Analysis;
590 Bergmeyer, H. V. Ed.; Verlag-Chemie: Weinheim, Germany, 416-424.
- 591 Richert, S., Wehr, N., Stadtman, E. and Levine, R., 2002. Assessment of skin carbonyl content
592 as a noninvasive measure of biological age. Arch Biochem Biophys. 397, 430-432.
- 593 Schimtz, K.J., Wohlschlaeger, J., Lang, H., Sotiropoulos, G.C., Malago, M., Steveling, K.,
594 Reis, H., Cicinnati, V.R., Schmid, K.W. and Baba, H.A., 2007. Activation of the ERK
595 and AKT signaling pathway predicts poor prognosis in hepatocellular carcinoma and ERK
596 activation in cancer tissue is associated with hepatitis virus infection. J Hepatol. 48, 83-
597 90.
- 598 Sen, N., Seaman, S. and McPherson, M., 1980. Further studies on the occurrence of volatile
599 and non-volatile nitrosamines in food. IARC Sci Publ. 457-460.
- 600 Sivaramakrishnan, V., Shilpa, P., Kumar, V. and Devaraj, S., 2008. Attenuation of *N*-
601 nitrosodiethylamine-induced hepatocellular carcinogenesis by a novel flavonol-Morin.
602 Chem Biol Interact. 171, 79-88.

- 603 Sreepriya, M. and Bali, G., 2005. Chemopreventive effects of embelin and curcumin against *N*-
604 nitrosodiethylamine/phenobarbital-induced hepatocarcinogenesis in Wistar rats.
605 *Fitoterapia*. 76, 549-555.
- 606 Sreepriya, M. and Bali, G., 2006. Effects of administration of embelin and curcumin on lipid
607 peroxidation, hepatic glutathione antioxidant defense and hematopoietic system during
608 *N*-nitrosodiethylamine/phenobarbital-induced hepatocarcinogenesis in Wistar rats. *Mol*
609 *Cell Biochem*. 284, 49-55.
- 610 Sundaresan, S. and Subramanian, P., 2008. Prevention of *N*-nitrosodiethylamine-induced
611 hepatocarcinogenesis by S-allylcysteine. *Mol Cell Biochem*. 310, 209-214.
- 612 Tessitore, L. and Bollito, E., 2006. Early induction of TGF- β 1 through a fasting–re-feeding
613 regimen promotes liver carcinogenesis by a sub-initiating dose of diethylnitrosamine.
614 *Cell Prolif*. 39, 105-116.
- 615 Thorgeirsson, S. and Grisham, J., 2002. Molecular pathogenesis of human hepatocellular
616 carcinoma. *Nat Genetics*. 31, 339-346.
- 617 Vinson, J., Proch, J., Bose, P., Muchler, S., Taffera, P., Shutta, D., Samman, N. and Agbor, G.,
618 2006. Chocolate is a powerful ex vivo and in vitro antioxidant, antiatherosclerotic agent
619 in an animal model, and a significant contributor to antioxidants in the European and
620 American diets. *J Agric Food Chem*. 54, 8071-8076.
- 621 Vitaglione, P., Morisco, F., Caporaso, N. and Fogliano, V., 2004. Dietary antioxidant
622 compounds and liver health. *Crit Rev Food Sci Nutr*. 44, 575-586.
- 623 Yadav, A. and Bhatnagar, D., 2007. Chemo-preventive effect of Star anise in *N*-
624 nitrosodiethylamine initiated and phenobarbital promoted hepato-carcinogenesis. *Chem*
625 *Biol Interact*. 169, 207-214.

- 626 Yamagishi, M., Natsume, M. and Nagaki, A., 2000. Antimutagenic activity of cacao: inhibitory
627 effect of cacao liquor polyphenols on the mutagenic action of heterocyclic amines. J
628 Agric Food Chem. 48, 5074-5078.
- 629 Yamagishi, M., Natsume, M., Osakabe, N., Nakamura, H., Furukawa, F., Imazawa, T.,
630 Nishikawa, A. and Hirose, M., 2002. Effects of cacao liquor proanthocyanidins on PhIP-
631 induced mutagenesis in vitro, and in vivo mammary and pancreatic tumorigenesis in
632 female Sprague-Dawley rats. Cancer Lett. 185, 123-130.
- 633 Yamagishi, M., Natsume, M., Osakabe, N., Okazaki, K., Furukawa, F., Imazawa, T.,
634 Nishikawa, A. and Hirose, M., 2003. Chemoprevention of lung carcinogenesis by cacao
635 liquor proanthocyanidins in a male rat multi-organ carcinogenesis model. Cancer Lett.
636 191, 49-57.
- 637 Yoshida, H., Yoshida, H., Shiina, S. and Omata, M., 2005. Early liver cancer: concepts,
638 diagnosis, and management. Int J Clin Oncol. 10, 384-390.
- 639

Figure Captions

Figure 1. Schematic representation of the experimental design used for the study.

Figure 2. Effect of cocoa on (A) glutathione levels and activity of (B) GPx, (C) GR, (D) CAT and (E) GST liver from control and DEN groups fed with standard (C) and cocoa-rich (Ca) diets. Data represent the means $\pm$ SD (n=8-10). Means without a common letter differ, $P < 0.05$.

Figure 3. Liver (A) protein carbonyl, (B) LDH and (C) caspase-3 levels in control and DEN groups fed with standard (C) and cocoa-rich (Ca) diets. Data represent the means $\pm$ SD (n=6-8). Means without a common letter differ, $P < 0.05$.

Figure 4. Effect of cocoa on levels of phosphorylated AKT (Ser473) and total AKT in controls and DEN-treated rats. (A) Bands of a representative experiment. (B) Percentage values of the p-AKT/AKT ratio relative to the control condition (means $\pm$ SD, n=6). Normalization of Western blots was ensured by β -actin. Means without a common letter differ, $P < 0.05$.

Figure 5. Effects of cocoa on the basal levels of phosphorylated ERK1/2 (Thr202/Tyr204), total ERK1/2, phosphorylated JNK1/2 (Thr183/Tyr185) and total JNK in controls and DEN-treated animals. (A) Representative blots of both MAPKs. Percentage data of (B) p-ERK/ERK and (C) p-JNK/JNK ratios relative to controls (means $\pm$ SD, n=7). The same liver homogenates were subjected to Western blot analysis using the corresponding non-phospho-

Protective effects of cocoa in liver damage

664 specific antibodies to detect total ERK or JNK. Equal loading of Western blots was ensured
665 by β -actin. Means without a common letter differ, $P < 0.05$.
666

667 Table 1. Composition of the experimental control (C) and cocoa-rich (Ca) diets (g/Kg
 668 dry weight).

	C	Ca
Casein	140	140
Dextrose	155	155
Sucrose	100	100
Fat	40	40
t-BHQ ¹	0.008	0.008
Mineral mix.	35	35
Vitamin mix.	10	10
L-Cys	1.8	1.8
Cholin bitartrate	2.5	2.5
Cellulose	100	45.6
Starch	415.7	310.1
Cocoa powder	-	160

669

670 ¹ *tert*-butylhydroquinone

671

672 Table 2. Body weight data, food intake and liver weight of rats in control and DEN groups fed with standard (C) and cocoa-rich (Ca) diets*.

	<i>Control groups</i>		<i>DEN groups</i>		
	1 (C)	2 (Ca)	3 (C)	4(Ca)	5(C-Ca)
Initial body weight (g)	95.00 ± 4.69 ^a	92.57 ± 3.41 ^a	102.75 ± 8.87 ^a	104.00 ± 7.77 ^a	105.88 ± 5.26 ^a
Body weight on day 14 (g)	135.17 ± 3.81 ^a	134.00 ± 7.83 ^a	135.00 ± 5.58 ^a	134.33 ± 6.68 ^a	137.75 ± 7.89 ^a
Final body weight (g)	205.83 ± 10.21 ^a	202.14 ± 6.36 ^a	183.17 ± 8.18 ^b	175.71 ± 12.09 ^{bc}	161.29 ± 10.81 ^c
Body weight gain (g in 42d)	110.83 ± 7.92 ^a	110.57 ± 4.58 ^a	81.20 ± 8.32 ^b	71.71 ± 16.98 ^{bc}	57.67 ± 17.15 ^c
Food Intake (g in 42d)	574.58 ± 3.66 ^a	574.40 ± 6.31 ^a	529.58 ± 25.66 ^b	534.91 ± 10.78 ^b	531.91 ± 16.53 ^b
Food Efficiency (Body weight gain/Food intake)	0.19 ± 0.02 ^a	0.19 ± 0.01 ^a	0.16 ± 0.02 ^b	0.13 ± 0.02 ^c	0.11 ± 0.03 ^c
Liver weight (g)	4.32 ± 0.28 ^a	4.42 ± 0.28 ^a	4.32 ± 0.17 ^a	4.20 ± 0.22 ^{ab}	3.79 ± 0.14 ^b
Liver-to body weight ratio	0.021 ± 0.002 ^a	0.022 ± 0.002 ^a	0.022 ± 0.001 ^a	0.024 ± 0.002 ^a	0.023 ± 0.002 ^a

685 * Data represent the means ± SD. Means in a row without a common letter differ, $P < 0.05$.

686 Table 3. Serum levels of total protein and activity of hepatic enzymes in control and DEN groups fed with standard (C) and cocoa-rich (Ca) diets*.

687

	<i>Control groups</i>		<i>DEN groups</i>		
	1 (C)	2 (Ca)	3 (C)	4 (Ca)	5 (C-Ca)
Total protein (mg/mL)	112.09 ± 11.61 ^a	119.09 ± 12.87 ^a	80.16 ± 11.65 ^b	80.96 ± 10.43 ^b	76.35 ± 16.55 ^b
ALT (U/L)	51.50 ± 10.67 ^a	59.29 ± 12.47 ^{ab}	75.50 ± 12.33 ^{bc}	85.00 ± 6.73 ^{cd}	86.75 ± 8.54 ^d
AST (U/L)	284.80 ± 41.84 ^a	278.17 ± 32.18 ^a	265.17 ± 28.99 ^a	479.67 ± 83.46 ^b	435.33 ± 67.33 ^b
ALP (U/L)	202.00 ± 20.82 ^a	221.00 ± 25.91 ^a	338.33 ± 33.43 ^b	315.00 ± 47.93 ^b	342.20 ± 54.67 ^b
GST (mU/mg protein)	0.58 ± 0.07 ^a	0.46 ± 0.08 ^a	0.56 ± 0.09 ^{ab}	0.71 ± 0.07 ^b	0.49 ± 0.09 ^a

695

696 * Data represent the means ± SD. Means in a row without a common letter differ, $P < 0.05$.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.