

# *Thermosinus carboxydivorans* gen. nov., sp. nov., a new anaerobic, thermophilic, carbon-monoxide-oxidizing, hydrogenogenic bacterium from a hot pool of Yellowstone National Park

Tatyana G. Sokolova,<sup>1</sup> Juan M. González,<sup>2†</sup>  
Nadezhda A. Kostrikina,<sup>1</sup> Nikolai A. Chernyh,<sup>1</sup>  
Tatiana V. Slepova,<sup>1</sup> Elizaveta A. Bonch-Osmolovskaya<sup>1</sup>  
and Frank T. Robb<sup>2</sup>

Correspondence  
Tatyana G. Sokolova  
tatso@mail.ru

<sup>1</sup>Institute of Microbiology, Russian Academy of Sciences, Prospect 60 Let Oktyabrya, 7/2,  
117811 Moscow, Russia

<sup>2</sup>COMB, Columbus Center, 701 E. Pratt St, Baltimore, MD 21202, USA

A new anaerobic, thermophilic, facultatively carboxydrophic bacterium, strain Nor1<sup>T</sup>, was isolated from a hot spring at Norris Basin, Yellowstone National Park. Cells of strain Nor1<sup>T</sup> were curved motile rods with a length of 2·6–3 µm, a width of about 0·5 µm and lateral flagellation. The cell wall structure was of the Gram-negative type. Strain Nor1<sup>T</sup> was thermophilic (temperature range for growth was 40–68 °C, with an optimum at 60 °C) and neutrophilic (pH range for growth was 6·5–7·6, with an optimum at 6·8–7·0). It grew chemolithotrophically on CO (generation time, 1·15 h), producing equimolar quantities of H<sub>2</sub> and CO<sub>2</sub> according to the equation CO + H<sub>2</sub>O → CO<sub>2</sub> + H<sub>2</sub>. During growth on CO in the presence of ferric citrate or amorphous ferric iron oxide, strain Nor1<sup>T</sup> reduced ferric iron but produced H<sub>2</sub> and CO<sub>2</sub> at a ratio close to 1 : 1, and growth stimulation was slight. Growth on CO in the presence of sodium selenite was accompanied by precipitation of elemental selenium. Elemental sulfur, thiosulfate, sulfate and nitrate did not stimulate growth of strain Nor1<sup>T</sup> on CO and none of these chemicals was reduced. Strain Nor1<sup>T</sup> was able to grow on glucose, sucrose, lactose, arabinose, maltose, fructose, xylose and pyruvate, but not on cellobiose, galactose, peptone, yeast extract, lactate, acetate, formate, ethanol, methanol or sodium citrate. During glucose fermentation, acetate, H<sub>2</sub> and CO<sub>2</sub> were produced. Thiosulfate was found to enhance the growth rate and cell yield of strain Nor1<sup>T</sup> when it was grown on glucose, sucrose or lactose; in this case, acetate, H<sub>2</sub>S and CO<sub>2</sub> were produced. In the presence of thiosulfate or ferric iron, strain Nor1<sup>T</sup> was also able to grow on yeast extract. Lactate, acetate, formate and H<sub>2</sub> were not utilized either in the absence or in the presence of ferric iron, thiosulfate, sulfate, sulfite, elemental sulfur or nitrate. Growth was completely inhibited by penicillin, ampicillin, streptomycin, kanamycin and neomycin. The DNA G+C content of the strain was 51·7 ± 1 mol%. Analysis of the 16S rRNA gene sequence revealed that strain Nor1<sup>T</sup> belongs to the *Bacillus–Clostridium* phylum of the Gram-positive bacteria. On the basis of the studied phenotypic and phylogenetic features, we propose that strain Nor1<sup>T</sup> be assigned to a new genus, *Thermosinus* gen. nov. The type species is *Thermosinus carboxydivorans* sp. nov. (type strain, Nor1<sup>T</sup> = DSM 14886<sup>T</sup> = VKM B-2281<sup>T</sup>).

## INTRODUCTION

Published online ahead of print on 9 July 2004 as DOI 10.1099/ijs.0.63186-0.

†Present address: Instituto de Recursos Naturales y Agrobiología, CSIC, PO Box 1052, 41080 Sevilla, Spain.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain Nor1<sup>T</sup> is AY519200.

Several phylogenetically diverse thermophilic prokaryotes perform the metabolic reaction CO + H<sub>2</sub>O → CO<sub>2</sub> + H<sub>2</sub> ( $\Delta G^0 = -20$  kJ). These are representatives of the bacterial genera *Carboxydothermus* (Svetlichny *et al.*, 1991, 1994), *Caldanaerobacter* (formerly *Carboxydibrachium pacificum*)

(Sokolova *et al.*, 2001; Fardeau *et al.*, 2004) and *Carboxydocella* (Sokolova *et al.*, 2002). Recently, a hyperthermophilic archaeon of the genus *Thermococcus* capable of growth at the expense of the same reaction was isolated from deep-sea hydrothermal vents (Sokolova *et al.*, 2004). The metabolism of *Carboxydothermus hydrogenoformans* has been studied at the enzymic level (Svetlitchny *et al.*, 2001; Dobbek *et al.*, 2001). This physiological group of prokaryotes has been proposed to be named 'hydrogenogens' (Svetlitchny *et al.*, 2001). CO-oxidizing hydrogenogenic prokaryotes were shown to possess various metabolic capacities. Growth of *Carboxydothermus thermautotrophica* was found to be obligately dependent on CO. Apart from growth on CO, *Carboxydothermus hydrogenoformans* is able to grow on pyruvate (Pusheva & Sokolova, 1995). *Caldanaerobacter subterraneus* subsp. *pacificus*, isolated from a deep-sea hydrothermal vent in the Okinawa Trough, besides having the capacity for anaerobic CO oxidation/H<sub>2</sub> formation, is capable of growing on several fermentable substrates (Sokolova *et al.*, 2001). *Thermococcus* strain AM4 grows on CO producing H<sub>2</sub>, or chemo-organotrophically with elemental sulfur (Sokolova *et al.*, 2004). Herein, we report the isolation of a novel anaerobic, CO-utilizing, H<sub>2</sub>-producing, thermophilic bacterium capable of iron reduction during growth on CO.

## METHODS

**Collection of samples.** A sample of mud and water was taken from a small pool in the neutral (wooded) part of Norris Basin in the Yellowstone National Park. Organic matter (rotting wood, scum) was present in the pool. The geographical coordinates of the sampling site were 44° 43·797 N, 110° 42·506 W. The sample temperature was 50 °C and its pH was 7·5 (at 50 °C).

**Culture conditions and strains.** Enrichment and isolation of anaerobic carboxydrophic bacteria were carried out on Medium 1 supplemented with a neutralized solution of ferric citrate or amorphous ferric iron oxide. Medium 1 was of the following composition (g l<sup>-1</sup>): NH<sub>4</sub>Cl, 1; MgCl<sub>2</sub>.2H<sub>2</sub>O, 0·33; CaCl<sub>2</sub>.6H<sub>2</sub>O, 0·1; KCl, 0·33; KH<sub>2</sub>PO<sub>4</sub>, 0·5; 1 ml of trace element solution (Kevbrin & Zavarzin, 1992); 1 ml of vitamin solution (Wolin *et al.*, 1963). After boiling, the medium was cooled under an N<sub>2</sub> atmosphere. Yeast extract (0·2 g l<sup>-1</sup>) and NaHCO<sub>3</sub> (0·5 g l<sup>-1</sup>) were added afterwards, and the pH was adjusted to 6·8–7·0 with 6 M HCl. A neutralized solution of ferric citrate or amorphous ferric iron oxide was added to a final concentration of 20 or 90 mM, respectively. Amorphous ferric iron oxide was prepared by titrating a solution of FeCl<sub>3</sub> with 10% NaOH to pH 9. Aliquots (10 ml) of the medium were placed in 50 ml bottles, and the head-space was filled with CO (100 kPa). Pure cultures were obtained from colonies on the same medium solidified with 5% agar in roll-tubes under CO in the gas phase. Growth of pure cultures and physiological tests were performed using Medium 2. Medium 2 had the same composition as Medium 1 except it was supplemented with Na<sub>2</sub>S.9H<sub>2</sub>O (0·5 g l<sup>-1</sup>).

The reference strains used in this study were *Carboxydothermus hydrogenoformans* Z-2901<sup>T</sup> (DSM 6008<sup>T</sup>) (Svetlitchny *et al.*, 1991), *Carboxydothermus thermautotrophica* 41<sup>T</sup> (DSM 12326<sup>T</sup>) (Sokolova *et al.*, 2002) and *Caldanaerobacter subterraneus* subsp. *pacificus* JM<sup>T</sup> (DSM 12653<sup>T</sup>) (Sokolova *et al.*, 2001; Fardeau *et al.*, 2004).

**Light and electron microscopy.** Light microscopy was carried

out using a phase-contrast microscope with a 90/1·25 oil immersion objective. Specimens of whole cells for electron microscopy were negatively stained with 2% phosphotungstic acid. For the preparation of thin sections, cells were fixed with 5% glutaraldehyde for 2 h and 1% OsO<sub>4</sub> for 4 h at 4 °C and then embedded in Epon-812. The thin sections were stained with uranyl acetate and lead citrate. Electron micrographs were taken with a JEM-100C electron microscope.

**Physiological studies.** To test the growth of the novel isolate with various substrates and electron acceptors, Medium 2 with 100% N<sub>2</sub> in the gas phase was used, supplemented with the following substrates (2 g l<sup>-1</sup>): peptone, yeast extract, sucrose, lactose, glucose, galactose, ethanol, methanol, sodium salts of citrate, acetate, formate or pyruvate. Possible electron acceptors – elemental sulfur (10 g l<sup>-1</sup>) and sodium salts of nitrate, sulfate, thiosulfate (2 g l<sup>-1</sup>), sulfite (2 mM) or selenate (2 mM) – were tested on Medium 2 with various growth substrates. Growth with amorphous ferric iron oxide (90 mM) or selenite (2 mM) on various substrates was tested in Medium 1.

Bacterial growth was determined by direct cell count under a phase-contrast microscope.

Determinations of CO, gaseous products of metabolism, short-chain organic acids and alcohols were performed by GC as described previously (Sokolova *et al.*, 2001). H<sub>2</sub>S was determined by colorimetric reaction (Trüper & Schlegel, 1964). Fe(III) reduction was determined by measuring the accumulation of Fe(II) in the growth medium. For that, a 0·5 ml sample was added to 5 ml of 0·6 M HCl and, after a 24 h extraction, HCl-soluble Fe(II) was determined by the reaction with 2,2'-dipyridyl (Balashova & Zavarzin, 1980).

**DNA isolation and base composition.** DNA was prepared as described by Marmur (1961). The DNA G+C content was determined by melting-point analysis (Marmur & Doty, 1962) using *Escherichia coli* K-12 DNA as a reference.

**rRNA gene sequence.** The 16S rRNA gene sequence was PCR-amplified by using the primer pair 519F (5'-GTT TCA GCM GCC GCG GTA ATW C-3') and 1522R (5'-AAG GAG GTG ATC CAG CCG CA-3'). The amplified DNA fragment was purified using the Qiagen PCR purification kit (Qiagen) and sequenced by the dideoxynucleotide chain-termination method on an ABI 373A sequencer (Applied Biosystems). Sequencing was performed using the primers 519F and 341F (5'-CC TAC GGG AGG CAG CAG-3'), forward strand, and primers 907R (5'-CCC CGT CAA TTC ATT TGA GTT T-3') and 1522R, reverse strand. Sequence alignment was performed using the software suite ARB (Ludwig *et al.*, 2004). The alignment was edited manually considering the expected sequence secondary structure. An unrooted phylogenetic tree was constructed by maximum-likelihood using the program FastDNAMl (Felsenstein, 1981) embedded in ARB. The obtained tree topology was reconstructed by quartet-puzzling using the program Treepuzzle (Strimmer & von Haeseler, 1996) also available in the ARB package. The quartet-puzzling tree represented a consensus tree showing well supported branching. It was based on 1000 puzzling trials. The reliability value of each internal branch indicates as a percentage how often the corresponding cluster was found. The GenBank/EMBL/DDBJ accession numbers of the 16S rRNA sequences used in this study are given in Fig. 4.

## RESULTS AND DISCUSSION


### Enrichment and isolation

For the enrichment of anaerobic, thermophilic, CO-oxidizing bacteria, 100 ml serum bottles containing 20 ml Medium 1 with ferric citrate and CO as the gas phase were

inoculated with about 1 g of sample. After 3 days incubation at 55 °C, the gas pressure in several bottles increased from 100 to 120–150 kPa. Growth of curved rod-shaped cells was observed. The CO content in the gas phase decreased; the resulting gas phase composition was about 25–30% CO, 30% H<sub>2</sub> and 30% CO<sub>2</sub>. The colour of the medium changed from yellow–brown to green, indicating the reduction of Fe(III) to Fe(II). When transferred to Medium 2, the culture retained the ability to grow by oxidation of CO to CO<sub>2</sub> and production of H<sub>2</sub> and had the same morphology. After seven passages performed by serial tenfold dilutions, the culture was transferred to solid medium in roll-tubes filled with CO. After 4 days incubation at 55–60 °C of tubes inoculated with aliquots from the 10<sup>6</sup> and 10<sup>7</sup> dilutions, round white colonies of about 0·5 mm in diameter developed. Several colonies were isolated and transferred to Medium 2 under 100% CO. From all the colonies the growth of motile curved rods was obtained. One isolate, designated Nor1<sup>T</sup>, was chosen for further characterization.

### Morphology

Cells of strain Nor1<sup>T</sup> were curved rods with a length of 2·6–3 µm and a width of about 0·5 µm, arranged singly or in pairs (Fig. 1a, b). Cells were motile due to lateral flagella (Fig. 1a). Electron microscopy of ultrathin sections revealed a Gram-negative cell wall structure (Fig. 1c). The outer membrane had a folded structure (Fig. 1b, c). Cells divided by binary transverse fission (Fig. 1b, c).



**Fig. 1.** Electron micrographs of cells of strain Nor1<sup>T</sup>. Thin sections (b, c) and negative staining of whole cells (a). Bars, 0·5 µm.

### Growth parameters


Strain Nor1<sup>T</sup> grew between 40 and 68 °C, with an optimum at 60 °C. No growth occurred at 37 or 70 °C. Growth was possible between pH 6·5 and 7·6; no growth was detected at pH 6·2 or 7·8. Optimum pH for growth was 6·8–7·0.

### Physiology of growth


Strain Nor1<sup>T</sup> was able to grow on Medium 1 with or without ferric iron with 100% CO in the gas phase, as well as on Medium 2 with 100% CO. CO oxidation was coupled with H<sub>2</sub> and CO<sub>2</sub> formation in equimolar quantities according to the equation CO + H<sub>2</sub>O → CO<sub>2</sub> + H<sub>2</sub> (Fig. 2 and Fig. 3). No methane, acetate or any other metabolic products were produced. The generation time of strain Nor1<sup>T</sup> grown on CO without ferric iron was 1·15 h. During the growth on CO in Medium 1 in the presence of ferric citrate or amorphous ferric iron oxide, strain Nor1<sup>T</sup> reduced ferric iron (Fig. 3); however, the amount of ferric iron reduced was not large enough to shift significantly the ratio of H<sub>2</sub> and CO<sub>2</sub> produced. The generation time of strain Nor1<sup>T</sup> during growth on CO in the presence of ferric citrate was 1·07 h. The isolate grew in Medium 1 on CO in the presence of selenite, reducing it to elemental selenium, visible as red precipitate, but producing H<sub>2</sub> and CO<sub>2</sub> in nearly equimolar quantities. The isolate did not grow under an H<sub>2</sub>/CO<sub>2</sub> (4:1) mixture either in Medium 2 or in Medium 1 supplemented with ferric citrate or amorphous ferric iron oxide.

Strain Nor1<sup>T</sup> was found to be an obligate anaerobe. It did not grow under mixtures of CO and air, which contained 0·5, 1·0, 1·5, 2, 5 or 10% molecular oxygen.

On Medium 2 (reduced with sodium sulfide), strain Nor1<sup>T</sup> was capable of growth with glucose, sucrose, lactose,



**Fig. 2.** Growth of strain Nor1<sup>T</sup> at 60 °C in Medium 2 supplemented with 200 mg yeast extract l<sup>-1</sup> under an atmosphere of CO: ○, cell number; ▲, CO consumption; ■, H<sub>2</sub> production. CO and H<sub>2</sub> are shown as their quantities in the gas phase per 1 ml of liquid culture.



**Fig. 3.** Growth of strain *Nor1<sup>T</sup>* at 60 °C in Medium 1 supplemented with 20 mM ferric citrate and 200 mg yeast extract l<sup>-1</sup> under an atmosphere of CO: ○, cell number; ▲, CO consumption; ■, H<sub>2</sub> production; ◇, ferrous iron production. CO and H<sub>2</sub> are shown as their quantities in the gas phase per 1 ml of liquid culture.

arabinose, maltose, fructose, xylose and pyruvate, but not with cellobiose, galactose, peptone, yeast extract, lactate, acetate, formate, ethanol, methanol or sodium citrate. Weak growth was detected on the medium with mannitol. During growth on glucose, strain *Nor1<sup>T</sup>* produced acetate, H<sub>2</sub> and CO<sub>2</sub>. On Medium 1 (without reduction with sodium sulfide), strain *Nor1<sup>T</sup>* was able to grow on organic substrates (glucose, sucrose and lactose) only in the presence of electron acceptors (ferric iron or thiosulfate), producing ferrous iron or H<sub>2</sub>S, respectively. Thiosulfate was found to stimulate the growth rate and cell yield of strain *Nor1<sup>T</sup>* on glucose. During growth on glucose in the presence of thiosulfate, H<sub>2</sub> was not produced; the products were acetate, CO<sub>2</sub> and H<sub>2</sub>S. No growth or H<sub>2</sub>S production was observed on the Medium 2 with peptone, lactate or acetate, in the presence of thiosulfate. The isolate showed poor growth and weak ferrous iron formation in Medium 1 with peptone or yeast extract and amorphous ferric iron oxide. The isolate did not reduce selenite or selenate during growth on peptone, sucrose or lactose.

Nitrate, sulfite and sulfate were not reduced during growth of strain *Nor1<sup>T</sup>* with CO or on organic substrates.

No growth was detected under an atmosphere of H<sub>2</sub> + CO<sub>2</sub> or H<sub>2</sub> + air.

Our tests of other hydrogenogenic carboxydrotrophs, *Carboxydotermus hydrogenoformans*, *Carboxydocella thermautotrophica* and *Caldanaerobacter subterraneus* subsp. *pacificus*, for the capacity to reduce Fe(III) during growth on CO showed that none of these three organisms was able to reduce ferric citrate or amorphous oxide of ferric iron.

### Sensitivity to antibiotics

Penicillin (100 µg ml<sup>-1</sup>), ampicillin (100 µg ml<sup>-1</sup>), streptomycin (100 µg ml<sup>-1</sup>), kanamycin (50 µg ml<sup>-1</sup>) and neomycin (50 µg ml<sup>-1</sup>) completely inhibited CO utilization and growth of strain *Nor1<sup>T</sup>*.


### DNA base composition

The DNA G+C content of strain *Nor1<sup>T</sup>* was 51.7 ± 1 mol%.

### 16S rRNA gene sequence analysis

Sequencing of the 16S rRNA gene from strain *Nor1<sup>T</sup>* placed this isolate in the domain *Bacteria*; this is in agreement with the profile of antibiotic inhibition of growth. A BLAST search (Altschul *et al.*, 1997) showed *Dendrosorobacter querciculus* (formerly *Clostridium quercicolum*) (Strömpl *et al.*, 2000) as the closest relative (89.6 % similarity). Other phyletic relatives of strain *Nor1<sup>T</sup>*, such as *Acetonema* (88.1 %), *Sporomusa* (87.4 %) and *Selenomonas* (86.2 %) species, showed lower similarity values but phylogenetic analysis presented them clustered with strain *Nor1<sup>T</sup>* (Fig. 4).

The ability to grow anaerobically on CO with the production of H<sub>2</sub> as the only reduced product was first observed in the mesophilic purple, non-sulfur bacteria *Rhodococcus gelatinosus* and *Rhodospirillum rubrum* (Uffen, 1976; Bonam *et al.*, 1989). *Carboxydotermus hydrogenoformans* was the first anaerobic, thermophilic, non-photosynthetic bacterium found that performed this process (Svetlichny *et al.*, 1991). Later, several other organisms performing the reaction of anaerobic CO oxidation with CO<sub>2</sub> and H<sub>2</sub> production were described (Svetlichny *et al.*, 1994; Sokolova *et al.*, 2001, 2004). Strain *Nor1<sup>T</sup>* is similar to *Caldanaerobacter subterraneus* subsp. *pacificus* (formerly *Carboxydi-brachium pacificum*) (Sokolova *et al.*, 2001; Fardeau *et al.*, 2004) in its ability to ferment some carbohydrates. However, they differ in the ability to reduce ferric iron. Strain *Nor1<sup>T</sup>* differs from *Caldanaerobacter subterraneus* subsp. *pacificus* JM<sup>T</sup> also in its cell morphology: *Nor1<sup>T</sup>* cells are short, curved, motile rods, while *Caldanaerobacter subterraneus* subsp. *pacificus* cells are non-motile, straight, long, thin rods, sometimes branching. The isolate described in this work differs from other previously described anaerobic CO-oxidizing hydrogenogens by its capacity for fermentative growth on several carbohydrates and for thiosulfate reduction. Unlike other anaerobic CO-oxidizing hydrogenogenic bacteria, strain *Nor1<sup>T</sup>* is able to reduce ferric iron during the growth on CO in the presence of ferric citrate or ferric iron amorphous oxide. All previously described CO-dependent, H<sub>2</sub>-generating bacteria show a cell wall structure typical of Gram-positive bacteria; they belong to the *Bacillus–Clostridium* phylum of Gram-positive bacteria and do not form a single phylogenetic cluster. Morphologically, strain *Nor1<sup>T</sup>* resembles *Thermaaerovibrio* species (Baena *et al.*, 1999; Zavarzina *et al.*, 2000), which are thermophilic, anaerobic organotrophs with vibrioid cells. As for strain *Nor1<sup>T</sup>*, *Thermaaerovibrio*



**Fig. 4.** Unrooted phylogenetic tree showing the position of strain *Nor1<sup>T</sup>*. The scale bar represents the expected number of changes per sequence position. Reliability values of internal branches are expressed as percentages. Hydrogenogenic carboxydrophic are shown in bold. GenBank/EMBL/DDBJ accession numbers for the 16S rRNA gene sequences are in parentheses.

species have curved cells, which are motile by means of a tuft of lateral flagella located on the concave side of the cell, Gram-negative, non-spore-forming and capable of growing chemo-organotrophically with fermentable substrates or lithoheterotrophically with molecular hydrogen and elemental sulfur, reducing sulfur to H<sub>2</sub>S (Zavarzina *et al.*, 2000). Strain *Nor1<sup>T</sup>* differs from *Thermaerovibrio* species by the ability to grow chemolithotrophically on CO, producing H<sub>2</sub> and CO<sub>2</sub>, and the phylogenetic distance between *Nor1<sup>T</sup>* strain and *Thermaerovibrio* species. A phylogenetic analysis revealed that strain *Nor1<sup>T</sup>* adds to the list of bacterial genera that show Gram-negative-type cell walls but belong to the phylogenetic lineage of Gram-positive bacteria (Lee *et al.*, 1978). By similarity percentage, the closest relative appears to be *D. querciculus* (Strömpl *et al.*, 2000). *D. querciculus*, formerly *Clostridium quercicolum* (Stankewich *et al.*, 1971), was isolated from discoloured tissue of living oak trees, and its cells are spore-forming, peritrichously flagellated rods. *D. querciculus* is able to ferment fructose or

glycerol, producing acetate and propionate. The G+C content of its genomic DNA is 52–54 mol%. Based on consensus tree topology, *Selenomonas ruminantium* was the closest species (86·2 % similarity) to strain *Nor1<sup>T</sup>* (Fig. 4). *Selenomonas ruminantium* is an amino-acid-fermenting anaerobic bacterium, generally found in the digestive tract of mammals (Bryant, 1956). Neither *D. querciculus* nor *Selenomonas ruminantium* was shown to grow on CO.

On the basis of its phenotypic and genotypic properties, we propose strain *Nor1<sup>T</sup>* as representative of the type species of a new genus and species, *Thermosinus carboxydovorans* gen. nov., sp. nov.

#### Description of *Thermosinus* gen. nov.

*Thermosinus* (Ther.mo.sin'us. Gr. adj. *thermos* hot; L. masc. n. *sinus* bend; N.L. masc. n. *Thermosinus* thermophilic curved rod).

Cells are motile, curved, non-spore-forming rods. Cell wall of Gram-negative type. Cells divide by binary transverse fission. Obligately anaerobic. Thermophilic. Neutrophilic. Ferment carbohydrates. DNA G+C content is  $51.7 \pm 1$  mol%. The habitat is terrestrial hot spring.

The type species is *Thermosinus carboxydovorans*.

### Description of *Thermosinus carboxydovorans* sp. nov.

*Thermosinus carboxydovorans* (car.bo.xy.di.vo'rans. N.L. neut. n. *carboxydum* carbon monoxide; L. part. adj. *vorans* devouring, digesting; N.L. part. adj. *carboxydovorans* digesting carbon monoxide).

Has the characteristics of the genus. Cells are curved rods with a length of 2.6–3 µm and a width of about 0.5 µm. Motile by means of lateral flagellation. Thermophile, grows in the temperature range 40–68 °C, with an optimum at 60 °C. Neutrophile, grows in the pH range 6.5–7.6, with an optimum at 6.8–7.0. Grows on glucose, sucrose, lactose, arabinose, maltose, fructose, xylose and pyruvate, but not on cellobiose, galactose, peptone, yeast extract, lactate, acetate, formate, ethanol, methanol or sodium citrate. During glucose fermentation produces acetate, H<sub>2</sub> and CO<sub>2</sub>. Grows chemolithotrophically on CO. Utilizes CO as the sole energy source with equimolar formation of H<sub>2</sub> and CO<sub>2</sub> according to the equation CO + H<sub>2</sub>O → CO<sub>2</sub> + H<sub>2</sub>. Reduces ferric iron during growth on CO, sucrose or lactose. Elemental sulfur, thiosulfate, sulfate and nitrate do not stimulate growth and these are not reduced during growth on CO. Thiosulfate enhances growth rate and cell yield during growth on glucose, sucrose or lactose; in this case, metabolic products are acetate, H<sub>2</sub>S and CO<sub>2</sub>. Does not utilize lactate, acetate, formate or H<sub>2</sub>, neither in the absence nor in the presence of ferric iron, thiosulfate, sulfate, sulfite, elemental sulfur or nitrate. Growth is completely inhibited by penicillin, ampicillin, streptomycin, kanamycin and neomycin. The DNA G+C content is  $51.7 \pm 1$  mol%.

The type strain is Nor1<sup>T</sup> (=DSM 14886<sup>T</sup>=VKM B-2281<sup>T</sup>); isolated from a hot pool at Norris Basin, Yellowstone National Park.

### ACKNOWLEDGEMENTS

We are grateful to A. I. Slobodkin, Institute of Microbiology, Russian Academy of Sciences, for fruitful discussion and to S. N. Gavrilov, Institute of Microbiology, Russian Academy of Sciences, for help in the determination of ferrous iron. Samples from the Yellowstone National Park were taken according to permit no. 4056. This work was supported by the NATO LST.CLG. 978269 grant, the CRDF RB2-2379-MO-02 grant and the Program 'Molecular and Cell Biology' of the Russian Academy of Sciences. J. M. G. acknowledges support from the Spanish Ministry of Science and Technology through a Ramon y Cajal contract and grant REN2002-00041.

### REFERENCES

- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**, 3389–3402.
- Baena, S., Fardeau, M.-L., Woo, T. H. S., Ollivier, B., Labat, M. & Patel, B. K. C. (1999). Phylogenetic relationships of three amino-acid-utilizing anaerobes, *Selenomonas acidaminovorans*, '*Selenomonas acidaminophila*' and *Eubacterium acidaminophilum*, as inferred from partial 16S rDNA nucleotide sequences and proposal of *Thermanaerovibrio acidaminovorans* gen. nov., comb. nov. and *Anaeromusa acidaminophila* gen. nov., comb. nov. *Int J Syst Bacteriol* **49**, 969–974.
- Balashova, V. V. & Zavarzin, G. A. (1980). Anaerobic reduction of ferric iron by hydrogen bacteria. *Microbiology* (English translation of *Mikrobiologiya*) **48**, 635–639.
- Bonam, D., Lehman, L., Roberts, G. P. & Ludden, P. W. (1989). Regulation of carbon monoxide dehydrogenase and hydrogenase in *Rhodospirillum rubrum*: effects of CO and oxygen on synthesis and activity. *J Bacteriol* **171**, 3102–3107.
- Bryant, M. P. (1956). The characteristics of *Selenomonas* isolated from bovine rumen contents. *J Bacteriol* **72**, 162–167.
- Dobbek, H., Svetlichnyi, V., Gremer, L., Huber, R. & Meyer, O. (2001). Crystal structure of a carbon monoxide dehydrogenase reveals a [Ni–4Fe–5S] cluster. *Science* **293**, 1281–1285.
- Fardeau, M.-L., Bonilla Salinas, M., L'Haridon, S., Jeanthon, C., Verhé, F., Cayol, J.-L., Patel, B. K. C., Garcia, J.-L. & Ollivier, B. (2004). Isolation from oil reservoirs of novel thermophilic anaerobes phylogenetically related to *Thermoanaerobacter subterraneus*: reassignment of *T. subterraneus*, *Thermoanaerobacter yonseiensis*, *Thermoanaerobacter tengcongensis* and *Carboxydibrachium pacificum* to *Caldanaerobacter subterraneus* gen. nov., sp. nov., comb. nov. as four novel subspecies. *Int J Syst Evol Microbiol* **54**, 467–474.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17**, 368–376.
- Kevbrin, V. V. & Zavarzin, G. A. (1992). The influence of sulfur compounds on the growth of halophilic homoacetic bacterium *Acetohalobium arabaticum*. *Microbiology* (English translation of *Mikrobiologiya*) **61**, 812–817.
- Lee, S. Y., Mabee, M. S. & Jangaard, N. O. (1978). *Pectinatus*, a new genus of the family *Bacteroidaceae*. *Int J Syst Bacteriol* **28**, 582–594.
- Ludwig, W., Strunk, O., Westram, R. & 29 other authors (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* **32**, 1363–1371.
- Marmur, J. (1961). A procedure for the isolation of desoxyribonucleic acid from microorganisms. *J Mol Biol* **3**, 208–218.
- Marmur, J. & Doty, P. (1962). Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. *J Mol Biol* **5**, 109–118.
- Pusheva, M. A. & Sokolova, T. G. (1995). Distribution of CO-dehydrogenase activity in anaerobic thermophilic carboxydrophic bacterium *Carboxydothermus hydrogenoformans* grown at the expense of CO or pyruvate. *Microbiology* (English translation of *Mikrobiologiya*) **64**, 491–495.
- Sokolova, T. G., Gonzalez, J. M., Kostrikina, N. A., Chernyh, N. A., Tourova, T. P., Kato, C., Bonch-Osmolovskaya, E. A. & Robb, F. T. (2001). *Carboxydobrachium pacificum* gen. nov., sp. nov., a new anaerobic, thermophilic, CO-utilizing marine bacterium from Okinawa Trough. *Int J Syst Evol Microbiol* **51**, 141–149.
- Sokolova, T. G., Kostrikina, N. A., Chernyh, N. A., Tourova, T. P., Kolganova, T. V. & Bonch-Osmolovskaya, E. A. (2002). *Carboxydocella thermautotrophica* gen. nov., sp. nov., a novel

- anaerobic, CO-utilizing thermophile from Kamchatkan hot spring. *Int J Syst Evol Microbiol* **52**, 1961–1967.
- Sokolova, T. G., Jeanthon, C., Kostrikina, N. A., Chernyh, N. A., Lebedinsky, A. V., Stackebrandt, E. & Bonch-Osmolovskaya, E. A. (2004).** The first evidence of anaerobic CO oxidation coupled with H<sub>2</sub> production by a hyperthermophilic archaeon isolated from a deep-sea hydrothermal vent. *Extremophiles* **8**, 317–323.
- Stankewich, J. P., Cosenza, B. J. & Shigo, A. L. (1971).** *Clostridium quercicolum* sp. n., isolated from discolored tissues in living oak trees. *Antonie van Leeuwenhoek* **37**, 299–302.
- Strimmer, K. & von Haeseler, A. (1996).** Quartet puzzling: a quartet maximum-likelihood method for reconstructing tree topologies. *Mol Biol Evol* **13**, 964–969.
- Strömpl, C., Tindall, B. J., Lunsdorf, H., Wong, T. Y., Moore, E. R. & Hippe, H. (2000).** Reclassification of *Clostridium quercicolum* as *Dendrosorobacter querciculus* gen. nov., comb. nov. *Int J Syst Evol Microbiol* **50**, 101–106.
- Svetlichny, V. A., Sokolova, T. G., Gerhardt, M., Ringpfeil, M., Kostrikina, N. A. & Zavarzin, G. A. (1991).** *Carboxydothermus hydrogenoformans* gen.nov., sp.nov., a CO-utilizing thermophilic anaerobic bacterium from hydrothermal environments of Kunashir Island. *Syst Appl Microbiol* **14**, 254–260.
- Svetlichny, V. A., Sokolova, T. G., Kostrikina, N. A. & Lysenko, A. M. (1994).** A new thermophilic anaerobic carboxydrophic bacterium *Carboxydothermus restrictus* sp. nov. *Microbiology* (English translation of *Mikrobiologiya*) **3**, 294–297.
- Svetlichny, V., Peschel, C., Acker, G. & Meyer, O. (2001).** Two membrane-associated NiFeS-carbon monoxide dehydrogenases from the anaerobic carbon-monoxide-utilizing eubacterium *Carboxydothermus hydrogenoformans*. *J Bacteriol* **183**, 5134–5144.
- Trüper, H. G. & Schlegel, H. G. (1964).** Sulfur metabolism in *Thiorhodaceae*. Quantitative measurements on growing cells of *Chromatium okenii*. *Antonie van Leeuwenhoek* **30**, 225–238.
- Uffen, R. L. (1976).** Anaerobic growth of *Rhodopseudomonas* species in the dark with carbon monoxide as sole carbon and energy substrate. *Proc Natl Acad Sci U S A* **73**, 3298–3302.
- Wolin, E. A., Wolin, M. J. & Wolfe, R. S. (1963).** Formation of methane by bacterial extracts. *J Biol Chem* **238**, 2882–2886.
- Zavarzina, D. G., Zhilina, T. N., Tourova, T. P., Kuznetsov, B. B., Kostrikina, N. A. & Bonch-Osmolovskaya, E. A. (2000).** *Thermaaerovibrio velox* sp. nov., a new anaerobic, thermophilic, organotrophic bacterium that reduces elemental sulfur, and emended description of the genus *Thermaaerovibrio*. *Int J Syst Evol Microbiol* **50**, 1287–1295.