Molecular phylogenetic analysis of the coccidian cephalopod parasites
***Aggregata octopiana* and *Aggregata eberthi* (Apicomplexa: Aggregatidae)**
from the NE Atlantic coast using 18S rRNA sequences

Sheila Castellanos-Martínez^a, Marcos Pérez-Losada^b, Camino Gestal^{a, *}

^aInstituto de Investigaciones Marinas, Consejo Superior de Investigaciones Científicas,
Eduardo Cabello,6, 36208 Vigo, Spain.

^bCIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do
Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal.

*Corresponding author at: Instituto de Investigaciones Marinas, Consejo Superior de
Investigaciones Científicas, Eduardo Cabello,6, 36208 Vigo, Spain.
Tel: +34 986 231 930; FAX: +34 986 292 762.

E-mail address: cgestal@iim.csic.es (C. Gestal)

Abstract

The coccidia genus *Aggregata* is responsible for intestinal coccidiosis in wild and cultivated cephalopods. Two coccidia species, *A. octopiana*, infecting the common octopus *Octopus vulgaris*, and *A. eberthi*, infecting the cuttlefish *Sepia officinalis*, are identified in European waters. Their morphology has been extensively studied and *A. octopiana* was redescribed in octopuses from the NE Atlantic Coast (NW Spain) to clarify confusing descriptions recorded in the past. The present study sequenced the 18S rRNA gene in *A. octopiana* and *A. eberthi* from NE Atlantic coast to assess their taxonomic and phylogenetic status. Phylogenetic analyses revealed conspecific genetic differences (2.5%) in 18S rRNA sequences between *A. eberthi* from the Ria of Vigo (NW Spain) and the Adriatic Sea. Larger congeneric differences (15.9%) were observed between *A. octopiana* samples from the same two areas, which suggest the existence of two species. Based on previous morphological evidence, host specificity data, and new molecular phylogenetic analyses, we suggest that *A. octopiana* from the Ria of Vigo is the valid type species.

Keywords: *Aggregata octopiana*; *Aggregata eberthi*; Coccidia; *Octopus vulgaris*; *Sepia officinalis*; 18S rRNA

Introduction

Coccidians are obligate intracellular parasites that cause severe injuries mainly in poultry and livestock (Levine, 1985), but are also able to infect marine fishes and molluscs causing a detrimental effect on their physiological condition (Kent and Hedrick, 1985; Lom and Dyková, 1992). Cephalopods are specifically infected by coccidians of the genus *Aggregata* (Hochberg, 1990), which are heteroxenous parasites transmitted through the food-web. Sexual stages (gamogony and sporogony) occur inside the digestive tract of the definitive cephalopod host, whereas asexual stages (merogony) can be found inside the digestive tract of the intermediate crustacean host (Hochberg, 1990).

The genus *Aggregata* has a complex taxonomic history. It was first described by Lieberkuhn (1854) as a gregarine infecting *Sepia officinalis*. Schneider (1875) described a similar parasite infecting *Octopus vulgaris*, and later the genus was correctly classified as a coccidium (Schneider 1883). Then, the genus *Aggregata* was assigned by Frenzel (1885), who described merogonic stages of the parasite in *Portunus arcuatus*. Finally, the cephalopod coccidia were classified into the family Aggregatidae by Labbé (1899). The taxonomy of the *Aggregata* species has been controversial (Hochberg, 1990), and confusing descriptions have been recorded in the past. The species *A. octopiana* was first described by Schneider (1875) in *O. vulgaris* from the English Channel and Western Mediterranean Sea (Banyuls-sur-Mer, France), and redescribed in samples from the NE Atlantic Ocean (Gestal et al., 1999b). Comparative ultrastructural studies revealed that the taxon described by others as *Aggregata spinosa* in the same host and locations using light microscopy (Moroff, 1908), was synonymous to *A. octopiana* (Gestal et al., 1999b). Consequently, ten *Aggregata* species have been described to date (see Table 1), and three of them are found in European waters: i) *A. eberthi*, which is the representative type-species of the genus *Aggregata* and infects the cuttlefish *Sepia officinalis* from the Mediterranean Sea, English Channel and NE Atlantic Ocean (Dobell, 1925); ii) *A. octopiana*, which infects the common octopus *O. vulgaris* and has

1 been re-described in hosts from the NE Atlantic Ocean (Gestal et al., 1999b); and iii) *A.*
2 *sagittata*, which infects the flying squid *Todarodes sagittatus* (Gestal et al., 2000).
3
4

5 Understanding cephalopod pathogens is particularly relevant to the worldwide aquaculture
6 of octopus species, which has to satisfy the global demand of cephalopods for human
7 consumption (Iglesias et al., 2004; Domingues et al., 2007; Solorzano et al., 2009). The
8 coccidian *A. octopiana* is known to cause heavy infections in the digestive tract of *O. vulgaris*
9 (Pascual et al., 1996). Gamogonic and sporogonic stages cause the host digestive tissue to
10 rupture (Gestal et al., 2002a). Malabsorption syndrome is a secondary effect of high infection
11 rates, reducing the growth and condition of infected octopuses (Gestal et al., 2002b) and
12 negatively impacting octopus culture (Gestal et al. 2007). Moreover, food sanitary regulations
13 forbid commercialization of parasitized fishery and aquaculture products; hence, although
14 *Aggregata* spp. are not zoonotic parasites, if oocysts are present in muscle, the octopus is
15 rejected for human consumption (Peñalver et al., 2008).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Due to the increasing importance of coccidian diseases, particularly those caused by
34 *Aggregata* species, the use of highly sensitive molecular methods for parasite diagnosis
35 becomes crucial. Furthermore, molecular approaches are also useful to characterize parasites,
36 complementing morphological descriptions, and phylogenetic classification (Jirků et al.,
37 2009; Rueckert et al., 2011). The species *A. octopiana* and *A. eberthi* have been identified and
38 characterized in the NE Atlantic coast according to morphological characters and host
39 specificity (Gestal et al., 1999b; 2002c; Gestal and Pascual, 2002). In contrast, very little is
40 known about their molecular classification and phylogenetic position, which could confirm
41 their taxonomic affiliation within the genus and validate conservative and robust phenotypic
42 characters used for species diagnosis. Kopečná et al. (2006) generated the first 18S rRNA
43 sequences for *A. octopiana* and *A. eberthi* from Croatia (Adriatic Sea); however, the
44 phylogenetic position of both coccidians remained unresolved.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 In this study, we generated new 18S rRNA nucleotide sequences for *A. octopiana* and *A.*
2 *eberthi* from the NE Atlantic coast (Galicia, NW Spain) to assess their phylogenetic position,
3
4 complement existing morphologic descriptions and validate their phenotypic characters.
5
6
7
8
9

10 **Material and Methods**

11 **Sampling and microscopic identification**

12
13
14 *Aggregata octopiana* was isolated from a pool of ten infected octopuses of the species
15
16
17 *Octopus vulgaris*, while *Aggregata eberthi* was isolated from a pool of ten infected
18
19
20 cuttlefishes of the species *Sepia officinalis*. Both cephalopod species were collected by traps,
21
22
23 an artisanal gear used by local fishermen from the Ria of Vigo, Spain (24° 14.09'N, 8°
24
25 47.18'W). The oocysts are easily observed as white spots on the digestive tract. Thus in the
26
27
28 laboratory, the presence of *Aggregata* was assessed macroscopically in each of the
29
30
31 cephalopod hosts, white oocysts were extracted from fresh caecum and intestine. Coccidians
32
33
34 were identified using light microscopy and Scanning Electron Microscopy (SEM) to analyse
35
36
37 morphology and dimensions of the fresh sporocyst and by histological analysis of the caecum,
38
39
40 which is the target organ of the infection. The infected tissue was fixed in Davidson,
41
42
43 embedded in paraffin wax and sectioned using a Microm HM-340 E microtome. Sections at 4
44
45
46 µm were stained with H-E according to standard procedures (Culling et al., 1985). For
47
48
49 Scanning Electron microscopy (SEM), purified oocyst suspension was fixed 4 h in 2.5%
50
51
52 glutaraldehyde in 0.2M cacodylate buffer (ph 7.4) at 4°C and washed for 30 min in the same
53
54
55 buffer. After dehydration in ethanol series, samples were critical point dried in CO₂ using a
56
57
58 Polaron E3000 and sputter-coated in a Polaron SC500 using 60% gold-palladium. Analysis
59
60
61 was performed with a Philips XC30 SEM operated at 10-20kV.
62
63
64
65

66 **Isolation and purification of the parasite**

1 The infected digestive tract of cephalopods was dissected and homogenized in 10 ml of
2 filtered sea water (FSW) 1% Tween80 using an electric tissue grinder (IKA-Ultra Turrax T-
3 25). Tissue homogenates were filtered twice with nylon meshes of 100 µm and 41 µm,
4
5 respectively, to remove tissue fragments. The filtrate was then centrifuged at 1000 x g for 5
6
7 min in a centrifuge Beckman GS-15R. The sporocyst were purified by density gradient
8
9 centrifugation method according to Gestal et al. (1999a), counted in a Neubauer chamber to
10
11 standardize the sample at 2×10^6 sporocyst/ml and finally, sporocyst were preserved in 70%
12
13 ethanol.
14
15
16
17
18
19

20 **DNA extraction**

21
22
23 Genomic DNA was extracted from *A. octopiana* and *A. eberthi* sporocysts. Sporocysts
24
25 were resuspended in 500 µl of extraction buffer (NaCl 100mM, EDTA 25mM pH 8, SDS
26
27 0.5%) and opened by sonication on ice (5 cycles, 40W, 50 s) to release sporozoites. After
28
29 Proteinase K (Sigma) digestion (1 mg ml^{-1}) at 37°C overnight, the DNA was purified
30
31 following the phenol:chloroform:isoamil alcohol extraction method, as described by
32
33 Sambrook et al. (1989). DNA was precipitated with ethanol and sodium acetate overnight at -
34
35 20 °C. The precipitated pellet was resuspended in 50 µl of Tris-EDTA (TE) buffer.
36
37
38
39
40

41 **DNA amplification, cloning and sequencing**

42
43
44 The small subunit 18S rRNA gene of both coccidia species was amplified by PCR using
45
46 conserved primers designed for *Aggregata* spp. (Kopečná et al. 2006) and derived from
47
48 GenBank sequences: (*Aggregata* 1-F: 5'-ATGATGAAACTGCGAAGAGC-3'; *Aggregata* 2-
49
50 R: 5'-CGACGGTATCTGATCGTCTT-3'; *Aggregata* 3-F: 5'-
51
52 GGGGGTATTTGTATTTAACAAGCA-3'; *Aggregata* 4-R: 5'-
53
54 CCTACGGAAACCTTGTTACGA-3'). *Aggregata* primers 1-2 (positions 76-1008) amplify
55
56 the initial 970 bp of the 18S rRNA gene, whereas *Aggregata* primers 3-4 (positions 871-1781)
57
58
59
60
61
62
63
64
65

1 amplify the next 915 bp. PCR reactions were performed in a total volume of 25 μ l containing
2 1 μ l 10mM dNTP mix, 0.25 μ l *Taq* DNA polymerase (Roche), 2.5 μ l *Taq* 10x buffer, 1 μ l 2.5
3 mM $MgCl_2$, 1 μ l of each primer (10 μ M) and 1 μ l of template DNA at 100 ng μ l⁻¹. The
4 temperature profile for primers 1-2 included an initial denaturation at 94°C for 10 min; 35
5 cycles of 94°C for 1 min, 57°C for 1 min and 72°C for 1 min, and a final extension at 72°C
6 for 10 min. For primers 3-4, we used an annealing temperature of 55°C. PCR products were
7 separated on 1% agarose in TAE 1x buffer gels (w/v), stained with ethidium bromide
8 including a 100-bp ladder size standard (Invitrogen) and visualized using UV light. Fresh
9 PCR products were cloned using a TOPO TA Cloning Kit (Invitrogen) according to the
10 protocol supplied by the manufacturers and transformed in TOP 10 F' competent bacteria
11 *Escherichia coli* (Invitrogen). Screening of clones carrying 18S rRNA-coding region
12 fragments was performed by PCR adding the positive colony directly to the PCR mixture
13 reaction using the corresponding *Aggregata* primers. Positive clones were purified by
14 digestion with the enzymes exonuclease I and shrimp phosphatase (SAP) (Amersham
15 Pharmacia Biothech) for 1h at 37°C. The enzymes were then denatured for 15 min at 80°C.
16 The purified PCR products were bi-directionally sequenced using the proper *Aggregata* pair
17 of primers and using ABI 3130 Genetic Analyzer according to the manufacturer's directions
18 (Applied Biosystems). Sequenced fragments from multiple clones belonging to each of the
19 two *Aggregata* species were then assembled together into two consensus sequences (see
20 below). Based on the obtained cloned sequences, the specific primers RV-F: 5'-
21 GCTTATTAAATCAGTTATAGTT-3'; RV-R: 5'ATATTTACACACATTCTAATTC-
22 3'(positions 20-1619) were designed and used to amplify almost complete 18S rRNA
23 sequences for each species (annealing temperature of 54°C). Primers *Aggregata* 5-F: 5'-
24 AAGCTCGTAGTTGCAGTTTTGA-3'; *Aggregata* 6-R: 5'
25 AACTAAGAACGGCCATGCAC-3' (positions 544-1178) equivalent to 662 bp were
26 designed to amplify the internal sequence at an annealing temperature of 54°C. All the sites in
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

these new two sequences were also present in the two entire consensus sequences assembled from multiple *Aggregata* clones.

Phylogenetic analysis

In addition to the new 18S rRNA sequences generated in this study for *A. octopiana* and *A. eberthi* from Ria of Vigo, sequences of 33 Apicomplexa taxa available in GenBank were used in the phylogenetic reconstruction. The GenBank accession numbers of the 18S rRNA gene sequences used are as follows: *Theileria buffeli* (AF236097), *Theileria* sp. (U97055), *Babesia* sp. (AY048113), *Babesia conradae* (AF158702), *Eimeria alabamensis* (AF291427), *Eimeria bovis* (U77084), *Eimeria falciformis* (AF080614), *Eimeria arnyi* (AY613853), *Cyclospora cayetanensis* (AF111183), *Cyclospora papionis* (AF111187), *Cyclospora colobi* (AF111186), *Isospora belli* (U94787), *Isospora felis* (L76471), *Goussia janae* (AY043206), *Goussia carpelli* (GU479640), *Goussia metchnikovi* (FJ009244), *Sarcocystis gracilis* (FJ196261), *Sarcocystis neurona* (U07812), *Toxoplasma gondii* (L37415), *Neospora caninum* (GQ899206), *Neospora* sp. (BPA1 U17345), *Hepatozoon canis* (EF622096), *Hepatozoon catesbiana* (AF130361), *Calyptospora spinosa* (FJ904637), *Calyptospora funduli* (FJ904645), *Adelina grylli* (DQ096836), *Adelina bambarooniae* (AF494059), *Adelina dimidiata* (DQ096835), *Tridacna hemolymph apicomplexan* (AB000912), *Klossia helicina* (HQ224955) clon 43, *Klossia helicina* (HQ224956) clone 26, *Aggregata octopiana* from the Adriatic Sea (DQ096837), *Aggregata eberthi* from the Adriatic Sea (DQ096838).

Representative species of *Babesia* and *Theileria* were used as outgroups.

All sequences were aligned in MAFFT v6 (Katoh et al., 2005; Katoh, 2008) under the Q-INS-i algorithm, which takes into account RNA secondary structure. Ambiguous regions in the resulting alignment were identified and removed using GBLOCKS 0.91b (Castresana, 2000). *Aggregata* phylogenetic relationships were inferred using maximum likelihood (ML) and Bayesian inference coupled with Markov chain Monte Carlo (BMCMC). ML trees were built

1 in RAxML v7.2.0 (Stamatakis et al., 2008) using 1,000 searches and 10 runs. JModelTest
2 v1.0.1 (Posada, 2009) was used to select the appropriate model of evolution under the Akaike
3 Information Criterion (Posada and Buckley, 2004). The general time reversible (GTR) model
4 (Tavaré, 1986), with invariable sites ($I = 0.13$) and gamma distribution ($G = 0.63$) to account
5 for the among site rate heterogeneity was chosen. Clade support was assessed using the non-
6 parametric bootstrap procedure (Felsenstein, 1985) with 5,000 bootstrap replicates run in the
7 portal CIPRES Science Gateway portal (Miller et al., 2010). BMCMC trees were built in
8 MrBayes v3.1.2 (Ronquist and Huelsenbeck, 2003). Three independent BMCMC analyses
9 were run in CIPRES with each consisting of four chains. Each Markov chain was started from
10 a random tree and run for 5×10^6 cycles, sampling every 1,000th generation. Model parameters
11 were unlinked and treated as unknown variables with uniform default priors and they were
12 estimated as part of the analysis. Convergence and mixing were monitored using Tracer v1.5
13 (Rambaut and Drummond, 2009). All sample points prior to reaching stationary were
14 discarded as burn-in. The posterior probabilities for individual clades obtained from separate
15 analyses were compared for congruence and then combined and summarized on a 50%
16 majority-rule consensus tree.

42 Results

45 Phenotypic identification of both *A. octopiana* and *A. eberthi* was performed by light
46 microscopy, histology and SEM (Fig. 1). The morphology of the analyzed oocysts, sporocyst
47 and sporozoites was consistent with those previously described as the type species from the
48 NE Atlantic (Dobell, 1925; Gestal et al., 1999b) (see Table 1). A total of 13 and nine 18S
49 rRNA partial sequence clones of *A. octopiana* and *A. eberthi*, respectively, were sequenced
50 and assembled to obtain two overlapping 50% majority-rule consensus partial 18S DNA
51 sequences of 1624 bp for *A. octopiana* and 1686 bp for *A. eberthi*. Variation among *A.*

1 *octopiana* clones was <0.55%, while variation among *A. eberthi* clones was <0.25%. In
2 addition, single sequences of similar length obtained in one single PCR amplification were
3
4 obtained for each species in order to confirm the assembled fragments. We used the consensus
5
6 sequences in all phylogenetic analyses to take into account intra-species genetic variation. The
7
8 consensus sequences of *A. octopiana* and *A.eberthi* were deposited in GenBank under the
9
10 accession numbers KC188342 and KC188343 respectively.
11
12
13
14

15 ML and BMCMC phylogenetic searches generated identical topologies, hence only the
16
17 ML tree with corrected branch lengths is presented (Fig. 2). In our analysis, two main
18
19 coccidian clades were recognized, and one of them, the adeleorinid clade, included *A.*
20
21 *octopiana* and *A. eberthi* (Fig. 2). The aggregatids from the Ria of Vigo and the Adriatic Sea
22
23 formed a highly supported monophyletic group [(bootstrap proportion (bp) = 100%, posterior
24
25 probability (pP) = 1.0)].
26
27
28
29

30 In our 18S rRNA ML tree, the minimum genetic divergence (corrected branch lengths)
31
32 observed between different recognized coccidian species pairs ranged from 0.1 to 15.1%, with
33
34 most cases above 3%. A genetic divergence of 15.9% was observed between *A. octopiana*
35
36 from the Ria of Vigo and *A. octopiana* from the Adriatic Sea, whereas a genetic divergence of
37
38 only 2.4% was found between *A. eberthi* from the Ria of Vigo and *A. eberthi* from the
39
40 Adriatic Sea (Fig. 2).
41
42
43
44
45
46
47
48

49 Discussion

50
51
52 According to their histological and ultrastructural features, *Aggregata octopiana* and *A.*
53
54 *eberthi*, have been successfully characterized in samples from the NE Atlantic Ocean by
55
56 Gestal et al. (1999b; 2002c) and Gestal and Pascual (2002). Now, the molecular
57
58 characterization carried out in this work showed a high identity with the two *Aggregata*
59
60
61
62
63
64
65

1 sequences available in GeneBank. Our phylogenetic analyses were consistent with what is
2 known from previous coccidian studies, where Eimeriidae and Sarcocystidae families formed
3 well-supported monophyletic groups (bp = 100%; pP = 1.00). Both ML and Bayesian
4 phylogenies strongly support (bp = 99%, pP = 0.99) a clade formed by Aggregatidae and
5 Adeleidae species. Within this clade, *Aggregata* species are evolutionary close to the
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
adeleorinid *Hepatozoon*, *Klossia* and *Adelina*, the latter being the most basal group, as also
suggested by the ML tree in Kopečná et al. (2006). This makes adeleorinids the most
primitive group of the Eucoccidiorida, as stated by Levin (1985), sharing with aggregatids the
formation of the sporocyst and the excystation through a longitudinal suture (Gestal et al.
1999b; Kopečná et al., 2006). However, our 18S rRNA tree, as in Kopečná et al. (2006),
cannot accurately discriminate the basal relationships and position of the genus *Aggregata*. As
has been previously suggested for other Apicomplexa (Barta et al., 2012), additional taxa and
new genetic markers will be required to resolve the relationships among these parasites.

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
Our tree shows that *A. octopiana* and *A. eberthi* from Ria of Vigo cluster with *A.*
octopiana and *A. eberthi*, respectively, from the Adriatic Sea (Fig. 2); however, the high
genetic divergence (15.9%) observed between the two *A. octopiana* samples suggests that
they represent different species (congeneric divergence). On the contrary, the genetic
divergence estimated between *A. eberthi* samples (2.5%) falls within the range observed
among populations from the same species (conspecific divergence).

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Coupled with molecular data, phenotypic characters are also required to classify
coccidians (Tenter et al., 2002). Among them, one of the most conspicuous characters is the
number of sporozoites per sporocyst (Lom and Dyková 1992). From the Adriatic Sea, scarce
and confusing records about the sporozoite number of coccidians infecting cephalopods exist.
Mladineo and Jozić (2005), for example, reported *O. vulgaris* infected by coccidian of the
genus *Aggregata* with four to five sporozoites. The Adriatic coccidia fit with the usual size
range of *A. octopiana* from the NE Atlantic (Gestal et al., 1999b), but it does not agree with

1 the number of sporozoites (eight sporozoites per sporocyst for *A. octopiana*), and spiny
2 sporocyst wall, which are the most noticeable specific features. Based on the number of
3 sporozoites, the *Aggregata* sp. from Adriatic Sea resembles *A. sagittata*, which infects only
4 the squid *Todarodes sagittatus* (Gestal et al., 2000), or *A. valdesensis*, which infects *Octopus*
5 *tehuatlensis* in SW Atlantic (Sardella et al., 2000). Interestingly, a second record by Mladineo
6 and Bočina (2007) also mentions coccidia with eight sporozoites infecting the Adriatic *O.*
7 *vulgaris*, which suggest the presence of *A. octopiana*. Thus, following morphologic
8 characters, these records suggest two different *Aggregata* species infecting *O. vulgaris* in the
9 Adriatic Sea. In addition, the absence of consistent and reliable morphological information
10 about the coccidia sequenced by Kopečná et al. (2006) makes it difficult to identify the
11 Adriatic *Aggregata* sp. correctly. Therefore, a detailed morphological characterization and
12 accurate identification of the *Aggregata* species occurring in the Adriatic Sea is needed.

13 *Octopus vulgaris* has a worldwide distribution including the Southern Indian Ocean
14 (Roper et al., 1984; Guerra et al., 2010). This octopod is now considered to form different
15 populations with differences in reproductive structures and parasite specificity (Mangold,
16 1998; Guerra pers. comm.). Because coccidia are host-specific parasites, the distinct number
17 of sporozoites in coccidia recorded from the Adriatic Sea (Mladineo and Jozić; 2005;
18 Mladineo and Bočina, 2007) suggests the possibility of different octopus populations
19 harboring different *Aggregata* parasites.

20 Therefore, based on previous morphological evidence (Gestal et al. 1999b; Gestal and
21 Pascual, 2002; Gestal et al. 2002c), host-specificity data and the new molecular phylogenetic
22 analyses herein presented, we conclude that the *Aggregata* species parasitizing the common
23 octopus *O. vulgaris* from the Ria of Vigo (NW Spain, NE Atlantic) is *A. octopiana*, the valid
24 type species. We also confirm the identification of *A. eberthi* infecting the cuttlefish *S.*
25 *officinalis* from the same locality, validating the known phenotypic characters as useful
26 diagnostic tools.

Further effort is needed to sample cephalopod hosts harboring *Aggregata* species at different geographic locations in the NE Atlantic and worldwide. Moreover, new genetic markers need to be combined with the 18S rRNA gene to improve phylogenetic analysis and complement the morphological taxonomy and classification of this poorly understood coccidian group.

Acknowledgements

This work has been funded by a research grant from the Galician Council Xunta de Galicia (10PXIB402116PR). Sheila Castellanos Martínez would like to thank CONACyT for her awarded scholarship.

References

- Barta, J.R., Ogedengbe, J.D., Martin, D.S., Smith, T.G., 2012. Phylogenetic position of the Adeleorinid coccidia (Myzozoa, Apicomplexa, Coccidia, Eucoccidiorida, Adeleorina) inferred using 18S rDNA sequences. *J. Eukaryot. Microbiol.* 59, 171-180.
- Castresana, J., 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* 17, 540-552.
- Culling, C.F.A., Allison, R.T., Barr, W.T., 1985. Cellular pathology techniques. Butterworth & Co. London.
- Dobell C.C., 1925. The life-history and chromosome cycle of *Aggregata eberthi* (Protozoa: Sporozoa: Coccidia). *Parasitol.* 17, 1-136.
- Domingues, P.M., López, N., Muñoz, J.A., Maldonado, T., Gaxiola, G., Rosas, C., 2007. Effects of a dry pelleted diet on growth and survival of the Yucatan octopus, *Octopus maya*. *Aquacult. Nutr.* 13, 273-280.
- Duszynski, D., Upton, S. J., 2001. Enteric protozoans: Cyclospora, Eimeria, Isospora, and Cryptosporidium spp. In: Samuel, W.M., Pybus, M.J., Kocan, A. (Eds.), Parasitic disease of wild mammals. 2nd edition. Wildlife Disease Association. Iowa State University Press. Ames, Iowa, pp. 416-459.
- Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783-791.
- Frenzel, J., 1885. Über einige in Seetieren lebende Gregarinen. *Arch. Milkrosk. Anat. EntwMech.* 24, 545.

- 1 Gestal, C., Abollo, E., Pascual, S., 1999a. Evaluation of a method for isolation and
2 purification of sporocysts of the cephalopod coccidian parasite *Aggregata* Frenzel, 1885
3 (Apicomplexa: Aggregatidae). *Iberus* 17, 115-121.
4
5
6
7
8 Gestal, C., Abollo, E., Pascual, S., 2002a. Observations on associated histopathology with
9 *Aggregata octopiana* infection (Protista: Apicomplexa) in *Octopus vulgaris*. *Dis. Aquat.*
10 *Org.* 50, 45-49.
11
12
13
14
15
16 Gestal, C., Guerra, A., Abollo, E., Pascual, S., 2000. *Aggregata sagittata* n. sp.
17 (Apicomplexa: Aggregatidae), a coccidian parasite from the European flying squid
18 *Todarodes sagittatus* (Mollusca: Cephalopoda). *Syst. Parasitol.* 47, 203–206.
19
20
21
22
23
24 Gestal, C., Guerra, A., Pascual, S., 2007. *Aggregata octopiana* (Protista: Apicomplexa): a
25 dangerous pathogen during comercial *Octopus vulgaris* ongrowing. *ICES J. Mar. Sci.* 64,
26 1743–1748.
27
28
29
30
31
32 Gestal, C., Nigmatullin, Ch. M., Hochberg, F.G., Guerra, A., Pascual, S., 2005. *Aggregata*
33 *andresi* n. sp. (Apicomplexa: Aggregatidae) from the ommastrephid squid *Martialia*
34 *hyadesi* in the SW Atlantic Ocean and some general remarks on *Aggregata* spp. in
35 cephalopod hosts. *Syst. Parasitol.* 60, 65-73.
36
37
38
39
40
41
42
43 Gestal, C., Paez, M., Pascual, S., 2002b. Malabsorption syndrome observed in the common
44 octopus *Octopus vulgaris* (Cephalopoda, Octopodidae) infected by *Aggregata octopiana*
45 (Protista: Apicomplexa). *Dis. Aquat. Org.* 51, 61-65.
46
47
48
49
50
51 Gestal, C., Pascual, S., 2002. Comparative x-ray microanalysis of the sporocyst wall of
52 *Aggregata octopiana* and *Aggregata eberthi* (Protista: Apicomplexa). *Eur. J. Protistol.*
53 38, 209–211.
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Gestal, C., Pascual, S., Corral, L., Azevedo, C., 1999b. Ultrastructural aspects of the
2 sporogony of *Aggregata octopiana* (Apicomplexa, Aggregatidae), a coccidian parasite of
3 *Octopus vulgaris* (Mollusca, Cephalopoda) from NE Atlantic Coast. Eur. J. Protistol. 35,
4 417-425.
5
6
7
8
9
- 10 Gestal, C., Serra, C., Guerra, A., Pascual, S., 2002c. Scratching the sporocysts surface:
11 characterization of European *Aggregata* species by Atomic Force Microscopy. Parasitol.
12 Res. 88, 242-246.
13
14
15
16
17
- 18 Gestal, C., Pascual, S., Hochberg, F.G., 2010. *Aggregata bathytherma* sp.nov. (Apicomplexa:
19 Aggregatidae), a new coccidian parasite associated with a deep-sea hydrothermal vent
20 octopus. Dis. Aquat. Org. 91, 237-242.
21
22
23
24
25
- 26 Guerra, A., Roura, A., González, A. F., Pascual, S., Cherel, Y., Pérez-Losada, M., 2010.
27 Morphological and genetic evidence that *Octopus vulgaris* Cuvier, 1797 inhabits
28 Amsterdam and Saint Paul Islands (southern Indian Ocean). ICES J. Mar. Sci. 67, 1401-
29 1407.
30
31
32
33
34
35
36
- 37 Gestal, C., Nigmatullin, Ch.M., Hochberg, F.G., Guerra, A., Pascual, S., 2005. *Aggregata*
38 *andresi* n. sp. (Apicomplexa: Aggregatidae) from the ommastrephid squid *Martialia*
39 *hyadesi* in the SW Atlantic Ocean and some general remarks on *Aggregata* spp. in
40 cephalopod hosts. Syst. Parasitol. 60, 65-73.
41
42
43
44
45
46
47
- 48 Hnida, J. A., Duszynskiy, D., 1999. Cross-Transmission Studies with *Eimeria arizonensis*, *E.*
49 *arizonensis*-like Oocysts and *Eimeria langebarteli* : Host Specificity at the Genus and
50 Species Level within the Muridae. J. Parasitol. 85, 873-877.
51
52
53
54
55
- 56 Hochberg, F.G., 1990. Diseases of Mollusca: Cephalopoda. Diseases caused by protistans and
57 metazoans. In: Kinne O. (Ed.), Diseases of marine animals, Vol. III. Cephalopoda to
58 Urochordata. Biologische Anstalt Helgoland, Hamburg, 47-227.
59
60
61
62
63
64
65

- Iglesias, J., Otero, J.J., Moxica, C., Fuentes, L., Sánchez, F. J., 2004. The completed life cycle of the octopus (*Octopus vulgaris*, Cuvier) under culture conditions: paralarval rearing using *Artemia* and zoeae, and first data on juvenile growth up to 8 months of age. *Aquacult. Int.* 12, 481-487.
- Jirků, M., Modrý, D., Šlapeta, J.R., Koudela, B., Lukeš, J., 2002. The phylogeny of *Goussia* and *Choleoeimeria* (Apicomplexa; Eimeriorina) and the evolution of excystation structures in coccidia. *Protist* 153, 380-389.
- Jirků, M., Jirků, M., Oborník, M., Lukeš, J., Modrý, D., 2009. A Model for taxonomic work on homoxenous coccidia: redescription, host Specificity, and molecular phylogeny of *Eimeria ranae* Dobell, 1909, with a review of anuran-host *Eimeria* (Apicomplexa: Eimeriorina). *J. Eukaryot. Microbiol.* 56, 39–51.
- Katoh, T., 2008. Recent developments in the MAFFT multiple sequence alignment program. *Brief. Bioinform.* 9, 286-298.
- Katoh, K., Kuma, K., Toh, H., Miyata, T., 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res.* 33, 511-518.
- Kent, M.L., Hedrick, R.P., 1985. The biology and associated pathology of *Goussia carpelli* (Leger and Stankovitch) in goldfish *Carassius auratus* (Linnaeus). *Fish Pathol.* 20, 485-494.
- Kopečná, J., Jirku, M., Oborník, M., Tokarev, Y. S., Lukeš, J., Modrý, D., 2006. Phylogenetic analysis of coccidian parasites from invertebrates: search for missing links. *Protist* 157, 173-183.
- Kuvarđina, O.N., Leander, B.S., Aleshin, V.V., Mil'nikov, A.P., Keeling, P.J., Simdyanov, T.G., 2002. *J. Eukaryot. Microbiol.* 49, 498-504.

- 1 Labbé, A., 1895. Sur le noyau et la division nucléaire chez les *Benedenia*. C. R. Acad. Sci.
2 Paris, CXX. pp. 381.
3
4
- 5 Labbé, A., 1899. Sporozoa. In: Bütschli, O. (Ed.), Das Tierreich, Lief. 5. Protozoa. R.
6 Friedlander und Sohn, Berlin, 1-180.
7
8
- 9
10
11 Levine, N.D., 1985. Phylum II. Apicomplexa Levine, 1970. In: Lee, J.J., Hutner, S.H., Bovee,
12 E. C. (Eds.), An illustrated guide to the protozoa. Society of Protozoologists, Lawrence,
13 pp. 332-374.
14
15
16
17
18
- 19 Lieberkuhn, N., 1854. Évolution des Grégarines. Mém. Couronnés et Mém. des savants
20 étrangers publ. Par L'Acad. Roy. de Belgique, XXVI. I.
21
22
23
24
- 25 Lom, J., Dyková, I., 1992. Protozoan parasites of fishes. Dev. Aquacult. Fish. Sci. 26, 1-315.
26
27
- 28 Mangold, K., 1998. The Octopodinae from the Eastern Atlantic Ocean and the Mediterranean
29 Sea. In: Voss, N.A., Vecchione, M., Toll, R.B., Sweeney, M.J. (Eds.), Systematics and
30 biogeography of cephalopods, Vol. II, Smithsonian. Contrib. Zool. 586, Washington, D.C.,
31 pp. 521-528.
32
33
34
35
36
37
38
- 39 Mathew, J.S., Van Den Bussche, R. A., Ewing, S.A., Malayer, J. R., Latha, B. R., Panciera, R.
40 J., 2000. Phylogenetic relationships of *Hepatoozon* (Apicomplexa: Adeleorina) based on
41 molecular, morphologic and life cycle characters. J. Parasitol. 86, 366-372.
42
43
44
45
46
- 47 Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for
48 inference of large phylogenetic trees. Paper presented at: Gateway Computing
49 Environments Workshop (GCE). New Orleans, LA.
50
51
52
53
54
- 55 Mladineo, I., Bočina, I., 2007. Extraintestinal gamogony of *Aggregata octopiana* in the reared
56 common octopus (*Octopus vulgaris*) (Cephalopoda: Octopodidae). J. Invertebr. Pathol.
57 96, 261–264.
58
59
60
61
62
63
64
65

- 1 Mladineo, I., Jozić, M., 2005. *Aggregata* infection in the common octopus, *Octopus vulgaris*
2 (Linnaeus, 1758), Cephalopoda: Octopodidae, reared in a flow-through system. Acta
3 Adriat. 46, 193-199.
4
5
6
7
8 Moroff, T., 1908. Die bei den cephalopoden vorkommenden Aggregataarten als grundlage
9 einer kritischen studie uber die physiologie des zellkernes. Arch. Protist 11, 1-224.
10
11
12
13 Narasimhamurti, C. C., 1979. The eimeriid *Aggregata kudo* n. sp. from *Sepia eliptica*.
14 Angew. Parasitol 20, 154-158.
15
16
17
18
19 Pascual, S., Gestal, C., Estévez, J.M., Rodriguez, H., Soto, M., Abollo, E., Arias, C., 1996.
20
21 Parasites in commercially-exploited cephalopods (Mollusca, Cephalopoda) in Spain: an
22 updated perspective. Aquaculture 142, 1–10.
23
24
25
26
27 Peñalver, J., Dolores, E. M., Muñoz, P., Cerezo, J., García, B., Viuda, E., 2008. Valoración
28 sobre la presencia y el control sanitario del coccidio *Aggregata octopiana* en el pulpo
29 común procedente de acuicultura. An. Vet. (MURCIA) 24, 57-62.
30
31
32
33
34
35
36 Posada, D., Buckley, T.R., 2004. Model selection and model averaging in phylogenetics:
37 Advantages of Akaike information criterion and Bayesian approaches over likelihood
38 ratio Tests. Syst. Biol. 53, 793-808.
39
40
41
42
43
44 Posada, D., 2009. Selection of models of DNA evolution with JModelTest. Methods Mol.
45 Biol. 537, 93-112.
46
47
48
49
50 Poynton, S., Reimschuesse, R., Stoskopf, M.K., 1992. *Aggregata dobelli* n.sp. and *Aggregata*
51 *millerorum* n.sp. (Apicomplexa: Aggregatidae) from two species of octopus (Mollusca:
52 octopodidae) from the Eastern North Pacific Ocean. J. Protozool. 39, 248-256.
53
54
55
56
57
58 Rambaut, A., Drummond, A. J., 2009. Tracer: MCMC trace analysis tool (Edinburgh,
59 Institute of Evolutionary Biology), pp. <http://tree.bio.ed.ac.uk/software/tracer/>.
60
61
62
63
64
65

- 1 Ronquist, F., Huelsenbeck, J. P., 2003. MrBayes 3: Bayesian phylogenetic inference under
2 mixed models. *Bioinformatics* 19, 1572-1574.
3
4
- 5 Roper, C. F. E., Sweeney, M. J., Nauen, C. E., 1984. FAO species catalogue. Cephalopods of
6 the world. An annotated and illustrated catalogue of species of interest to fisheries. FAO
7 Fish. Synop. 125, pp. 277.
8
9
- 10 Rueckert, S., Villette, P. M.A.H., Leander, B. S., 2011. Species boundaries in gregarine
11 apicomplexan parasites: a case study-comparison of morphometric and molecular
12 variability in *Lecudina* cf. *tuzetae* (Eugregarinorida, Lecudinidae). *J. Eukaryot.*
13 *Microbiol.* 58, 275-283.
14
15
- 16 Sambrook, J., Fritsch, E.F., Maniatis, T., 1989. Molecular cloning: a laboratory manual. 2^a ed.
17 Cold Spring Harbor Laboratory Press. New York, pp. 142-143.
18
19
- 20 Sardella, N.H., Ré, M.E., Timi, J. T., 2000. Two new *Aggregata* species (Apicomplexa:
21 *Aggregatidae*) infecting *Octopus tehuelchus* and *Enteroctopus megalocyathus* (Mollusca:
22 *Octopodidae*) in Patagonia, Argentina. *J. Parasitol.* 86, 1107-1113.
23
24
- 25 Schneider, A., 1875. Note sur la sporospermies oviformes du poulpe. *Archs. Zool. Exp. Gén.*
26 (Notes et Rev.) 4, 11-14.
27
28
- 29 Schneider, A., 1883. Nouvelles observations sur la sporulation du *Klossia octopiana*. *Archs.*
30 *Zool. Exp. Gen. (ser. 2)* 1, 77-104.
31
32
- 33 Solorzano, Y., Viana, M. T., López, L. M., Correa, J. G., True, C. C., Rosas, C., 2009.
34 Response of newly hatched *Octopus bimaculoides* fed enriched *Artemia salina*: Growth
35 performance, ontogeny of the digestive enzyme and tissue amino acid content.
36 *Aquaculture* 289 (1-2), 84-90.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Stamatakis, A., Hoover, P., Rougemont, J., 2008. A rapid bootstrap algorithm for the RAxML
2 Web servers. *Syst. Biol.* 57, 758-771.
3
4

5 Tavaré, S., 1986. Some probabilistic and statistical problems in the analysis of DNA
6 sequences, Vol. 17. American Mathematical Society, Providence, RI.
7
8
9

10 Tenter, M., Barta J.R., Beveridge, I., Duszynski, DW., Mehlhom, H., Morrison, D.A.,
11 Andrew-Thompson, R.C., Conrad, P.A., 2002. The conceptual basis for a new
12 classification of the coccidia. *Int. J. Parasitol.* 32, 595-616.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Fig. 1. Morphology of *Aggregata octopiana* and *Aggregata eberthi*. (A) Histological section of the digestive tract of *Octopus vulgaris* showing sporocyst of *A. octopiana* containing 8 sporozoites. (B) Histological section of the digestive tract of *Sepia officinalis* showing sporocyst of *A. eberthi* containing 3 sporozoites. (C) SEM photograph of *A. octopiana* sporocyst showing the spiny wall. (D) SEM photograph of *A. eberthi* showing the smooth sporocyst wall.

Fig. 2. Maximum likelihood cladogram of Apicomplexa evolutionary relationships. Corrected branch lengths estimated under the GTR+G+I evolutionary model are shown above branches and bootstrap proportions (if $\geq 70\%$)/posterior probability (if ≥ 0.95) are shown in bold below branches.

Figure

Table 1. *Aggregata* species recorded from cephalopod hosts. Length and width measurements are given as ranges (- denotes no data available).

<i>Aggregata</i> species	Host	Locality (Ocean/Sea)	Sporocysts			Sporozoites		References
			length	width	cyst wall	n°	length	
<i>octopiana</i>	<i>Octopus vulgaris</i>	NE Atlantic, W Mediterranean	11-15	11-15	spiny	8	16-24	Gestal et al. (1999b) Schneider (1875)
<i>Aggregata</i> sp.	<i>O. vulgaris</i>	E Mediterranean (Adriatic Sea)	-	-	-	4-5 8	-	Mladineo and Jozic (2005) Mladineo and Bocina (2007)
<i>dobelli</i>	<i>Enteroctopus dofleini</i>	NE Pacific	18-31	15-27	smooth	9-22	18-23	Poynton et al. (1992)
<i>milleroorum</i>	<i>O. bimaculoides</i>	NE Pacific	12-20	11-17	smooth	8-10	18-31	Poynton et al. (1992)
<i>patagonica</i>	<i>E. megalocyatus</i>	SW Atlantic	13	12	smooth	8	18	Sardella et al. (2000)
<i>valdesensis</i>	<i>O. tehuelchus</i>	SW Atlantic	10	10	-	4-8	17	Sardella et al. (2000)
<i>bathitherma</i>	<i>Vulcanoctopus hydrothermalis</i>	NE Pacific	27-32	24-32	smooth; thick	14-17	49	Gestal et al. (2010)
<i>sagittata</i>	<i>Todarodes sagittatus</i>	NE Atlantic	17	15	smooth; thick	4-8	12	Gestal et al. (2000)
<i>andresi</i>	<i>Martialia hyadesi</i>	SW Atlantic	9.7	8.2	smooth; thick	3	16-20	Gestal et al. (2005)
<i>eberthi</i>	<i>Sepia officinalis</i>	NE Atlantic, W Mediterranean	8-9	-	smooth	3	15-17	Labbé (1895)
<i>kudoï</i>	<i>S. elliptica</i>	NW Indian	9-14	-	smooth	6-12	16-18	Narasimhamurti (1979)