The neglected navigating web of the incomprehensibly emerging and re-emerging *Sarcoptes* mite

Samer Alasaad ^{a,b,*}, Luca Rossi ^d, Jorg Heukelbach ^e, Jesús M. Pérez ^f, Omar Hamarsheh ^g, Moses Otiende ^h, Xing-Quan Zhu ^{c,*}

^a Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC), Avda. Américo Vespucio s/n, 41092 Sevilla, Spain

^b Institute of Evolutionary Biology and Environmental Studies (IEU), University of Zürich Winterthurerstrasse 190, 8057 Zürich, Switzerland

^c State Key Laboratory of Veterinary Etiological Biology, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu Province 730046, China

^d Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, Università degli Studi di Torino, Via Leonardo da Vinci 44, I-10095 Grugliasco, Italy

^e Department of Community Health, School of Medicine, Federal University of Ceará, Brazil

^f Departamento de Biología Animal, Biología Vegetal y Ecología, Universidad de Jaén, Campus Las Lagunillas, s/n, E-23071 Jaén, Spain

^g Al-Quds University, Department of Biological Sciences, P.O. Box 51000, Jerusalem, Palestine

^h Kenya Wildlife Service, Veterinary Services Department, P.O. Box 40241-00100, Nairobi, Kenya

A B S T R A C T

through which parasites move from one to another host. The appropriate assimilation of parasite navigating web is pivotal for a better understanding of pathogen flow in the ecosystem, with implications for disease control. *Sarcoptes* mite has been approached from medical, veterinary, entomological, physiological and, recently, molecular sides, to understand its epidemiological navigating web between isolates from different hosts and geographical regions. The obtained conclusions are still a matter of debate. *Sarcoptes* navigating web (*Sarcoptes*-NW) is intricate and uncertain, with unexplainable pathogenic flow. In this review we summarize by which routes, under what conditions and at what level the *Sarcoptes* mite moves among its hosts.

Keywords:

Sarcoptes scabiei

Scabies

Sarcoptes mite

Parasite navigating web (Parasite-NW)

Parasite flow

Emerging and re-emerging disease

Contents

1. Introduction.....	254
2. Emerging and re-emerging <i>Sarcoptes scabiei</i>	254
3. <i>Sarcoptes</i> transmissions.....	254
3.1. Intra-mammalian groups transmission.....	255
3.1.1. Intra-humans transmission	255
3.1.2. Intra-domestic animals transmission	255
3.1.3. Intra-wild animals transmission.....	255
3.2. Inter-mammalian groups transmission.....	256
3.2.1. Humans-wild/domestic animals transmission.....	256
3.2.2. Domestic animals-wildlife transmission.....	256
4. Predictions of emergence and re-emergence.....	256
5. Public awareness.....	257
Acknowledgements.....	257
References	257

* Corresponding authors. Address: Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC), Avda. Américo Vespucio s/n 41092 Sevilla, Spain. Tel.: +41 446354914; fax: +41 446355711 (S. Alasaad), tel./fax: +86931 8342837 (X.-Q. Zhu).

E-mail addresses: samer@ebd.csic.es (S. Alasaad), xingquanzhu1@hotmail.com (X.-Q. Zhu).

1. Introduction

The neglected parasite *Sarcoptes scabiei* affects humans and a wide range of mammalian hosts worldwide (Bornstein et al., 2001; Pence and Ueckermann, 2002; Walton et al., 2004b; Alasaad et al., 2011a), entailing significant mortality in both wild and domestic animals, with considerable economic losses (Bornstein, 1995; Pence and Ueckermann, 2002; Heukelbach and Feldmeier, 2006; Dagleish et al., 2007), and ravages in human populations (Walton et al., 2004b; Hay et al., 2012).

There are no accurate estimates of the prevalence of *Sarcoptes* mite in the many animal populations worldwide, particularly in sylvatic animals. In humans, estimations indicate that hundreds of million people are infested with scabies worldwide (WHO, 2009). The prevalence of scabies in African children can be as high as 40–80%, and in remote indigenous communities in northern Australia, up to 50% of children and 25% of adults were found to be infested (Kristensen, 1991; Carapetis et al., 1997; Terry et al., 2001). Considering increasing resistance of topical chemotherapy, there is an urgent need to develop new control strategies (Currie et al., 2004; Bradberry et al., 2005; Sanderson et al., 2007), and the increasing need of sensitive and reliable diagnostic tests for humans and many domestic and wild animals (Haas et al., 2005; Heukelbach and Feldmeier, 2006).

Sarcoptes mite has been approached from morphological (Fain, 1978), medical (Carapetis et al., 1997; Feldmeier et al., 2009), veterinary (Bornstein, 1995), entomological (Fain, 1978), physiological (Arlian, 1989; Arlian et al., 1996; Haas et al., 2005) and, recently, molecular points of view (Zahler et al., 1999; Walton et al., 1997, 1999, 2004a; Alasaad et al., 2008b, 2009a,b, 2011b, 2012c; Rasero et al., 2010; Gakuya et al., 2011) aiming to understand its epidemiology between isolates from different hosts and geographical regions, but this issue is still a matter of vivid debate.

The objective of the present review is to summarize available evidence on *Sarcoptes* movement among hosts, in order to further elucidate transmission dynamics and provide an evidence-based rationale for sustainable control.

2. Emerging and re-emerging *Sarcoptes scabiei*

Parasite navigating webs (Parasite-NW) are the complex webs through which zoonotic parasites move from one host to another within the ecosystem (Polley, 2005). Pivotal to an appreciation of the function of parasite webs is an understanding of parasite flow: by which routes, under what conditions and at what level the parasite flows among its various hosts, of the same or different species (including vectors where applicable), and between the hosts and the environment (Daszak et al., 2000).

Based on the definition of emergence and re-emergence of diseases by Lederberg et al. (1992), *Sarcoptes* mite emergence could be the result of the spread of *Sarcoptes* mites in one human/animal population from an infectious origin, or simply the realization that an infection has been present in a population but undetected for several reasons. The re-emergence of *Sarcoptes* is defined as the reappearance of *Sarcoptes* after a decline in incidence. Accordingly:

(i) A *Sarcoptes* mite mange outbreak can be 'genuine' emergence of an infestation, which is new to that particular and naive animal/human population. In this case, other infested hosts (humans, domestic and/or wild animals) share space with the non-infested population are suspected to be the reservoir and source of the mites through cross-infestation. In ruminants, the possibility that mites, adapted to a "main" reservoir host, may infest other sympatric species has been documented in mange foci in Europe (León-Vizcaíno et al., 1999; Rossi et al., 2007; Oleaga et al., 2008), and field evidence has been supported by the results of

experimental infestations (Meneguz and Rossi, 1995; Lavín et al., 2000). Migration of human populations may also be a driving force for genuine emergence of scabies.

(ii) A *Sarcoptes* mite outbreak can be 'apparent' emergence/re-emergence, where *Sarcoptes* infestation is pre-existing, and the newly recognition is a result of increased detection opportunities (Kutz et al., 2003; Oleaga et al., 2008) and/or the modification of a pre-existing host-parasite equilibrium (Pence et al., 1983; Lloyd, 1995). In this second case, destabilizing modifications may have occurred on the host side, e.g. a diminished herd resistance, and/or on the parasite side, e.g. the selection of a more pathogenic strain or drug resistance (Pence and Windberg, 1994; Leung and Grenfell, 2003).

Further epidemiological studies should consider a third category, which is a mixture of genuine and apparent emergence of *Sarcoptes* mite outbreaks. Such mixture category can be revealed using both epidemiological and population genetic approaches.

3. *Sarcoptes* transmissions

It is thought that *Sarcoptes* mite originated from a human ancestor and then spread to domestic and then free-living animals (Fain, 1968; Walton et al., 2004b). *Sarcoptes* mites lack free-living stages, and individual hosts, depending on their susceptibility and behaviour, are essentially ephemeral habitats providing patchy environments that hamper random mating (Price, 1980; Criscione et al., 2005). All mites on an individual host may in fact form an 'intra-population' (Bush et al., 1997; Alasaad et al., 2008b) that has a number of recurrent generations. The number of generations is influenced by the short generation interval of this parasite (about three weeks), as well as by the infected host's life expectancy and susceptibility (Bornstein et al., 2001).

Transmission of *S. scabiei* may occur by direct or indirect contact. Larvae and nymphs of *S. scabiei* frequently leave their burrows and wander on the skin (Arlian and Vyszenski-Moher, 1988), which may, in the case of crusted scabies, harbour hundreds to several thousands of mites/cm² (Zeh, 1974; Arlian et al., 1988; Pérez et al., 2011). Some may become dislodged from the host and fall off (Arlian and Vyszenski-Moher, 1988). In controlled environments, mites may survive up until a few weeks if conditions (microclimate) are optimal (Arlian, 1989). Cooler temperatures and high humidity prolong off-host survival of *S. scabiei*, presumably because under these conditions mite metabolism is reduced (Davis and Moon, 1987; Arlian et al., 1989).

However, *Sarcoptes* navigating web (*Sarcoptes*-NW) is intricate and uncertain with fragility of parasite flow, and consequently transmission dynamics are not clear. For example, the mode of scabies transmission in humans is still discussed—whereas some authors define scabies as a sexually-transmitted disease, others suggest close body contact between children and their mothers to be the main route of transmission in endemic communities (Jackson et al., 2007). Clearly, these aspects depend on many interacting factors and differ from setting to setting.

The time needed for *S. scabiei* var. *hominis*, immediately transferred from one host to another, to initiate penetration into the stratum corneum was ~10 min (Arlian et al., 1984), and it took the mites ~35 min to become completely submerged into the epidermis. The time required for complete penetration into the stratum corneum increased as a function of the time the mites had been off their host. Experiments indicate that mites remain infective at least one-half to two-thirds of their survival time when dislodged from their host (Arlian, 1989). In historical but classical studies, Mellanby (1944) believed that the stage responsible for transmission was the young, newly fertilised adult female, which wandered around on the skin surface before burrowing. His studies

on transmission of scabies determined that patients with a high parasite load in excess of 100 adult female mites are more likely to spread the disease than those harbouring a lower number of parasites.

Additional studies on *Sarcoptes* transmission and off-host survival under different host species, climatic conditions and fomite materials are still needed (Arlian, 1989).

From the other side, the co-infection between *Sarcoptes* mite and other ecto- and endo-parasites still not well-understood and need further studies (Alasaad et al., 2008; Ryser-Degiorgis et al., 2002). The better understanding of mixed infection, involving *S. scabiei*, is of pivotal interest for the better understanding of *Sarcoptes* epidemiology, diagnosis and treatment.

3.1. Intra-mammalian group transmission

3.1.1. Intra-human transmission

Epidemics of human scabies have been hypothesised to occur on a worldwide basis in 15–25 year cycles (Falk, 1982; Arlian, 1989; Lassa et al., 2011). Nonetheless, this possible “fluctuations” of disease occurrence is not corroborated by appropriate data (Orkin and Maibach, 1978).

Estimations indicate that 300 million people, especially children, are affected (WHO, 2009). Prevalence of scabies in African children can be as high as 40–80% (Kristensen 1991; Terry et al., 2001). In low and middle income countries, scabies is a significant public health problem because it is highly prevalent and complications are frequent (Heukelbach et al., 2005; Hay et al., 2012; Heukelbach et al., 2012). Presence and severity of scabies are associated with young age, presence of many children in the household, illiteracy, low family income, poor housing, sharing clothes and towels, and irregular use of shower (Feldmeier et al., 2009). The high association with sharing clothes and towels points to a substantial contribution of fomites in the transmission of mites between humans, at least in certain settings (Zeh, 1974). However, in practice (and in endemic communities) it is not possible to quantify the importance of each single factor in epidemiological studies. Crusted scabies (also called Norwegian scabies), which is a scabies mite hyper-infection with hundreds of thousands of parasites on a single host, is rarely diagnosed in healthy individuals, but sometimes is seen in elderly, immune-suppressed patients with chronic disease, or patients suffering from malnutrition. These high transmitter individuals, who because of behaviours or mite load or some other factor, may provide a much higher risk of transmitting infestations (de Almeida Barbosa et al., 1996).

In an urban slum in Bangladesh, the incidence of scabies in children younger than 5 years was 952/1000 per year, which means that almost all children experienced at least one infestation per year (Stanton et al., 1987). This is in contrast to the situation in industrialized countries, where the disease occurs sporadically in all age groups, particularly in sexually active adults, or causes epidemics in institutions and nursing homes (Andersen et al., 2000; Heukelbach et al., 2005; Ariza et al., 2012).

3.1.2. Intra-domestic animal transmission

In Africa, mites from sheep were successfully transferred to goats (Abu-Samra et al., 1984), and transmission of mites from goats to sheep was also achieved (Ibrahim and Abu-Samra, 1987). In both cases, infested goats and sheep developed severe lesions, particularly in moistened areas, and observation of skin scrapings revealed large numbers of mites belonging to all developmental stages, which demonstrated active reproduction of transferred mites in the new host. Similar results were obtained with experimental cross-infection between camels and goats (Nayel and Abu-Samra, 1986). Conversely, other attempts to transfer *Sarcoptes* mites between domestic species failed, like those

Table 1

Experimental cross-transmission of *Sarcoptes scabiei*, host species and outcomes.

Host of origin	Receiver host	Outcome	Citation
Man	Macacus rhesus	Not successful	Ruch (1959)
Man	Domestic sheep	Not successful	Pirilä et al. (1967)
Man	Cattle	Not successful	Pirilä et al. (1967)
Domestic goat	Northern chamois	Successful	Fiebiger (1917) and Kerschagl (1955)
Domestic goat	Camel	Successful	Nayel and Abu-Samra (1986)
Domestic goat	Domestic sheep	Successful	Ibrahim and Abu-Samra (1987)
Domestic goat	Southern chamois	Successful	Lavín et al. (2000)
Domestic goat	Donkey	Successful	Abu-Samra et al. (1984)
Northern chamois	Domestic goat	Successful	Meneguz and Rossi (1995)
Southern chamois	Cattle	Not successful	Lavín et al. (2000)
Southern chamois	Domestic sheep	Not successful	Lavín et al. (2000)
One humped camel	Domestic goat	Successful	Nayel and Abu-Samra (1986)
Dog	Domestic sheep	Not successful	Arlian et al. (1988)
Dog	Domestic goat	Not successful	Arlian et al. (1988)
Dog	Cattle	Not successful	Arlian et al. (1988)
Dog	Cat	Not successful	Arlian et al. (1988)
Dog	Rabbit	Successful	Arlian et al. (1984)
Dog	Guinea pig	Not successful	Arlian et al. (1984)
Dog	House mouse	Not successful	Arlian et al. (1984)
Dog	Pig	Not successful	Arlian et al. (1984)
Dog	Rat	Not successful	Arlian et al. (1984)
Dog	Man	Not successful	Kutzer and Grunberg (1970)
Coyote	Dog	Not successful	Samuel (1981)
Coyote x dog hybrid	Red fox	Successful	Stone et al. (1972)
Wolf	Dog	Not successful	Samuel (1981)
Red fox	Dog	Not successful	Samuel (1981)
Red fox	Dog	Successful	Stone et al. (1972) and Bornstein (1991)
Red fox	Coyote x dog hybrid	Successful	Stone et al. (1972)
Red fox	Coyote	Successful	Samuel (1981)
Red fox	Man	Not successful	Kutzer and Grunberg (1970)
Tapir	Man	Not successful	Kutzer and Grunberg (1970)

involving dogs, sheep, goats, cattle and cats (Arlian et al., 1988) (Table 1).

3.1.3. Intra-wild animal transmission

Mange, caused by *S. scabiei*, has been reported from many species of wild mammals worldwide. Some of the most remarkable hosts include (1) canids in North America (Pence et al., 1983; Little et al., 1998); (2) red foxes and other canids in Europe (Mörner, 1992; Ippen et al., 1995; Gortazar et al., 1998); (3) red foxes and dingoes in Australia (McCarthy, 1960); (4) chamois, ibex and a variety of other ungulates in Europe (Ippen et al., 1995; Fernández Morán et al., 1997; Rossi et al., 2007); (5) felids in Europe (Mörner,

1992) and Africa (Young, 1975); (6) wild boar in Europe (Ippen et al., 1995); (7) wombats in Australia (Skerratt et al., 1998); and (8) a range of ungulates, primates, and canids in Africa (Young, 1975; Kalema et al., 1998; Munang'andu et al., 2010; Gakuya et al., 2012b). New outbreaks in wildlife are being continually reported worldwide (Alasaad et al., 2012a).

The molecular analyses of sympatric wild animals in Europe show unambiguously that there has been a lack of gene flow or recent admixture between carnivore-, herbivore-, and omnivore-derived *Sarcoptes* populations (Rasero et al., 2010). Host-taxon-derived effect seems stronger driver of intra-specific differentiation than geographical separation, and it seems temporally stable (Alasaad et al., 2011b). The existence of host-taxon-derived *Sarcoptes* mites could explain why mange-free populations can live in sympatry with mangy animals, as is the case of the mange-free *Capra ibex* and *Rupicapra rupicapra* of the western Italian Alps that live in close proximity with the endemically mangy population of red fox *Vulpes vulpes*, and wild boar *Sus scrofa* (Rasero et al., 2010). *Vice versa*, this effect could be the immediate reason why cross-transmission easily occurs, in Europe, between foxes, dogs and other canids, as well as felids including domestic European cats, when these hosts are exposed to the same vulpine-derived strain (Bornstein 1995) (Table 1).

The studied wild animals in Europe (from which the host-taxon was coined) lacked of clear predator-prey interaction and putative inter-specific transmission models. Recent epidemiological and molecular studies revealed potential prey-to-predator gene flow between wild animals from Africa. Cheetahs were infected from both wildebeest and Thomson's gazelle, while lions were only infected from wildebeest. In a predator/prey ecosystem, like Masai Mara in Kenya, it seems that *Sarcoptes* infestation is from prey-to-predator, in relation to the predator's "favourite prey", which could be attributed to the direct skin-to-skin prolonged contact between the predator and the prey during hunting process (Gakuya et al., 2011, 2012a).

3.2. Inter-mammalian group transmission

3.2.1. Humans-wild/domestic animal transmission

Humans are occasionally infected with *S. scabiei* of animal origin (Arlian, 1989). This includes human infection from camel, cat, chamois, coyote, dog, ferret, fox, wombat, gazelle, goat, horse, llama, pig, sheep, water buffalo and cattle (Kutzer, 1970; Fain, 1978; Chakrabarti et al., 1981; Samuel, 1981; Folz, 1984; Chakrabarti, 1990; Mitra et al., 1993; Morsy et al., 1995; Skerratt and Beveridge, 1999; Menzano et al., 2004; Bazargani et al., 2007). Those infected include farmers, personnel working in slaughterhouses researchers, veterinarians, wildlife biologist, and pet owners.

Most human infections from animal sources involve only topographically circumscribed body regions, are short-lived and self-limiting, lasting from a few days to several weeks (Arlian, 1989). As infestation is less severe and clinical features are different from infestation with *Sarcoptes scabiei* var. *hominis*, this condition is usually referred in the medical literature as "pseudoscabies". Lesions are frequently seen on the trunk, arms, and abdomen and rarely on the finger webs and genitalia. In addition, the incubation period is markedly shorter, and mite burrows are not regularly seen (Orkin and Maibach, 1991). Pruritus may be as intense as in classical scabies, but symptoms usually wane within a maximum of 2–4 weeks. The human host is not a source of infection to other humans. Reportedly, non-human mites most frequently originate from dogs infested with *S. scabiei* var. *canis* (30–50% of cases) (Thomsett, 1968; Folz, 1984; Aydingöz and Mansur, 2011).

The short-life and self-limiting infection of *Sarcoptes* mites of animal origin when transmitted to humans was confirmed by

molecular analysis. Walton et al., 1999, 2004a used multi-locus genotyping applied to microsatellite markers to substantiate previous findings that gene flow between human- and dog-derived mite populations is extremely rare in scabies-endemic Aboriginal communities in Australia.

3.2.2. Domestic animals-wildlife transmission

Considering the increasing deforestation and changing ecosystems worldwide, there has been more and more contact of domestic animals with sylvatic animals. In fact, the introduction of infected domestic animals and the success of the *Sarcoptes* mite in adapting to new highly susceptible wild hosts have been proposed as the origin of *Sarcoptes* mite epizootics in previously mange-free wildlife populations (Arlian, 1989). This scenario was also proposed for a *Sarcoptes* mite outbreak in Iberian ibex in Sierra Nevada mountain range, Spain in the 1970s, possibly from domestic infected goats (Pérez et al., 1997).

Molecular studies have been carried out to understand *Sarcoptes* transmission between humans and animals, and between wild host species, but never between domestic and wild animals, and hence effort should be made in this direction. We know very little of interspecific transmission between (putative) domestic reservoirs and wildlife, while we know much more on wildlife to wildlife transmission.

In the Sierras de Cazorla, Segura y Las Villas Natural Park (southern Spain) mange has become an endemic disease. At the end of the 1980s, it firstly affected Iberian ibex (*Capra pyrenaica*) causing devastating mortality rates with a reduction of the population by 95%. Then, the disease reached other sympatric wild ungulate species: European mouflon (*Ovis aries*), red deer (*Cervus elaphus*) and fallow deer (*Dama dama*). The origin of this epizooty is attributed to infested domestic goat herds (see, for instance, León-Vizcaíno et al., 1999). Regarding this situation, an attempt to experimentally infect chamois (*Rupicapra rupicapra*) with *Sarcoptes* mites derived from naturally-infested domestic goats was successful (Lavín et al., 2000). The opposite route of transmission was also previously demonstrated under experimental conditions (Meneguz and Rossi, 1995). A recent outbreak of *Sarcoptes* mite in domestic goats from northern Italy has been attributed to presumable contact with wild animals under natural conditions (Menzano et al., 2007).

Domestic-wild animal transmission usually produce high mortality and morbidity rates, probably because of insufficient management actions.

4. Predictions of emergence and re-emergence

Sarcoptes emergence and re-emergence could be predicted by different approaches, such as (i) endemic cycles, (ii) increasing agglomeration of people, wild and domestic animals, and increasing contact between human-wildlife-livestock. Notwithstanding, the prediction of *Sarcoptes* emergence and re-emergence could be affected by (a) the emerging of resistant against some acaricides (Currie et al., 2004), (b) the lack of effective diagnosis method for many host species especially the wild ones (Alasaad et al., 2012b), and (c) the totally neglected role of fomites in *Sarcoptes* transmission.

Better prediction of scabies and mange emergence and re-emergence could be achieved through better understanding of interspecific transmission patterns, which requires further studies including the effect of (i) travelling, sexual and familiar contact, biomedical manipulation and migration in intra-human transmission; (ii) agricultural intensification and animal translocation in intra-domestic animals transmission; (iii) translocation and animal stress in intra-wild animals transmission; (iv) technology and

Fig. 1. The neglected *Sarcoptes* navigating web (*Sarcoptes*-NW), including multiple putative routes of parasite flow between mammalian hosts, following authors' criteria. Dotted lines: flow through fomites. Solid lines: flow through direct contact (host-to-host). Circular arrow: flow within animal group. Question mark: questionable parasite flow. Curved lines: flow between groups.

industry manipulation in humans-domestic animal transmission; (v) human encroachment, tourism, and ecological manipulation in humans-wild animal transmission; and (vi) animal encroachment and animal introduction in domestic animals-wild life transmission (Fig. 1).

5. Public awareness

A better understanding of *Sarcoptes* navigating web is of pivotal interest for the public health. Unfortunately many facets of *Sarcop*-tes navigating web regarding humans are not yet clear, including human-to-human transmission, especially the sexually transmission and family care, and wild and domestic animals to humans transmission. Integral programs should be carried out including epidemiological and genetic studies of *Sarcoptes* navigating web between humans and wild/domestic animals, and review of knowledge and practices regarding *Sarcoptes* infection in humans (Gakuya et al., 2012b). Effective and operator-friendly methods for direct and indirect diagnosis of scabies are still missing and the resistance of *Sarcoptes* to drugs is now becoming of major concern.

Acknowledgements

Project support was provided by Proyecto de Excelencia RNM 06400 (Junta de Andalucía, Spain) and Juan de la Cierva Grant (Ministerio Innovación y ciencia, Spain). XQZ is supported by the International Science & Technology Cooperation Program of China

(Grant No. 2013DFA31840), the Science Fund for Creative Research Group of Gansu Province (Grant No. 1210RJIA006) and the Special Fund for Agro-scientific Research in the Public Interest (Grant No. 201303037).

References

- Abu-Samra, M.T., Ibrahim, K.E.E., Abdel Aziz, M., 1984. Experimental infection of goats with *Sarcptes scabiei* var. *ovis*. *Ann. Trop. Med. Parasitol.* 78, 55–61.
- Alasaad, S., Granados, J.E., Cano-Manuel, F.J., Meana, A., Zhu, X.Q., Pérez, J.M., 2008a. Epidemiology of fasciolosis affecting Iberian ibex (*Capra pyrenaica*) in southern Spain. *Parasitol. Res.* 102, 751–755.
- Alasaad, S., Soglia, D., Sarasa, M., Soriguer, R.C., Pérez, J.M., Granados, J.E., Rasero, R., Zhu, X.Q., Rossi, L., 2008b. Skin-scale genetic structure of *Sarcptes scabiei* populations from individual hosts: empirical evidence from Iberian ibex-derived mites. *Parasitol. Res.* 104, 101–105.
- Alasaad, S., Rossi, L., Soriguer, R.C., Rambozzi, L., Soglia, D., Pérez, J.M., Zhu, X.Q., 2009a. *Sarcptes* mite from collection to DNA extraction: the lost realm of the neglected parasite. *Parasitol. Res.* 104, 723–732.
- Alasaad, S., Soglia, D., Spalenza, V., Maione, S., Soriguer, R.C., Pérez, J.M., Rasero, R., Ryser-DeGiorgis, M.P., Nimmervoll, H., Zhu, X.Q., Rossi, L., 2009b. Is ITS-2rDNA suitable marker for genetic characterization of *Sarcptes* mites from different wild animals in different geographic areas? *Vet. Parasitol.* 159, 181–185.
- Alasaad, S., Holt, D., Walton, S., Rossi, L., Bornstein, S., Abu-Madi, M., Soriguer, R.C., Fitzgerald, S., Zhu, X.Q., Zimmermann, W., Ugbomoiko, U.S., Heukelbach, J., 2011a. *Sarcptes*-World Molecular Network (*Sarcptes*-WMN): integrating research on scabies. *Int. J. Infect. Dis.* 15, 294–297.
- Alasaad, S., Oleaga, A., Casais, R., Rossi, L., Molinar Min, A., Soriguer, R., Gortazar, C., 2011b. Temporal stability in the genetic structure of *Sarcptes scabiei* under the host-taxon law: empirical evidences from wildlife-derived *Sarcptes* mite in Asturias, Spain. *Parasit. Vectors* 4, e151.
- Alasaad, S., Ndeereh, D., Rossi, L., Bornstein, S., Permunian, R., Soriguer, R.C., Gakuya, F., 2012a. The opportunistic *Sarcptes scabiei*: a new episode from giraffe in the drought-suffering Kenya. *Vet. Parasitol.* 185, 359–363.

- Alasaad, S., Permian, R., Gakuya, F., Mutinda, M., Soriguer, R.C., Rossi, L., 2012b. Sarcoptic-mange detector dogs used to identify infected animals during outbreaks in wildlife. *BMC Vet. Res.* 8, e110.
- Alasaad, S., Schuster, R.K., Gakuya, F., Theneyan, M., Jowers, M.J., Maione, S., Molinar Min, A., Soriguer, R.C., Rossi, L., 2012c. Applicability of molecular markers to determine parasitic infection origins in the animal trade: a case study from Sarcopites mites in wildebeest. *Forensic Sci. Med. Pathol.* 8, 280–284.
- Andersen, B.M., Haugen, H., Rasch, M., Haugen, A.H., Tageson, A., 2000. Outbreak of scabies in Norwegian nursing homes and home care patients: control and prevention. *J. Hosp. Infect.* 45, 160–164.
- Ariza, L., Walter, B., Worth, C., Brockmann, S., Weber, M.L., Feldmeier, H., 2012. Investigation of a scabies outbreak in a kindergarten in Constance, Germany. *Eur. J. Clin. Microbiol. Infect. Dis.* [Epub ahead of print].
- Arlian, L.G., Runyan, R.A., Estes, S.A., 1984. Cross infectivity of *Sarcoptes scabiei*. *J. Am. Acad. Dermatol.* 10, 979–986.
- Arlian, L.G., Vyszenski-Moher, D.L., 1988. Life cycle of *Sarcoptes scabiei* var. *canis*. *J. Parasitol.* 74, 427–430.
- Arlian, L.G., Ahmed, M., Vyszenski-Moher, D.L., 1988. Effects of *Sarcoptes scabiei* var. *canis* (Acari: Sarcopitidae) in blood indices of parasitized rabbits. *J. Med. Entomol.* 25, 360–369.
- Arlian, L.G., 1989. Biology, host relations, and epidemiology of *Sarcoptes scabiei*. *Ann. Rev. Entomol.* 34, 139–161.
- Arlian, L.G., Vyszenski-Moher, D.L., Pole, M.J., 1989. Survival of adults and developmental stages of *Sarcoptes scabiei* var. *canis* when off the host. *Exp. Appl. Acarol.* 6, 181–187.
- Arlian, L.G., Morgan, M.S., Arends, L.L., 1996. Immunologic cross-reactivity among various strains of *Sarcoptes scabiei*. *J. Parasitol.* 82, 66–72.
- Aydingöz, I.E., Mansur, A.T., 2011. Canine scabies in humans: a case report and review of the literature. *Dermatology* 223, 104–106.
- Bazargani, T.T., Hallan, J.A., Nabian, S., Rahbari, S., 2007. Sarcopit mangle of gazelle (*Gazella subgutturosa*) and its medical importance in Iran. *Parasitol. Res.* 101, 1517–1520.
- Bradberry, S.M., Cage, S.A., Proudfoot, A.T., Vale, J.A., 2005. Poisoning due to pyrethroids. *Toxicol. Rev.* 24, 3–106.
- Bornstein, S., 1991. Experimental infection of dogs with *Sarcoptes scabiei* derived from naturally infected wild red foxes (*Vulpes vulpes*): clinical observations. *Vet. Dermatol.* 2, 151–159.
- Bornstein, S., Mörner, T., Samuel, W.M., 2001. *Sarcoptes scabiei* and sarcopit mangle. In: Samuel, W.M., Pybus, M.J., Kocan, A.A. (Eds.), *Parasitic Diseases of Wild Mammals*, second ed. Iowa State University Press, Ames, pp. 107–119.
- Bornstein, S., 1995. *Sarcoptes scabiei* infections of the domestic dog, red fox and pig. PhD thesis. Department of Veterinary Microbiology, Section of Parasitology, Swedish University of Agricultural Sciences and National Veterinary Institute, Uppsala, Sweden.
- Bush, A.O., Lafferty, K.D., Lotz, J.M., Shostak, A.W., 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J. Parasitol.* 83, 575–583.
- Carapetis, J.R., Connors, C., Yarmirr, D., Krause, V., Currie, B.J., 1997. Success of a scabies control program in an Australian aboriginal community. *Pediatr. Infect. Dis. J.* 16, 494–499.
- Chakrabarti, A., 1990. Pighandler's stitch. *Int. J. Dermatol.* 29, 205–206.
- Chakrabarti, A.N., Chatterjee, A.A., Chakrabarti, K., Sengupta, D.N., 1981. Human scabies from contact with water buffaloes infested with *Sarcoptes scabiei* var. *bubalis*. *Ann. Trop. Med. Parasitol.* 75, 353–357.
- Criscione, C.D., Poulin, R., Blouin, S., 2005. Molecular ecology of parasites: elucidating ecological and microevolutionary processes. *Mol. Ecol.* 14, 2247–2257.
- Currie, B.J., Harumal, P., McKinnon, M., Walton, S.F., 2004. First documentation of in vivo and in vitro ivermectin resistance in *Sarcoptes scabiei*. *Clin. Infect. Dis.* 39, 8–12.
- Dagleish, M.P., Ali, Q., Powell, R.K., Butz, D., Woodford, M.H., 2007. Fatal *Sarcoptes scabiei* infection of blue sheep (*Pseudois nayaur*) in Pakistan. *J. Wildl. Dis.* 43, 512–517.
- Daszak, P., Cunningham, A.A., Hyatt, A.D., 2000. Emerging infectious diseases of wildlife – threats to biodiversity and human health. *Sci.* 287, 443–449.
- Davis, D.P., Moon, R.D., 1987. Survival of *Sarcoptes scabiei* (De Geer) stored in three media at three temperatures. *J. Parasitol.* 73, 661–662.
- de Almeida Barbosa, A., Correia Silva, T.M., Ribeiro Santos, M.I., Gomez Garrido, M.F., Niljay Patel, B., Cavalcanti, M., Oliveira Riccio, M.C., 1996. Coexistence of an unusual form of scabies and lepromatous leprosy. A case report. *Pathol. Res. Pract.* 192, 88–90.
- Fain, A., 1968. Étude de la variabilité de *Sarcoptes scabiei* avec une révision des Sarcopitidae. *Acta Zool. Pathol. Antverp.* 47, 1–196.
- Fain, A., 1978. Epidemiological problems of scabies. *Int. J. Dermatol.* 17, 20–30.
- Falk, E.S., 1982. Scabies: some aspects of its relationship to the immune mechanisms. PhD Thesis. Tromsø University, Tromsø.
- Feldmeier, H., Jackson, A., Ariza, L., Lins-Calheiros, C.M., Soares, V.L., Oliveira, F.A., 2009. The epidemiology of scabies in an impoverished community in rural Brazil: presence and severity of disease are associated with poor living conditions and illiteracy. *J. Am. Acad. Dermatol.* 60, 436–443.
- Fernández Morán, J., Gómez, S., Ballesteros, F., Quirós, P., Benito, J.L., Feliu, C., Nieto, J.M., 1997. Epizootiology of sarcopit mangle in a population of Cantabrian chamois (*Rupicapra pyrenaica parva*) in northwestern Spain. *Vet. Parasitol.* 73, 163–171.
- Fiebigler, J., 1917. Neue Untersuchungen über Gamsenräude. Übertragung der Ziegenräude auf Gamsen. *Wien. tierärztl. Mschr.* 4, 433–450.
- Folz, S.D., 1984. Canine scabies (*Sarcoptes scabiei*) infestation. *Compend. Contin. Educ. Vet.* 6, 176–180.
- Gakuya, F., Rossi, L., Ombui, J., Maingi, N., Muchemi, G., Ogara, W., Soriguer, R.C., Alasaad, S., 2011. The curse of the prey: *Sarcoptes mite* molecular analysis reveals potential prey-to-predator parasitic infestation in wild animals from Masai Mara, Kenya. *Parasit. Vectors* 4, e193.
- Gakuya, F., Ombui, J., Maingi, N., Muchemi, G., Ogara, W., Soriguer, R.C., Alasaad, S., 2012a. Sarcopit mangle and cheetah conservation in Masai Mara (Kenya): epidemiological study in a wildlife/livestock system. *Parasitology* 139, 1587–1595.
- Gakuya, F., Ombui, J., Heukelbach, J., Maingi, N., Muchemi, G., Ogara, W., Mijele, D., Alasaad, S., 2012b. Knowledge of mangle among Masai pastoralists in Kenya. *PLoS ONE* 7, e43342.
- Gortazar, C., Villafuerte, R., Blanco, J.C., Fernandez-de-Luco, D., 1998. Enzootic sarcopit mangle in red foxes in Spain. *Z. Jagdwiss.* 44, 251–256.
- Haas, N., Wagemann, B., Hermes, B., Henz, B.M., Heile, C., Schein, E., 2005. Crossreacting IgG antibodies against fox mite antigens in human scabies. *Arch. Dermatol. Res.* 296, 327–331.
- Hay, R.J., Steer, A.C., Engelman, D., Walton, S., 2012. Scabies in the developing world – its prevalence, complications, and management. *Clin. Microbiol. Infect.* 18, 313–323.
- Heukelbach, J., Wilcke, T., Winter, B., Feldmeier, H., 2005. Epidemiology and morbidity of scabies and pediculosis capitis in resource-poor communities in Brazil. *Br. J. Dermatol.* 153, 150–156.
- Heukelbach, J., Feldmeier, H., 2006. Scabies. *Lancet* 367, 1767–1774.
- Heukelbach, J., Mazigo, H.D., Ugbomoiko, S., 2012. Impact of scabies in resource-poor communities. *Clin. Microbiol. Infect.* 18, 313–323.
- Ibrahim, K.E.E., Abu-Samra, M.T., 1987. Experimental transmission of a goat strain of *Sarcoptes scabiei* to desert sheep and its treatment with ivermectin. *Vet. Parasitol.* 26, 157–164.
- Ippen, R., Nickel, S., Schröder, H.D., 1995. Krankheiten des Jagdbaren Wildes. Deutscher Land-wirtschaftsverlag, Berlin, pp. 189–195.
- Jackson, A., Heukelbach, J., Feldmeier, H., 2007. Transmission of scabies in a rural community. *Braz. J. Infect. Dis.* 11, 386–387.
- Kalema, G., Koch, R.A., Macfie, E., 1998. An outbreak of sarcopit mangle in free-ranging mountain gorillas (*Gorilla gorilla berengei*) in Bwindi Impenetrable National Park, South Western Uganda. In: Proceedings AAZV and AAWV Joint Conference, Omaha, Nebraska, USA, p. 438.
- Kerschagl, W., 1955. Übertragbarkeit der Gamsräude auf andere Wildarten. *Anblick* 10, 295–296.
- Kristensen, J.K., 1991. Scabies and pyoderma in Lilongwe, Malawi. Prevalence and seasonal fluctuation. *Int. J. Dermatol.* 30, 699–702.
- Kutz, S.J., Hoberg, E.P., Nagy, J., Polley, L., Elkin, B., 2003. Emerging parasitic infections in arctic ungulates. *Integrat. Comp. Biol.* 44, 109–118.
- Kutzer, E., 1970. *Sarcoptes-Milben und Sarcoptes-räude der Haustiere*. Merkblätter über angewandte Parasitenkunde und Schädlingsbekämpfung. *Angew. Parasitol.* 11, 1–22.
- Kutzer, E., Grunberg, W., 1970. Transmission of sarcopit mangle from animals to man. *Vet. Bull.* 40, 4.
- Lavín, S., Ruiz-Bascaran, M., Marco, I., Fondevila, M.D., Ramis, A.J., 2000. Experimental infection of chamois (*Rupicapra pyrenaica parva*) with *Sarcoptes scabiei* derived from naturally infected goats. *J. Vet. Med.* 47, 693–769.
- Lassa, S., Campbell, M.J., Bennett, C.E., 2011. Epidemiology of scabies prevalence in the U.K. from general practice records. *Br. J. Dermatol.* 164, 1329–1334.
- Lederberg, J., Shope, R.E., Oaks, S.C. (Eds.), 1992. *Emerging Infections: Microbial Threats to Health in the United States*. National Academies Press, Washington, DC.
- León-Vizcaíno, L., Ruíz de Ybáñez, M.R., Cubero, M.J., 1999. Sarcopit mangle in Spanish ibex from Spain. *J. Wildl. Dis.* 35, 647–659.
- Leung, B., Grenfell, B.T., 2003. A spatial stochastic model simulating a scabies epidemic and coyote population dynamics. *Ecol. Model.* 166, 41–52.
- Little, S.E., Davidson, W.R., Howerth, E.W., Rakich, P.M., Nettles, V.F., 1998. Diseases diagnosed in red foxes from Southeastern United States. *J. Wildl. Dis.* 34, 620–624.
- Lloyd, S., 1995. Environmental influences on host immunity. In: Grenfell, B.T., Dobson, A.P. (Eds.), *Ecology of Infectious Diseases in Natural Populations*. Cambridge University Press, Cambridge, UK, pp. 327–361.
- McCarthy, P.H., 1960. The presence of sarcopit mangle in the wild fox (*Vulpes vulpes*) in Central Queensland. *Aust. Vet. J.* 36, 359–360.
- Mellanby, K., 1944. The development of symptoms, parasitic infection and immunity in human scabies. *Parasitology* 35, 197–206.
- Meneguz, P.G., Rossi, L., 1995. Experimental transmission of *Sarcoptes scabiei* from chamois to domestic goat. *Proc. Soc. It. Sci. Vet.* 49, 755–756.
- Menzano, A., Rambozi, L., Rossi, L., 2004. Outbreak of scabies in human beings, acquired from chamois (*Rupicapra rupicapra*). *Vet. Rec.* 155, 568.
- Menzano, A., Rambozi, L., Rossi, L., 2007. A severe episode of wildlife derived scabies in domestic goats in Italy. *Small Rumin. Res.* 70, 154–158.
- Mitra, M., Mahanta, S.K., Sen, S., Ghosh, C., Hati, A.K., 1993. *Sarcoptes scabiei* in animals spreading to man. *Trop. Geogr. Med.* 45, 142–143.
- Morsy, T.A., Bakr, M.E., Ahmed, M.M., Kotb, M.M., 1995. Human scabies acquired from a pet puppy. *J. Egypt. Soc. Parasitol.* 24, 305–308.
- Mörner, T., 1992. Sarcopit mangle in Swedish wildlife. *Rev. Sci. Tech.* 11, 115–121.
- Munang'andu, H.M., Siamudaala, V.M., Matandiko, W., Munyeme, M., Chembensofu, M., Mwase, E., 2010. *Sarcoptes mite* epidemiology and treatment in African

- buffalo(*Syncerus caffer*) calves captured for translocation from the Kafue game management area to game ranches. *BMC Vet. Res.* 6, 29.
- Nayel, N.M., Abu-Samra, M.T., 1986. Experimental infection of the one-humped camel (*Camelus dromedarius*) with *Sarcoptes scabiei* var. *cameli* and *S. scabiei* var. *ovis*. *Ann. Trop. Med. Parasitol.* 80, 553–561.
- Oleaga, A., Casais, R., González-Quirós, P., Prieto, M., Gortazar, C., 2008. Sarcoptic mange in red deer from Spain: improved surveillance or disease emergence? *Vet. Parasitol.* 154, 103–113.
- Orkin, M., Maibach, H.I., 1978. Scabies in children. Symposium on Pediatric Dermatology. *Pediatr. Clin. North Am.* 25, 371–386.
- Orkin, M., Maibach, H.I., 1991. Ectoparasitic diseases. In: Orkin, M., Maibach, H., Dahl Norwalk, M.V. (Eds.), *Dermatology*, vol. 3. Appleton and Lange, VA, pp. 205–214.
- Pence, D.B., Windberg, L.A., Pence, B.C., Sprowls, R., 1983. The epizootiology and pathology of sarcoptic mange in coyotes, *Canis latrans*, from South Texas. *J. Parasitol.* 69, 1100–1115.
- Pence, D.B., Windberg, L.A., 1994. Impact of a sarcoptic mange epizootic on a coyote population. *J. Wildl. Mgmt.* 58, 624–633.
- Pence, D.B., Ueckermann, E., 2002. Sarcoptic mange in wildlife. *Rev. Sci. Tech.* 21, 385–398.
- Pérez, J.M., Ruiz-Martínez, I., Granados, J.E., Soriguer, R.C., Fandos, P., 1997. The dynamics of sarcoptic mange in the ibex population of Sierra Nevada in Spain— influence of climatic factors. *J. Wildl. Res.* 2, 86–89.
- Pérez, J.M., Granados, J.E., Sarasa, M., Serrano, E., 2011. Usefulness of estimated surface area of damaged skin as a proxy of mite load in the monitoring of sarcoptic mange in free-ranging populations of Iberian wild goat, *Capra pyrenaica*. *Vet. Parasitol.* 176, 258–264.
- Pirilä, V.V., Salo, O.P., Kiistala, R., 1967. Scabies Norvegica. *Acta Derm. Venereol.* 47, 267–268.
- Polley, L., 2005. Navigating parasite webs and parasite flow: emerging and re-emerging parasitic zoonoses of wildlife origin. *Int. J. Parasitol.* 35, 1279–1294.
- Price, P.W., 1980. *Evolutionary Biology of Parasites*. Princeton University Press, Princeton, NJ.
- Rasero, R., Rossi, L., Soglia, D., Maione, S., Sacchi, P., Rambozzi, L., Sartore, S., Soriguer, R.C., Spalenza, V., Alasaad, S., 2010. Host taxon-derived *Sarcoptes* mite in European wild animals revealed by microsatellite markers. *Biol. Conserv.* 143, 1269–1277.
- Rossi, L., Fraquelli, C., Vesco, U., Permian, R., Somavilla, G.M., Carmignola, G., Da Pozzo, M., Meneguz, P.G., 2007. Descriptive epidemiology of a scabies epidemic in chamois in the Dolomite Alps, Italy. *Eur. J. Wildl. Res.* 53, 131–141.
- Ruch, T.C., 1959. *Diseases of laboratory primates*. W.B. Saunders Company, Philadelphia.
- Ryser-Degiorgis, M.P., Ryser, A., Bacciarini, L.N., Angst, C., Gottstein, B., Janovsky, M., Breitenmoser, U., 2002. Notoedric and sarcoptic mange in free-ranging lynx from Switzerland. *J. Wildl. Dis.* 2002 (38), 228–232.
- Samuel, W.M., 1981. Attempted experimental transfer of sarcoptic mange (*Sarcoptes scabiei*, Acarina: Sarcoptidae) among red fox, coyote, wolf and dog. *J. Wildl. Dis.* 17, 343–347.
- Sanderson, H., Laird, B., Pope, L., Brain, R., Wilson, C., Johnson, D., 2007. Assessment of the environmental fate and effects of ivermectin in aquatic mesocosms. *Aquat. Toxicol.* 85, 229–240.
- Skerratt, L.F., Martin, R.W., Handasyde, K.A., 1998. Sarcoptic mange in wombats. *Aust. Vet. J.* 76, 408–410.
- Skerratt, L.F., Beveridge, I., 1999. Human scabies of wombat origin. *Aust. Vet. J.* 9, 607.
- Stanton, B., Khanam, S., Nazrul, H., Nurani, S., Khair, T., 1987. Scabies in urban Bangladesh. *J. Trop. Med. Hyg.* 90, 219–226.
- Stone, W.B., Parks, E., Weber, B.L., Parks, F.J., 1972. Experimental transfer of sarcoptic mange from red foxes and wild canids to captive wildlife and domestic animals. *N.Y. Fish Game J.* 19, 1–11.
- Terry, B.C., Kanjah, F., Sahr, F., Korteque, S., Dukulay, I., Gbakima, A.A., 2001. *Sarcoptes scabiei* infestation among children in a displacement camp in Sierra Leone. *Public Health* 115, 208–211.
- Thomsett, L.R., 1968. Mite infestation of man contracted from dogs and cats. *Br. Med. J.* 3, 93–95.
- Walton, S.F., Currie, B.J., Kemp, D.J., 1997. A DNA fingerprinting system for the ectoparasite *Sarcoptes scabiei*. *Mol. Biochem. Parasitol.* 85, 187–196.
- Walton, S.F., Choy, J.L., Bonson, A., Valle, A., McBroom, J., Taplin, D., Arlian, L., Mathews, J.D., Currie, B., Kemp, D.J., 1999. Genetically distinct dog-derived and human-derived *Sarcoptes scabiei* in scabies-endemic communities in northern Australia. *Am. J. Trop. Med. Hyg.* 61, 542–547.
- Walton, S.F., Dougall, A., Pizzutto, S., Holt, D., Taplin, D., Arlian, L.G., Morgan, M., Currie, B.J., Kemp, D.J., 2004a. Genetic epidemiology of *Sarcoptes scabiei* (Acari: Sarcoptidae) in northern Australia. *Int. J. Parasitol.* 34, 839–849.
- Walton, S.F., Holt, D.C., Currie, B.J., Kemp, D.J., 2004b. Scabies: new future for a neglected disease. *Adv. Parasitol.* 57, 309–376.
- WHO (World Health Organisation), 2009. Available from: <http://www.who.int/en/>.
- Young, E., 1975. Some important parasitic and other diseases of lion, *Panthera leo*, in the Kruger National Park. *J. S. Afr. Vet. Assoc.* 46, 181–183.
- Zahler, M., Essig, A., Gothe, R., Rinder, H., 1999. Molecular analyses suggest monospecificity of the genus *Sarcoptes* (Acari: Sarcoptidae). *Int. J. Parasitol.* 29, 759–766.
- Zeh, J.B., 1974. Infestation of sarcoptic mange on red fox in New York. *New York Fish Game J.* 21, 182–183.