Title:

Lacustrine carbonates of Iberian Karst Lakes: sources, processes
and depositional environments

Valero - Garcés, Blas (1); Morellón, Mario (2); Moreno, Ana (1); Corella, Juan
Pablo (3); Martín-Puertas, Celia (4); Barreiro, Fernando (1); Pérez, Ana (1);
Giralt, Santiago (5); Mata-Campo, María Pilar (6)

(1) Instituto Pirenaico de Ecología, CSIC, Avda Montañana 1005, Apdo 13034,
50080 Zaragoza, Spain

(2) Instituto de Geociencias, CSIC, UCM. José Antonio Nováis, 2, 3ª planta, 3b.
Facultad de Ciencias Geológicas, Univ. Complutense. 28040 Madrid, Spain.

(3) Museo Nacional de Ciencias Naturales (MNCN-CSIC), C/ Serrano 115bis,
28006 Madrid, Spain

(4) Helmholtz Centre Potsdam, GFZ German Research Centre for
Geosciences, Section 5.2 Climate Dynamics and Landscape Evolution,
Telegrafenberg D-14473, Potsdam, Germany

(5) Institute of Earth Sciences Jaume Almera, CSIC, Lluís Solé I Sabarís s/n,
08028 Barcelona, Spain

(6) Instituto Geológico y Minero de España, C/ La Calera, 1, 28760 Tres Cantos
– Madrid, Spain

Abstract

Carbonates are the main components of Iberian Quaternary lake sediments. In this review we summarize the main processes controlling carbonate deposition in extant Iberian lakes located in Mesozoic and Tertiary carbonate-dominated regions and formed through karstic activity during the Late Quaternary. The lakes, relatively small (1 ha to 118 ha) and relatively shallow ($Z_{\text{max}} = 11$ to 40 m) provide examples of the large variability of sedimentary facies, depositional environments, and carbonate sources. Hydrology is dominated by groundwater inflow except those directly connected to the fluvial drainage. Nine lakes have been selected for this review and the main facies in palustrine, littoral and profundal environments described and interpreted.

Clastic carbonates occur in all Iberian lakes due to the carbonate composition of the bedrocks, surface formations and soils of the watersheds. Low temperatures and dilute meteoric waters seem responsible for the low carbonate content of sediments in high elevation lakes in the glaciated terrains in the Pyrenees and the Cantabrian Mountains. Clastic carbonates are dominant in small karst lakes with functional inlets where sediment infill is dominated by fining upward sequences deposited during flood events. Re-working of littoral carbonates is common in shallow environments and during low lake level stages. In most lakes, endogenic carbonate production occurs in two settings: i) littoral platforms dominated by *Chara* and charophyte meadows and ii) epilimnetic zone as biologically-mediated calcite precipitates. Continuous preservation of varves since the Mid-Holocene only occurs in one of the deepest lake (Montcortès Lake, up to 30 m) where calcite laminae textures (massive, fining upward and coarsening upward) reflect seasonal changes in limnological conditions. However, varves have been formed and preserved in

most of the lakes during short periods associated with increased water depth and more frequent meromictic conditions.

Most Iberian lakes are in a mature stage and karstic processes are not very active. An outstanding example of a lake with intense karstic activity is Banyoles Lake where increased spring discharge after long rainy periods causes large remobilization and re-suspension of the sediments accumulated in the deepest areas, leading to the deposition of thick homogeneous layers (homogeinites).

The Iberian karst lake sequences underline the large variability of facies, carbonate sources, and depositional environments in small lake systems. They illustrate how lake types evolve through the existence of a lake basin at centennial or even smaller time scales. Hydrology is the paramount control on facies and depositional environment patterns distribution and lake evolution and, consequently, a lake classification is proposed based on hydrology and sediment input. A correct interpretation of carbonate sources and depositional history is key for using lake sequences as archives of past global changes.

Keywords: limnogeology, karst, lacustrine facies, Late Quaternary, endogenic and clastic carbonates, Iberian Peninsula

1. Introduction

Lakes occur almost in every geographic, geologic, and climatic context where geomorphic factors enable the creation of accommodation space (basin) and hydrologic balance is adequate to accumulate water. A variety of processes are able to create and maintain lakes on Earth: tectonics, eolian, fluvial, karstic, volcanic, impact events and even anthropogenic activities (Gierlowski-Kordesch and Kelts, 2000; Cohen, 2003). In carbonate and evaporite bedrock, karstic processes such as dissolution and collapse are very effective in creating centripetal drainage patterns and depressions for lake development. Although karst lakes are particularly abundant in some regions such as North and Central America (Florida and Yucatan), southern China (Yunnan Province), and the Mediterranean Basin (Balkans, Italian and Iberian Peninsula), they occupy less than 1% of the total global lake area (Cohen, 2003). The dynamics of these lakes, particularly the hydrology and hydrochemistry are greatly controlled by exogenic and endogenic karstic processes. This activity leads to the generation of funnel-shaped dolines with steep margins, which have high depth/water surface ratios (Palmquist, 1979; Cvijic, 1981; Gutiérrez-Elorza, 2001). These steep-sided lakes are conducive to water stratification and contribute to a dynamic depositional environment with abrupt changes in geochemical and limnological factors, such as water chemistry, temperature, light penetration or oxic/anoxic conditions at the lake bottom (Renaut and Gierlowski-Kordesch, 2010)- In addition, the direct connection to aquifers makes these systems very sensitive to regional hydrological balances, experiencing considerable lake level, water chemistry, and biological fluctuations in response to changes in effective moisture (Nicod, 2006; Morellón et al., 2009a;

93 Sondi and Jura Ci, 2009). The development of lakes on evaporite and limestone
94 substrates favors sulfate and carbonate-rich waters. Lakes within continental
95 evaporitic provenance include Lac de Besse, France (Nicod, 1999); Lago di
96 Pergusa, Italy (D'Amore, 1983); Lake Demiryurt, Turkey (Alagöz, 1967) and
97 Laguna Grande de Archidona, (Pulido-Bosch, 1989), Lake Banyoles (Julià, 1980),
98 and Lake Estanya (Morellón et al., 2009a) in Spain. Generally carbonate-rich and
99 chloride-rich lake waters develop on lakes within marine formations, e.g., Lake
100 Vrana, Croatia (Schmidt et al., 2000) and Lake Zoñar, Spain (Valero-Garcés et al.,
101 2003).

102 Lakes are active agents in the carbon cycle (Dean and Gorham, 1998;
103 Schrag et al., 2013) and, in particular, karst lakes are key elements in the recycling
104 of old carbonate and evaporite formations. Besides endogenic carbonate
105 deposition, allochthonous siliciclastic and carbonate material is delivered to the
106 lakes from the watersheds and re-distributed in the lake through waves in the
107 littoral areas, and turbidite – type (Moreno et al., 2008; Valero Garcés et al., 2008)
108 and mass-wasting processes (Morellón et al., 2009a) in the distal realm.

109 Lacustrine depositional models have been extensively described in the
110 literature (Kelts and Hsü, 1978; Dean, 1981; Dean and Fouch, 1983; Eugster and
111 Kelts, 1983; Wright, 1990; Gierlowski-Kordesch and Kelts, 1994; Talbot and Allen,
112 1996; Gierlowski-Kordesch and Kelts, 2000) and some reviews have been recently
113 published (Gierlowski-Kordesch, 2010; Renaut and Gierlowski, 2010; Last and
114 Last, 2012). Massive, fine-grained carbonates with abundant fauna and flora
115 remains and laminated carbonates are common facies in the geological lacustrine
116 record as well (Gierlowski-Kordesch and Kelts, 1994; Gierlowski-Kordesch and

Kelts, 2000). Most lake sediments contain some carbonate and the variety of lacustrine facies reflects the different settings: carbonate-rich lakes (Platt and Wright, 1991), ephemeral and shallow saline lakes (Last, 1990; Smoot and Lowenstein, 1991; Renault and Last, 1994), volcanic-related lakes (Negendank and Zolitschka, 1993; Pueyo et al., 2011), glacial and periglacial lakes (Kelts, 1978; Hsü and Kelts, 1985; Davaud and Girardclos, 2001). Karst lakes have provided numerous paleohydrologic and paleoclimate reconstructions e.g., Lago d'Accesa, Italy (Magny et al., 2006; Magny et al., 2007); Lake Banyoles, Spain (Pérez-Obiol and Julià, 1994; Valero-Garcés et al., 1998; Höbig et al., 2012); Lago di Pergusa, Italy (Sadori and Narcisi, 2001; Zanchetta et al., 2007); Lake Zoñar, Spain (Martín-Puertas et al., 2008; Martín-Puertas et al., 2009). Several sequences have been described in detail in the last decade: Petén Itza (Anselmetti et al., 2006; Hodell et al., 2008), Ohrid (Lézine et al., 2010; Lindhorst et al., 2010; Vogel et al., 2010), Estanya (Morellón et al., 2009b), however, comprehensive facies and depositional models for karstic lakes are scarce (see Morellón et al., 2009a).

In this paper we review the available sedimentary sequences from extant permanent karst lakes in Spain. In the Iberian Peninsula these lakes are relatively small (1 – 118 ha surface area) and not very deep (max depth range between 11 and 40 m) but they occur in a variety of geographic, geologic and climatic contexts. These relatively small, perennial Iberian karst lakes may serve as facies analogs for larger systems and help to identify sources and processes controlling lacustrine carbonate deposition in modern lakes and pre-Quaternary lacustrine formations. Detailed facies analyses and depositional models would also help to untangle the

endogenic and clastic contribution to carbonate budget. These depositional models provide a dynamic framework for integrating all paleolimnological data necessary to decipher the high-resolution paleoenvironmental information archived in these lake sequences (Last and Smol, 2001; Renaut and Gierlowski-Kordesch, 2010).

2. Geologic and geographic settings

The Iberian Peninsula is composed of three main geological units (Gibbons and Moreno, 2002): i) the Iberian Massif, made up of Palaeozoic and Proterozoic rocks, ii) the Alpine Ranges (Pyrenees, Betics and Iberian Mountains), composed of Mesozoic and Cenozoic sedimentary formations affected by the Alpine orogeny, and iii) large tectonic Cenozoic basins, such as the Ebro, Duero, Tagus and Guadalquivir basins. Karst lakes occur in the Alpine Ranges and the Cenozoic basins where carbonate bedrock is dominant (Fig. 1).

The climate of the Iberian Peninsula is varied, displaying strong temperature and humidity gradients due to the altitudinal variability, the presence of mountain ranges, and the interplay of Mediterranean and Atlantic processes (Sumner et al., 2001). The inland areas and the mountain ranges experience a moderate continental climate while oceanic climate dominates in the north and west and a warm Mediterranean climate along the Mediterranean coast. Average annual temperatures range from 0 °C in the northern mountains (Pyrenees, Cantabrian) to 18 °C in the southern and eastern areas; highest precipitations are recorded on

the northern mountains (> 1500 mm/yr), while the inland and southern regions are drier (< 500 mm/yr) (Capel Molina, 1981).

Permanent karst lakes occur in a variety of geological settings in Spain (Figs. 1 and 2): 1) glaciated terrains (Lakes Enol, Basa de la Mora and Marboré), spring and tufa – dammed areas (Lakes Taravilla, Basturs, Ruidera and Banyoles), fluvial drainages (Guadiana River and Daimiel National Park), carbonate formations in the Iberian Range (Lakes of Cañada del Hoyo and El Tobar), the Pre-Pyrenees (Lakes Estanya and Montcortès), and the Guadalquivir Basin (Lakes Zóñar, Archidona, Medina and Fuentepiedra), and in halokinetic salt structures (Lake Arreo). Some ephemeral saline lakes in Tertiary continental basins (e.g., Monegros in the Ebro Basin) formed by combined karstic and aeolian processes. A few karstic systems – Lake Banyoles (Sanz, 1981; Brusi et al., 1990), Lake Estanya (Pérez-Bielsa et al., 2012), Lake Zóñar (Valero-Garcés et al., 2006) - have quantitative water balances that demonstrate groundwaters are the main input to these lakes. Qualitative hydrogeological models suggest that this is common to all karstic lakes in Spain except those directly connected to fluvial drainages (Lakes Taravilla, and Ruidera).

For this review we have selected lakes with detailed facies descriptions that have been studied by our research group during the last decade following a similar approach and methodology (see methods section). The selected case studies illustrate different hydrological and limnological situations in four carbonate-rich regions: i) The Iberian Range, ii) high altitude lakes in the Pyrenees and the

Cantabrian Mountains, iii) mid-altitude lakes in the western Ebro Basin and the Pre-Pyrenees and iv) the Guadalquivir Basin (Table 1, Figs. 1 and 2).

i) The karst lakes in the Iberian Range formed during an Upper Pliocene - Pleistocene karstification phase affecting mostly Jurassic limestones (Gutierrez-Elorza and Peña Monné, 1979). The karstification led to the formation of sinkholes, examples include the seven dolines in Cañada del Hoyo (39°N, 1°52'W, 1000 m a.s.l.), the Lake Taravilla (40°39' N, 1°58' W, 1100 m asl) and Lake El Tobar (40°32'N; 3°56'W, 1200 m asl). Sediment cores are available for four of the Cañada del Hoyo lakes: La Cruz, Lagunillo del Tejo, El Tejo and La Parra (see Fig. 1 for location). **Laguna de la Cruz** (surface area = 1.4 ha; 132 m of diameter; $Z_{\max}$ = 25 m) exhibits meromixis and the occurrence of 'whittings' during the summer (Vicente and Miracle, 1987; Dasi and Miracle, 1991; Miracle et al., 1992; Rodrigo et al., 1993; Julià et al., 1998; Romero-Viana et al., 2008; Romero-Viana et al., 2011). Short cores have been recently obtained from Lake **Lagunillo del Tejo** (López-Blanco et al., 2011; Romero-Viana et al., 2009), and the study of long cores from Lakes **Tejo** and **La Parra** is in progress by our team (Barreiro-Lostres et al., 2011; Barreiro-Lostres et al., 2013).

Lake El Tobar is the largest karstic lake in the region, with a maximum depth of 20 m and a surface area of 70 ha. It is composed of two sub-basins, a deeper, meromictic with an anoxic hypersaline (NaCl) hypolimnion and a shallower main basin (Vicente and Miracle, 1987; Miracle et al., 1992; Vicente et al., 1993; Romero Viana, 2007). Exokarstic processes, particularly tufa damming are responsible for the genesis of other lakes such as **Lake Taravilla**, a shallow ($Z_{\max}$

= 12 m) lake, dammed behind a tufa barrage in a small tributary of the Tagus River. The lake is hydrologically open with a strong fluvial influence and also a large palustrine area.

ii) In the Cantabrian Mountains, the **Lake Enol** basin (43°11'N, 4°09'W, 1070 m asl) was deepened by a glacier, though previous karstic processes within these Carboniferous limestone (Valdeteja and Picos de Europa) and sandstone (Amieva series) formations were involved in the origin of the basin (Marquínez and Adrados, 2000). In the Pyrenees, we have selected two examples: Lakes Marboré (00°23' E, 42°41' N) and Basa de la Mora (42°32' N 0°19' E). Lake **Marboré** is a high altitude (2590 m asl), small (14.3 ha) and relatively deep ($Z_{\max} = 27$ m) lake located in Mesozoic – Tertiary carbonate formations. The lake basin was also carved by glaciers and, although the last remnants of the Monte Perdido Glaciers occur in the peaks surrounding the basin, there is no hydrological connection between the lake and the glaciers nowadays. Lake Marboré has an extreme oligotrophic environment similar to other high altitude lakes in the Pyrenees with low alkalinity waters (Catalán et al., 2006). Lake **La Basa de la Mora** (1914 m a.s.l) is a shallow ($Z_{\max} < 4$ m) lake developed behind a terminal moraine in the Cotiella Massif.

iii) In the mid-altitude areas of the western Ebro Basin and Pyrenean domain there are numerous examples of karst lakes, with four highlighted here:

Lake Arreo (42°46' N, 2°59' W; 655 m asl) is located on the southwestern edge of the Salinas de Añana diapir, an ellipsoidal, 5.5 x 3.2 km halokinetic structure developed in Upper Triassic evaporite formations in the NW Ebro River

Basin (Garrote Ruíz and Muñoz Jiménez, 2001). Lake Arreo is the best example in Spain of a karst lake formed in evaporites and constitutes the deepest body of water ($Z_{\max}= 24.8$ m) with gypsum substrate in the Iberian Peninsula (Martín-Rubio et al., 2005). The lake (6.57 ha) has a funnel-shaped deep basin and a shallow platform occupies 2/3 of the lake's total surface area (Rico et al., 1995). The lake is hydrologically open, with a small stream that enters the lake from the east and a small ephemeral outlet, a tributary of the Ebro River, which flows southward.

Balsas de Estanya ($42^{\circ}02'$ N, $0^{\circ}32'$ E; 670 m asl) is a karstic lake complex located in a relatively small endorheic basin covering 18.8 ha (López-Vicente et al., 2009) that belongs to a larger Miocene structure, the Saganta-Estopiñan polje (Sancho-Marcén, 1988). The karstic system consists of three dolines with water depths of 7, 12 and 20 m, and one that is seasonally flooded. The largest lake is composed of two coalescent dolines, 20 and 12 m deep with no surface inlets or outlets. **Lake Montcortès** ($42^{\circ} 19'$ N, $0^{\circ} 59'$ E, 1027 m asl) is one of the deepest karst lakes in the Iberian Peninsula ($Z_{\max}= 30$ m) (Alonso, 1998). The lake is almost circular in shape (maximum length = ~525 m, maximum width = ~450 m), has no permanent inlet, and an outlet stream located on the north shore controls maximum lake level (Camps et al., 1976). The lake is oligotrophic and meromictic, with permanently anoxic freshwater monimolimnion below 18 m in summer and 20 m in winter (Modamio et al., 1988).

Lake Banyoles ($42^{\circ}1'$ N; $2^{\circ}4'$ E, 173 m asl) is the largest karst lake complex in Spain (118 ha surface area) with a maximum depth of 40 m, although some active sinkholes may reach up to 80 m. The basin developed in Tertiary evaporitic

formations contains eleven karstic depressions (Bischoff et al., 1994; Canals et al., 1990) connected by their epilimnetic waters, but with isolated hypolimnions, which show differential anoxic periods, from 1 to 12 months/year (Prat and Rieradevall, 1995). The lake is hydrologically open, connected to a large Paleogene limestone and gypsum aquifer (Sanz, 1981; Brusi et al., 1990). About 80 % of the inflow is through groundwaters. Active springs in the sinkholes are responsible for sediment re-mobilization and development of sediment plumes (Serra et al., 2005).

iv) In the Guadalquivir Basin, **Lake Zoñar** (37°29' N, 4°41' W, 300 m a.s.l.) is the deepest (14.5 m) and largest lake (37 ha, surface area) (Valero-Garcés et al., 2006). The origin of this lake basin is related to karstic activity along some fault structures (Sánchez et al., 1992). The lake has no surface outlet and the inlets are temporary; groundwater is the main input to the lake.

3. Methods

For this review, we have selected nine lakes to summarize their sediment sequences from the twelve karst lakes listed in Table 1. Sedimentological data have been previously published except for Lakes La Parra, Banyoles, El Tejo and Marboré. All the sequences have been analyzed following similar methodologies and the detailed analyses and techniques applied to each lake are described in publications listed in Table 1. Lake watersheds were identified and mapped using topographic and geologic maps and aerial photographs. Surface sediments were sampled and long cores were retrieved using modified Kullenberg piston coring equipment and Uwitec® corers and platform. Physical properties (magnetic susceptibility and density) were measured with a Geotek Multi-Sensor Core Logger (MSCL) every 1 cm. The cores were subsequently split lengthwise in two halves

and imaged with high-resolution cameras mounted in core scanners. Color parameters (L^* , a^* , b^*) were extracted using Adobe Photoshop® and ImageJ software. Sedimentary facies were defined after visual description of core sections and microscopic observation of smear slides, following the methodology described in Schnurrenberger et al. (2003). This classification protocol is more adequate for unconsolidated soft sediments because it integrates meso-scale (visual descriptions) and micro-scale (microscopic) observations and it includes a semi-quantification of the main components (organic remains, mineral grains, biological content). For the purposes of this paper the facies described for each lake have been synthesized according to compositional and textural characteristics, mainly grain size and lamination. Compositional measurements include total organic (TOC) and inorganic (TIC) carbon and total nitrogen (TN). Bulk sediment mineralogy was characterized by X-ray diffraction and scanning electron microscope (SEM) observations completed textural and compositional characterizations. Large thin sections (100 x 15 x 35 mm) were prepared using the freeze-dry technique and subsequent impregnation with epoxy resin under vacuum conditions (Brauer and Casanova, 2001; Lotter and Lemcke, 1999).

The chronology of the selected sequences has been constrained with radiocarbon and $^{210}\text{Pb}/^{137}\text{Cs}$ techniques (see references in Table 1). They include the last 30000 years in Lake Enol, 22000 years in Lake Estanya, most of the Holocene in Lakes Banyoles, Marboré, and Basa de La Mora; the mid to late Holocene in Lake Montcortès (ca. 6000 years) and Lake Zoñar (ca. 4000 years) and the last millennia in Lakes Arreo, Tejo, La Parra and Taravilla.

4. Facies associations and depositional environments

Lacustrine environments are defined by depth and energy levels and include littoral, sublittoral and profundal (Renaut and Gierlowski-Kordesch, 2010). Littoral environments group the shallow-water zones around the lake margins affected by waves; sublittoral environments are transitional to deeper waters but still within the photic zone; finally, distal-profundal environments are below wave base with minimal to no sunlight penetration (Renaut and Gierlowski-Kordesch, 2010). Palustrine environments are transitional between lacustrine and terrestrial, and affected by significant periods of subaerial exposure and pedogenesis (Alonso-Zarza and Wright, 2010). Due to the small size of most Spanish karstic lakes, “distal” environments are quite close to the shoreline and we prefer to maintain the double name: distal-profundal. Also, we have included sublittoral and littoral in the same environment with examples of both, gently (bench, terrace or platform) and steeply sloping margins. Considering the characteristics of the Iberian karstic lakes (relatively small and shallow, significant clastic input) we have identified three main depositional environments: palustrine, littoral - sublittoral and profundal - distal). Gierlowski-Kordesch (2010) considered five main types of carbonate lacustrine facies; laminated, massive, microbial, marginal and open-water and Alonso-Zarza and Wright (2010) defined eleven palustrine carbonate facies. We have grouped the variety of facies in Iberian karstic lakes using key sedimentological (lamination and grain size) and compositional (carbonate versus siliciclastic and organic) properties (Table 2 and Figures 3 to 14). Detailed descriptions of sedimentary facies for each lake are provided elsewhere (see references in Table 1). Three

facies associations and depositional environments can be identified in most Recent Iberian karstic lake basins (Fig. 15): i) palustrine, ii) littoral-sublittoral and iii) profundal- distal.

4.1 Palustrine Facies Associations

Facies.

There are two main types of palustrine sedimentary facies (P, Table 2):

1. **Clastic Facies.** Clastic facies are well developed in palustrine areas with alluvial inlets such as Lakes Banyoles, Taravilla, and Arreo. They include gravels and sands (Palustrine Coarse, PC) and fine sands and silts (Palustrine Fine, PF). In Lake Arreo massive to banded conglomerate layers with a sandy matrix and silty sand layers with abundant plant remains occur at the base of the littoral sequences. Fine, massive carbonate facies have edaphic textures, bioturbation and evidence of subaerial exposure (root traces, cracks) (Fig. 4). In Lake Taravilla the fining-upwards sequences are composed of gravel beds up to 25 cm thick composed of Cretaceous rock fragments and tufa clasts with a carbonate and quartz sandy matrix and coarse silts with abundant plant remains (Figs. 3A and 5).

Interpretation. Coarse facies arranged in fining-upward sequences represents deposition in palustrine areas and proximal littoral zones with a strong alluvial/deltaic influence. Finer facies represent deposition in wetlands with lower alluvial influence and subsequent subaerial exposure, colonization by terrestrial vegetation and development of incipient soils. These coarse clastic layers show similarities with carbonate-filled channels facies in modern wetlands and ancient

palustrine settings (Alonso-Zarza and Wright, 2010).

2. Organic – rich (PO). Cm to dm thick, massive organic-rich silts with abundant plant remains occur at the top of fining-upward sequences in Lake Taravilla and are common in shallow palustrine settings at Lake Arreo (Fig. 4).

Interpretation. These facies represent organic accumulation in low energy palustrine settings (wetland type) with abundant vegetation. They are similar to organic-rich marlstone and clays facies defined by Alonso-Zarza and Wright (2010)

4.2. Littoral Facies Associations

Facies

1. Coarse clastic (LC). Coarse (gravel, sand and coarse silt size) facies occur in most sequences, although they are minor components. These facies are well represented only in Lake Banyoles, where shallow (ca. 10 m), flat platforms are extensively developed in the northern sub-basin. In Lake Banyoles, the littoral sediments are banded to massive carbonate-rich (up to 95 % calcite) (Pérez-Obiol and Julià, 1994) organized in sand-silt-mud fining-upward sequences (Valero-Garcés et al., 1998; Höbig et al., 2012). These facies contain bioclasts (*Chara* fragments, gastropods, ostracods and reworked carbonate coatings with plant remains (Morellón et al., in review). In Lake Arreo, massive silty sand layers occur at the southern margin, with high carbonate content (up to 80%) and significant amounts of siliciclastic material, (up to 10 %) and organic matter (up to 6% of TOC; Fig. 4). In the northern margin, some massive layers occurred at the feet of the steep scarps. The lacustrine sequence in Lake Taravilla contains meter-thick fining upward sequences with three components (Fig. 5): i) decimeter-thick, massive dark

grey carbonate-rich silty-sand to coarse sand; ii) centimeter- to decimeter-thick, massive to faintly laminated light grey carbonate silt and iii) centimeter to decimeter-thick, massive to faintly laminated, dark grey carbonate silt and mud. Rarely centimeter-thick silt layers with abundant plant macroremains are also deposited. Coarse facies (gravels and sands) are relatively common in La Parra sequence (Fig. 6) and include dm-thick massive gravel beds, cm-thick fine gravels and coarse sands and cm-thick, massive fine sands arranged in fining-upward sequences. Coarse facies are minor in topographically – closed basins with no surface inlet. In Lake Enol, they only occur as cm-thick thick carbonate sandy layers with abundant shell fragments.

Interpretation: Coarse carbonate facies arranged in fining-upward sequences and with sedimentary structures indicative of sub-aqueous transport (lamination) represents deposition in littoral zones with a strong alluvial/deltaic influence. They are better developed close to the surface inlets feeding the lakes in Taravilla and Banyoles. Massive layers at the feet of steep scarps (e.g., northern margin of Lake Arreo) are interpreted as debris deposits related to cliff erosion.

2. Massive to banded carbonate – rich silts and muds (LF).

Carbonate-rich silts and muds are the most common facies in Iberian karst lakes. This group of facies shows a large variability depending on stratification features (massive, banded and laminated), amount of carbonates and siliceous minerals and the abundance of littoral biota and flora remains. If present, sedimentary structures indicate current or wave action. In Lake Taravilla, deposition in littoral areas close to the tufa barrage is characterized by massive to faintly laminated, dark grey, organic-rich (up to 10% in organic matter) carbonate

silts with gastropods and mollusks and charophyte remains (Fig. 5). Finer facies are deposited in deeper areas. Most littoral sediments in Lake Banyoles are made up of these facies (Valero-Garcés et al., 1998; Höbig et al., 2012). In Lake Enol, Holocene sediments are composed of banded to finely laminated coarse silts with variable carbonate content (mostly coarse-silt calcitic detrital particles), silicate minerals, macrophytes and terrestrial organic matter (Fig. 13). In Lake Arreo the sequence from the littoral core is composed of alternating fine and coarse silts with some intercalating clastic carbonate silts with fining upwards texture (Fig. 4). In Lake Montcortès, cm-thick, greyish-brownish layers of carbonate silty mud with abundant organic matter, *Chara* and gastropod fragments are indicative of deposition in carbonate-producing littoral environments (Corella et al., 2011b). Similarly to Lake Arreo, some facies are dominated by grey, silty, carbonate layers, pointing to higher carbonate clastic input from the drainage basin and re-working of carbonates in the littoral platform, while other are richer in organic matter suggesting higher organic productivity in the littoral zone. In Lake Estanya, banded to laminated dm-thick silt layers are mainly composed of calcite as well as minor amounts of quartz and clay minerals (Fig. 11). Carbonates are biogenic grains (*Chara* fragments, micrite oncoids, carbonate coatings) and small crystals derived from the re-working of particles produced in the littoral environments, although some euhedral morphologies point to direct precipitation in the waters. In Lake Basa de la Mora, carbonate-rich silts (2 – 7% TIC) with variable, but relatively high organic matter content (TOC 1-3%) and relatively low magnetic susceptibility, are arranged in massive to banded deposits with thin intercalations of organic-rich silt (Fig. 7).

Interpretation. Banded silty mud and carbonate silty sand with abundant plant remains, gastropods, mollusks and charophytes reflect deposition in a carbonate-producing littoral environment with restricted clastic input from the watershed. The lack of sedimentary textures is interpreted as a result of bioturbation in well-oxygenated, shallow waters. Banded sediments represent re-working and re-deposition of the charophyte meadows and other carbonate-encrusted littoral sediments. These facies are similar to marl/micrite facies ubiquitous in littoral settings of carbonate lakes (Magny et al., 2006; Magny et al., 2007; Renaut and Gierlowski-Kordesch, 2010). Finer silts with faint lamination and less fossil content suggest relatively deeper, littoral to sublittoral depositional environments. In more profundal areas, darker, massive to faintly laminated silt and mud represents deposition during periods of low energy, less fluvial influence, restricted water mixing, and episodic occurrence of anoxic bottom conditions. Massive or graded silty layers deposited in the littoral realm during periods of higher run-off in the watershed.

3. Massive to banded carbonate – rich silts and muds with mottling and evaporite minerals (gypsum) (LFs).

These facies are similar to the previous LF facies but may include gypsum nodules, other carbonates minerals (aragonite, dolomite, Mg-calcite) and bioturbation features (root traces, coarse plant remains, mottling). Good examples occur in the Lake Arreo and Zoñar sequences (Figs. 4 and 9).

Interpretation. These facies deposited in shallow, sometimes ephemeral environments and were affected by prolonged periods of subaerial exposure, soil development and evaporative mineral formation. They are similar to some of the

palustrine facies identified by Alonso-Zarza and Wright (2010) as mottled, nodular and brecciated, and with root cavities.

4.3. Distal-Profundal Facies Associations

Facies

1. Banded to laminated carbonate silts (DB)

These facies are the most common in open – water, relatively deep environments but include a large range in grain size (from fine to coarse silts), sedimentary textures (banded to laminated), color (grey, black, brown) and composition (from carbonate-rich to silicate-rich). Although biogenic components (gastropods, mollusks, *Chara*) are present, they are less abundant than in littoral facies. Cm- to dm- thick sequences are usually defined by grain size variability. Examples of these facies occur in most lakes, such as in Lake Basa de la Mora, where thin carbonate-rich massive silts were deposited during the late Holocene (Fig. 7). Sedimentation in the distal areas of Lake Estanya during the last 1000 years (Morellón et al., 2009a; Morellón et al., 2011) (Fig. 11) and Lake Banyoles (Fig. 14) are also dominated by banded carbonate silts.

Interpretation. This facies association represents deposition in the distal open water areas of permanently or seasonally oxygenated lakes. They include the massive (structureless) and laminated carbonate muds and silts from open water – offshore facies (Renaut and Gierlowski-Kordesch, 2010).

2. Finely laminated (DL)

Finely laminated sediments (layers thinner than 1 cm) are the most characteristic lacustrine facies. They vary in thickness and lateral extent (continuous to discontinuous) and form when sediment supply changes with time

(seasonally, annually) and preservation conditions are met (Glenn and Kelts, 1991). In the Iberian lakes we have identified three main facies associations: i) annually laminated (varves), ii) finely laminated (laminites), and iii) rhythmites,

2.1. Varves (DLV). Intervals with finely, laminated facies occur in most lake sequences, although only in some (Lakes Zoñar, Montcortés, Arreo, La Cruz) they are annual in nature. We have identified two main types:

- With calcite laminae (DLV(c)). These facies occur in most karst lakes, but are best developed in Lakes Montcortès and Zoñar (Figs. 3, 8, 9 and 10). They are composed of mm-thick triplets of three laminae: i) white layers, mostly made of rhombohedral calcite crystals of variable (4–20 μm -long) sizes, ii) greenish-brownish, organic-rich laminae with abundant diatoms, amorphous organic matter and variable detrital content and iii) clastic grey laminae made up of allochthonous carbonate, clay minerals, and quartz grains, commonly with a fining-upward texture. Grey laminae are not always present. In some intervals, varves are poorly preserved; laminae are thicker and clastic laminae more common and thicker too. Varved facies occur through the whole sequence in Lake Montcortès (last 5000 years) (Corella, 2011b) (Fig. 8). In Lakes Zoñar, Arreo, El Tejo, La Cruz, El Tobar and La Parra laminated facies are restricted to specific intervals and in Lake Banyoles only in the northernmost sub-basin. In Lake Zoñar they are well developed in the lower part of the sequence (2600-1600 cal yr BP) (Fig. 9) and some intervals contain varves poorly preserved and without the organic-rich layers (Martín-Puertas et al., 2008). In Lake Arreo, varves composed of mm-thick triplets of endogenic calcite, organic, and detrital laminae frequently occur in the whole sequence of the distal core (Corella et al., 2011a; Corella et al., 2013a) (Fig. 12).

Several microfacies can be distinguished in the calcite layers (Fig. 10). In Lakes Montcortès and Zoñar there are three main types: i) fining upward (C-F), with a lower sub-layer of coarser calcite crystals (averaging 12 μm in thickness) and an upper sub-layer of finer crystals (average length of 5 μm), ii) coarsening-upward (F-C), and, iii) homogeneous coarse crystals. Most of the sub-layers show an abrupt boundary, indicating two distinct calcite precipitation pulses (Martín-Puertas et al., 2008; Corella et al., 2012). In Lake Zoñar, *Botryococcus* blooms are associated with the calcite sub-layer (Fig. 10b). Two main microfacies also occur (Martín-Puertas et al., 2009): (1) microfacies 1c is composed of thicker calcite layers (1-2 mm), higher crystal density and smaller calcite grain size (5-7 μm) than (2) microfacies 1a (0.1-0.8 mm and 10-15 μm).

- without calcite laminae (DLV(o)) In Lake Montcortès this facies is composed of couplets of two laminae: i) mm-thick, brown to greenish, organic-rich laminae with higher content of amorphous aquatic organic matter, diatoms and clay minerals, and ii) mm-thick terrigenous grey silt layers composed of clay minerals, carbonate, and quartz grains (Fig. 3l). This laminated facies has the highest TOC values in the sediment sequence (up to 18%) and thicker diatom- and organic matter-rich laminae, suggestive of high biological productivity and organic matter preservation in the lake. In Lake Zoñar, there is also a finely laminated microfacies (microfacies 1b) characterized by the absence of a calcite layer (Fig. 9). In El Tobar some of the finely-laminated organic facies are made of detrital and organic – rich laminae, but the annual nature has not yet been demonstrated.

Interpretation. The development of long periods of anoxia, possibly related to the development of temporary meromictic conditions, the strong seasonality of the

518 climate in Mediterranean areas, and the occurrence of algal blooms in spring and
519 summer allowed the formation and preservation of biogenic varves in these small,
520 relatively deep karst lakes. Varves with three sublayers correspond to the classic
521 biogenic varves described in lakes located in carbonate bedrock (Brauer, 2004).
522 Varves have also been described in other settings: volcanic lakes (e.g., Eifel maar
523 lakes (Brauer, 2004); Nar Golü, Turkey, (Jones et al., 2006)); glaciated terrains of
524 British Columbia, (Renault and Last, 1994), and coastal lagoons (Ariztegui et al.,
525 2010). Calcite crystals originate from endogenic precipitation in the lake epilimnion
526 caused by ion saturation enhanced by algal blooms and increasing water
527 temperatures (Brauer, 2004; Kelts and Hsü, 1978; Romero-Viana et al., 2008).
528 Epilimnetic formation of calcite during spring/summer is typical in lakes emplaced
529 in carbonate-rich bedrock areas (Brauer et al., 2008). Similar late spring to summer
530 calcite precipitation has been described in hard water lakes in Germany (e.g.,
531 (Koschel, 1997)), Switzerland (e.g., (Kelts and Hsü, 1978)), and the Iberian
532 Peninsula (Lake La Cruz, (Julià et al., 1998; Romero-Viana et al., 2008)). Calcite
533 grain size and texture give evidence on past temperatures, seasonality, and
534 anthropogenic activities in the lake's catchment (Teranes et al., 1999a; Teranes et
535 al., 1999b). Occurrence of larger calcite grain size has been interpreted in some
536 lacustrine environments as precipitation during spring season responding to higher
537 phosphate content (Teranes et al., 1999a; Teranes et al., 1999b). Thicker calcite
538 laminae and smaller crystal size may also suggest either that saturation conditions
539 or organic productivity periods lasted longer (warmer summers) or that the
540 concentration of calcium and bicarbonate ions in the waters was higher (increased
541 supply due to higher winter precipitation and aquifer recharge) as shown in Lake

La Cruz (Romero-Viana et al., 2008). Reduced calcite thickness and development of fine to coarse (F-C) calcite sub-layering in Lake Montcortès is interpreted as colder temperatures and prolonged winter conditions (Corella et al., 2012).

The organic-rich lamina represents deposition after the period of calcite precipitation. The variable content in clay minerals within the organic lamina would reflect seasonal changes in sediment delivery to the lake, likely controlled by runoff and rainfall. The detrital laminae were deposited when sediment delivery from the watershed to the lake was intensified during stronger, rainier periods. The thickness and number of those layers contain information on storm seasonality and frequency (Corella et al., 2012).

2.2.Laminites (DLL). Finely laminated facies with no clear indication of annual lamination are included in this category. They are gypsum-rich -DLL(g)- and/ or organic – rich DLL(o) - facies.

Gypsum is a common mineral in Iberian karst lakes because of the presence of evaporites in the bedrock, which often constitutes the karstified substrate. The higher salinity of the groundwaters and the strong evaporation in some of these hydrologically – closed lakes is conducive to the formation of gypsum. Gypsum-rich facies DLL(g) in open water environments are dominated by endogenic gypsum crystals and they occur in Lakes Estanya, Zoñar, and Arreo. In Lake Arreo, these laminated facies are composed of mm-thick triplets (average thickness 2-3 mm) of organic matter, terrigenous, and gypsum layers with endogenic prismatic gypsum crystals (Fig. 3E). Calcite laminae are also present in some triplets (Figs. 3F and 12).

Lake Estanya contains the only sequence located in the Iberian Peninsula with a thick interval of finely-laminated gypsum layers; these facies contain prismatic and nodular gypsum and intercalate with carbonate mud layers with a mixed mineralogy (aragonite, calcite, high -Mg calcite, dolomite) (Fig. 11). In Lake Zoñar, gypsum – rich facies occur as cm-thick layers intercalated in the varved facies (Fig. 9). They are composed of gypsum crystals around 10-30 µm long, also appears as irregular cm-thick nodular gypsum layers, and cm-thick layers of carbonate silt with mm-long gypsum nodules and isolated longer gypsum crystals.

There are two types of finely laminated organic-rich facies DLL(o): sapropels and microbial mats. In Lake Estanya, both facies are present and always associated with gypsum. Brown, massive sapropel layers associated with gypsum laminae composed of idiomorphic, well-developed crystals ranging from 25 to 50 µm in length are common in the Holocene sediments. Cm-thick microbial mats also occur (Fig. 11). In Lake Zoñar, microbial – algal mats only occur in a thin interval in the upper part of the sequence and they are not associated with evaporite minerals (Fig. 9). These microbial- algal mats are the only microbial carbonate facies (after (Gierlowski-Kordesch, 2010) in Iberian karst lakes.

Interpretation: The pyramidal shapes, the homogeneous size of crystals, and the random distribution of the crystals in the DDL(g) laminae suggest gypsum precipitation within the water column (Smoot and Lowenstein, 1991), a consequence of chemically concentrated lake waters and saturated conditions for sulfates. Interestingly, the presence of planktonic saline diatoms suggests relatively high lake levels during deposition of these facies in Lake Estanya (Morellón et al., 2009a). The nodular textures indicate gypsum formation from the sediment

interstitial brine waters (Hardie et al., 1978) that may occur in both, shallow ephemeral settings and relatively “deep” and concentrated hypolimnetic waters in meromictic saline lakes (Last, 1994).

Organic-rich sediments and saline minerals (gypsum) point to deposition in lake environments with a high microbial/algal bioproductivity and strong evaporation processes for DDL (o). Preservation of laminated facies in shallow lakes occurs commonly in saline environments which reduced bioturbation (Last, 1990; Schreiber and Tabakh, 2000).

2.3. Rhythmites (DLR). Laminated, siliciclastic carbonate-poor facies only occur in high altitude lakes with a mix siliciclastic and carbonate provenance (Lakes Basa de la Mora, Enol and Marboré) and where glacier and periglacial processes occur. In Lake Enol laminated facies (up to 4 % TIC) occur in the pre-Holocene sequence (Moreno et al., 2011) (Figs. 3J and 13). Lamination is composed of three laminae: (a) grey, clay – rich, (b) light brown with 10-15% of silty calcitic grains, and (c) brown, coarser with calcite and siliciclastic particles. In Lake Marboré, the Holocene sediments are made up of banded to laminated, carbonate-poor, coarse and fine silts. Carbonate content is low (TIC < 1 %) and composition is dominated by quartz and clay minerals, with low organic matter content. Finer-grained laminae are lighter in color and coarser laminae are darker with more biotites. In Lake La Basa de la Mora, carbonate – poor (< 2 % TIC) sediments, with also low TOC and MS occur in laminated or banded intervals of fine silts (Fig. 7). Although all these lakes are present in carbonate bedrock, TIC percentages are surprisingly low (2-3 %).

Interpretation. Sedimentological and textural features of the Lake Enol facies are similar to proglacial lake sediments (Leonard and Reasoner, 1999; Ohlendorf et al., 2003). They are interpreted as rhythmites deposited in lakes fed by glacier meltwater with a strong seasonality: the calcite-rich, coarse silt laminae deposit during the melting season and the fine-grained clay-rich laminae during the winter, when the lake was ice-covered. Coarser laminae were deposited during years with stronger melting pulses. In proglacial environments, better development of laminated sediments would occur during periods of stronger seasonality with higher melting and run-off discharges during spring and summer and longer ice-covered winters (Ohlendorf and Sturm, 2001; Ohlendorf et al., 2003).. More massive facies are finer-grained and clay-rich with no carbonate. They would deposit during periods with less marked seasonality (colder summers), less available water for run-off when only fine glacial sediment were mobilized ("glacial flour"). Higher carbonate content in Lake Enol proglacial sediments is clearly associated with coarser sediment fraction since finer fraction includes more clay minerals (Moreno et al., 2011).

In Lakes Marboré and Basa de la Mora, lamination reflects changes in annual clastic input to the lake and changes in the seasonality: coarser clastic materials are delivered to the lake during the summer and finer sediments are deposited during the ice-covered winter.

3. Massive and Graded Facies

Cm- to dm-thick massive (DM) and graded (DG) facies are common in distal areas of karst lakes. Graded facies are common in relatively deep lakes, such as Taravilla, Montcortés, Arreo, Estanya and El Tobar. Both facies are characterized

by high magnetic susceptibility values, graded textures, and mixed carbonate and silicate composition. In most lakes the facies occur as i) mm-thick laminae and ii) cm- to dm-thick layers. The cm- to dm-thick layers show fining-upward textures, erosive basal surfaces with abundant plant remains, and a sandy basal sub-layer. Some layers incorporate sub-angular calcite particles in the base, likely related to the re-working of older carbonates in the catchment.

Graded facies are particularly well developed in Lake Taravilla deposits, where they occur as up to dm-thick massive, graded, fining-upward layers, ranging from coarse sands with plant remains at the base to silts at the top (Valero Garcés et al., 2008) (Figs. 3A and 5). In Lake Arreo, massive, cm-thick silty sand layers with abundant plagioclase and pyroxene crystals, erosive basal surfaces, and high magnetic susceptibility values occur in the profundal areas (Fig. 4). In Lake Zoñar these facies are less common and occur as cm-thick layers composed of massive to graded coarse sand and silt sediments with abundant amorphous aquatic and terrestrial organic matter remains (Fig. 9).

Homogeneites facies (DH) only appear in the Lake Banyoles sequence (Fig. 14). They occur as up to 75 cm thick homogeneous layers composed of fine calcitic mud, intercalated within distal, blackish, massive fine-grained silts in the cores located in the 20 m deep flat platforms surrounding the deepest and most active sinkholes.

Interpretation. The sedimentological features of graded and massive facies are characteristic of turbidite-type or storm-related deposits (Mangili et al., 2007; Moreno et al., 2008). The graded nature indicates deposition by turbidity currents that separate the coarse bed load from the fine grains in suspension. The coarse

basal layer and the erosional surfaces likely result from underflow current processes (Sturm and Matter, 1978). Fine particles with a fining upwards texture would have been deposited by settling afterwards. The presence of reworked carbonate material may indicate erosion of the littoral platform during the periods of more intense flooding in the watershed. The fine, massive texture of some facies indicates rapid deposition in distal areas of the lake of suspended clay-rich materials that were transported by creeks that drain the catchment during flooding episodes. These facies are similar to those described in alpine lakes (Wilhelm et al., 2012). In Lake Taravilla, turbidites DG result from delta collapse episodes associated to flood events in the lake (Moreno et al., 2008). In Lake Arreo massive facies DM are similar to recent deposition of massive, sandy facies related to documented scarp failures in the ophyte outcrops located along the northern shore of the lake.

Homogeneites DH facies are only present in cores recovered from the platforms surrounding the deepest sinkholes in Banyoles with active groundwater flow processes. Morellón et al (in review) propose that sediment fluidification events during periods of higher groundwater discharge are likely responsible of the episodic deposition of these homogeneous layers in the internal platforms. This depositional mechanism has been documented during the last decades (Colomer et al., 2002; Serra et al., 2002; Serra et al., 2005): after intense and prolonged rains in the catchment, groundwater flow in the depressions increases, sediments are mobilized and transported by turbidite currents sweeping over the southern platforms of Lake Banyoles. The occurrence of homogeneites in the Banyoles

sequence demonstrate that groundwater activity may be a significant factor controlling sediment deposition in karstic lakes.

4. Gravitational deposits facies (G)

In several sequences (Lakes El Tobar, Montcortès, Enol) discrete intervals show convoluted and disrupted textures with folds, microfractures, and microslides (Facies GF). Coarser, chaotic facies with cm long sandstone and limestone clasts within a sandy matrix also occur at the base of some gravitational units in Lake Montcortès (Facies GC).

In particular, Lake Montcortès provides a remarkable example of several major slide units, delimited by surfaces with little deformation and slide structures (Corella et al., 2011b) (Fig. 8). In Lake Estanya a large mass flow (5 m thick, 150 m in maximum length) was identified in the deepest sub-basin by a geophysical survey (Morellón et al., 2009a) coinciding with a major change in sedimentation, from organic-rich into clastic-rich sediments (Fig. 11). Smaller deposits occurred in the north-western basin, likely associated with early stages of development of the lake basin. Gravitational deposits have also been identified in Lake Enol by multiple core correlation (Fig. 13) (Moreno et al., 2010; Moreno et al., 2011).

Interpretation. Convoluted and chaotic facies are interpreted as a result of gravitational processes affecting fine laminated or coarse lacustrine deposits. Local instability in submerged sediment-covered slopes (Strasser et al., 2007) originates from various processes such as erosion, rapid sedimentation, gas release or migration and earthquake shaking (among others) (Hampton et al., 1996; Locat and Lee, 2002; Girardclos et al., 2007; Corella et al., 2013b). Mass wasting processes are generated when gravitational downslope forces overcome the

static threshold of the shear stress in the sediments (Hampton et al., 1996). These processes are frequent in lakes with steep margins (Chapron et al., 2004; Girardclos et al., 2007).

4.4. Lacustrine environments

Carbonate sedimentary facies in Iberian karst lakes group in three main environments (Fig. 15): palustrine, littoral-sublittoral and distal-profunda

Palustrine environments are associated with hard-water lakes and wetland settings (Alonso-Zarza and Wright, 2010). The Florida Everglades, USA (Platt and Wright, 1991) and the Tablas de Daimiel wetlands, Spain (Alonso-Zarza et al., 2006) are examples of extensive palustrine environments. In Iberian karstic lakes, they occur in the transitional zone between the submerged littoral and the emerged areas surrounding the lake. Their surface area is controlled by lake basin topography and hydrology and they are characterized by fluctuating water levels with alternating periods of flooding and subaerial exposure. They are better developed in lakes with an extensive shallow area as Lakes Taravilla and Arreo. More than half of the surface area of the Lake Taravilla lake basin is occupied by a palustrine environment (Fig. 2B and 7) while a significantly large wetland occurs in the southern part of Lake Arreo (Fig. 2I and 5). In these relatively large shallow areas, vegetation develops and organic matter production and accumulation processes led to deposition of organic-rich facies and peat layers as in Lake Taravilla (Valero Garcés et al., 2008), Arreo (Corella et al., 2013a) and Banyoles (Höbig et al., 2012). Lakes with inlets have littoral carbonate and siliciclastic facies intercalated with the carbonate and organic-rich facies. Coarse palustrine carbonates are composed of extraclasts from the watershed, endoclasts (bioclasts,

coated grains, intraclasts, Chara fragments, shell material) and endogenic carbonate precipitated by microbial and algal activity (Alonso-Zarza and Wright, 2010). The vegetation belt stabilizes the substrate and acts as a barrier for allocthonous material transported into the lakes by run-off as shown in many case-studies (Platt and Wright, 1991; Alonso-Zarza et al., 2006; Alonso-Zarza and Wright, 2010).

The littoral environment constitutes a relatively shallow flat to sloping area, partially colonized by vegetation, that provides support for epiphytic fauna, and largely contributes to the production of carbonate particles (Renaut and Gierlowski-Kordesch, 2010). In the Iberian lakes, the extension of the present-day littoral areas differs enormously in size, although sedimentological and compositional features of the littoral facies in all lakes are relatively similar. In Lake Estanya there is an internal littoral platform, 5 to 10 m wide, extending from the vegetation belt to the lake shoreline. Sediment is mainly composed of light grey, massive, bioturbated (root casts, worm traces, mottling sediment textures) carbonate – rich (up to 10 % TIC) coarse silts with abundant plant fragments. The external littoral platform (0 to 4.5 m water depth) permanently submerged between the shoreline and the slope is colonized by macrophytes and charophytes. In Lake Arreo, the littoral realm occupies > 60 % of the lake total area mostly in the southern area of the lake and it is very narrow in the northern side due to the presence of a steep margin. In contrast, the steep margins in Lake Montcortès limit the littoral, carbonate-producing sub-environment to 10% of the lake total area. In Lake Taravilla, a very narrow littoral zone developed with thick aquatic vegetation overhanging at the edge of the steep lake margins and submerged plants coated

with carbonate (Valero Garcés et al., 2008).

This littoral environment is the main carbonate factory in the Iberian karstic lakes, comprising biogenic carbonates (ostracods, gastropods and *Chara* sp. particles) and non-biogenic carbonates (coatings around submerged macrophytes and the lake substrate). Detrital carbonates are also present as a result of re-working of the endogenic carbonate and also as clastic sediment delivered from the watershed. Storms and wave activity lead to the re-working of these particles, as indicated by the rare presence of ripples (Morellón et al., 2008; Corella et al., 2013a)..

Sedimentation in littoral areas is mainly governed by physical processes (waves and currents) (Imboden, 2007) and by changes in lake levels (Renaut and Gierlowski-Kordesch, 2010). The oscillations in the hydrological dynamics would play an important role in these sensitive environments as it is directly related to water depth, chemical composition, and light penetration. Thus, increasing water salinity and/or reduced light penetration would control the existence of different biota sensitive to changes in the limnological conditions (Wright, 1990). However, the sediment input into the lake, controlled by hydrology, climate (increased precipitation), and human activities (changes in land use) would also affect the water turbidity – which is also a limiting factor in the development of algal blooms (Reynolds and Walsby, 1975) - and nutrients inputs into the lake. The combination of these factors (fluctuating lake levels, water chemistry, clastic input) are inter-related and would respond to internal thresholds that triggers the changes in the lateral and vertical sedimentary facies distribution in littoral environments (Wright, 1990; Valero-Garcés and Kelts, 1995).

779 There are numerous examples of recent and Holocene carbonates deposited
780 in littoral settings. The classical bench margins models of Murphy and Wilkinson,
781 (1980) and Treese and Wilkinson (1982) contain charophyte – rich facies with
782 numerous intraclasts and rippled carbonates. Tucker and Wright (1990) and Platt
783 and Wright (1991) present facies models for carbonate shorelines depending on
784 the slope (benches in moderately steep margins and ramps in gentler slopes) and
785 the energy conditions. Littoral settings in Iberian karst lakes are always low-energy,
786 but they present similarities with both: steep margins (bench) as in Lake
787 Montcortés and gradual margins (ramp) in Lakes Taravilla and Arreo. Low-energy
788 bench margins contain massive and banded carbonate facies in the littoral zones
789 and bedload – transported carbonates with slumps, turbidites, and re-sedimented
790 carbonates from the littoral zone. The mid-Holocene Lake Montcortès sedimentary
791 record (Corella et al., 2011b) is an example of this type of margin. The Lake
792 Taravilla and Arreo littoral sequences are examples of low energy ramp lake
793 margins with palustrine facies, subaerial exposure textures, and some fluvial
794 intercalations.

795 The distal-profundal environments include the deeper zone of the transitional
796 talus and the central, deepest, and relatively flat areas. The transitional talus is a
797 narrow area characterized by steep morphology, limited presence of vegetation
798 due to the lack of light, and the occurrence of small mass movements as a result of
799 sediment destabilization. Transport processes are dominant over sedimentation in
800 this spatially-restricted environment. Carbonates originating in the littoral platform
801 are transported to the talus and the distal areas. Slope instability in the talus is
802 more common in deeper lakes with steeper morphologies where small and large

slides and mass wasting deposits have been identified with seismic and multiple core stratigraphic analyses (Lakes Estanya, Montcortès and Enol). In addition, turbidite currents frequently remobilize and erode talus sediments and bring the eroded material to distal areas.

Distal areas are characterized by the deposition of darker, massive to laminated, fine-grained carbonate silts. Laminated fine sediments occur in the distal areas of all studied lakes. Carbonate content is variable (0.1-10 % TIC) reflecting the distance to the producing littoral areas and to the dissolution processes that remove small carbonate particles. According with the predominantly oxic or anoxic conditions at the bottom of the lake, and the salinity, three main sub-environments are identified: i) Dominantly oxic hypolimnion with only seasonal anoxic conditions characterized by the deposition of banded carbonate silts; ii) Anoxic, freshwater hypolimnion, with deposition of varves (Lake Montcortès); iii) Anoxic, saline hypolimnion with deposition of laminated facies with gypsum (Lakes Arreo, Tobar, Estanya)

Seasonal or permanent anoxic hypolimnetic conditions greatly reduce bioturbation processes and enhance organic matter preservation (up to 8 % TOC) and sulfide formation. Recent deposition of finely laminated sediments occurs in Lakes Montcortès, Arreo, and La Cruz. Precipitation of small calcite crystals in the epilimnion associated with algal bloom seems to be a smaller contributor to total carbonate production in most lakes.

The size of these sub-environments differs largely in comparison with the littoral sub-environments (80 % of the total surface in Lake Montcortès and 15% of the total area in Lake Arreo). The morphology of these basins consisting of funnel-

shaped dolines with steep margins and a high depth/area ratio and the limnological and hydrological features lead to the development of anoxic conditions during most of the whole annual cycle, with limited bottom bioturbation, allowing the preservation of finely laminated sediments (O'Sullivan, 1983; Brauer, 2004; Zolitschka, 2007). These facies occur in several Iberian lakes (e.g., Lake La Cruz; Julià et al., 1998; Romero-Viana et al., 2008), Lake Zoñar, (Valero-Garcés et al., 2006; Martín-Puertas et al., 2008b), Lake Banyoles, meromictic sub-basins.

Bottom morphology in distal areas of most lakes is flat because sediments have covered the original topography having more irregular karstic features (sinkholes). In active karstic lake systems, such as Lake Banyoles, lake bed morphology is more complex. Several distal, flat platforms (ca. 20 m water depth) occur between the main sinkholes (>30 m deep) where fluidized fine-grained sediments of varying densities are dominant. These sinkhole areas are characterized by flat bottom in seismic surveys (Canals et al., 1990; Morellón et al., in review) that are the uppermost surface of suspensate sediment clouds sustained by upwards phreatic inflow of warmer waters.

5. Discussion: The carbonate factory in Iberian karst lakes

5.1. Sources and Processes.

Carbonates are the main sediment component of Iberian karst lakes, except those located in glacial/periglacial settings. Carbonate minerals are formed in most depositional environments in the lake systems as: i) tufa deposits associated with springs ii) coatings in macrophytes and aquatic vegetation in palustrine and littoral settings, and iii) endogenic precipitates in the epilimnion. Carbonate particles are

also transported into the lake as iv) alluvial/fluvial and v) aeolian input. Finally, early diagenetic processes within the sediments (authigenic minerals) and internal re-working also contribute to carbonate deposition in lakes.

Carbonates in Iberian karst lakes have three main origins: chemical, biogenic and clastic (alluvial and aeolian)

i) Chemical formation. Carbonate precipitation as a consequence of direct chemical water concentration has been considered a main process in many freshwater and saline lakes (Eugster and Kelts, 1983; Last and Smol, 2001). Waters may become supersaturated in calcite (or other carbonate phases) because of changes in temperature and increased evaporation. However, even with supersaturated waters, microbial or algal activity in the epilimnion is necessary to trigger calcite formation (Stabel, 1986; Gierlowski-Kordesch, 2010). In littoral and palustrine settings, with larger changes in chemistry caused by lake level variations and stronger evaporation, direct precipitation could occur. However, precipitation of carbonate minerals in shallow marginal areas is also predominantly biochemical, mediated by microbes, macrophytes, algae or even fauna (Sanz-Montero et al., 2008; Gierlowski-Kordesch, 2010;). Some of the coatings and thin crusts in littoral vegetation in Lakes Estanya (Morellón et al., 2009a) and Arreo (Corella et al., 2013a) may respond to this process. Precipitation of carbonate nodules in mudflat – type environments with subaerial exposure and pedogenic processes have occurred in Lakes Estanya and Zoñar during low lake levels at the onset of the Holocene and between 4 – 2.5 ka BP, respectively. Aragonite formation in such ephemeral and saline settings (Lake Zoñar, Martín-Puertas et al., 2008) could be mostly chemically – controlled when waters reached a high (>1)

Mg/Ca ratios (Hardie et al., 1978). Occurrence of high magnesium calcite and dolomite in laminated intervals in Lake Estanya could reflect the effect of evaporation and chemical water – enrichment during periods of increased aridity (Fig. 11). However the association with microbial mats suggests that biomediation plays a significant role as a whole; direct chemical precipitation is a minor process in carbonate formation in Iberian karst lakes, and most likely microbe activity could be responsible for most of the endogenic carbonate formation.

ii) Biogenic. Biogenic - mediated carbonate precipitation occurs in all depositional environments in karst lakes. In littoral areas, shells (bivalves, gastropods, ostracods) are a significant fraction of massive and banded facies in all lakes. Charophyte remains are a particularly significant component in Lakes Taravilla, Estanya, Banyoles and Arreo where *Chara* meadows extend to deeper waters. Carbonate coatings around plants and nearshore vegetation are major contributions in lakes with a relatively large littoral zone (Lakes Estanya and Arreo). Although old tufa buildups are present in some of the watersheds (Lakes Taravilla, and Banyoles), and subaerial and subaqueous springs occur in all of them, no spring – related deposits have been found in any of the lakes. The absence of carbonate accumulations associated to seepage of spring zones could be caused by the similar hydrochemistry of groundwaters and lakewaters.

Calcite laminae in varved facies demonstrate the widespread precipitation of calcite in epilimnetic waters. Whittings have only been observed in La Cruz (Romero et al., 2006; Romero-Viana et al., 2008; Romero-Viana et al., 2011). Sediment traps in Lakes Estanya, Zoñar and Banyoles became quickly covered with algal and microbial mats encrusted with calcite. Varve microstructure and composition

indicate a clear connection with biological activity in Lakes Montcortès (diatom blooms) (Scussolini et al., 2011; Corella et al., 2012;) and Zoñar (*Botryococcus* blooms) (Martín-Puertas et al., 2008). Laminated facies composed of irregular organic (microbial mats) and calcite layers as those in the Holocene in Lake Estanya also suggest calcite formation in shallower, saline settings associated with microbial activity (Morellón et al., 2009a).

iii) Clastic. Lake sediments greatly reflect the geology of the watershed. Karst lakes are developed in carbonate terrains and, consequently, the erosion of surface and bedrock formations is responsible for the observed dominance of carbonates in these lake sediments. However, the amount of carbonate reaching the lake from the catchment depends on several factors: i) the size of the watershed, ii) the availability of erodible materials and iii) the existence of an organized drainage network. The availability of sediments in the catchment area is determined by the size of the watershed, geology, climate, vegetation cover, and in more recent times, land uses and human impact. In Lake Taravilla, the large tufa build-ups in the catchment are one of the main suppliers of carbonate particles to the lake. Lakes with small watersheds and no surface drainage network, such as Lakes Tejo and La Cruz, have reduced clastic input, mostly as coarse materials derived from the littoral zones. Lakes of similar size but with a relatively larger watershed and a small ephemeral inlet like Lake La Parra, have more clastic coarser facies. Lakes with intermediate watersheds (Lakes El Tobar and Montcortès) have more clastic facies, particularly fine-grained. Lakes with larger watersheds (Lakes Banyoles, Zoñar and Taravilla) have the highest percentages of clastic facies. Evidence of flood events and their sedimentological signature

(turbidite- type facies) are found in the distal facies associations of all Iberian karstic lakes.

A unique characteristic of Iberian lakes is the large increase in clastic input during historical times, particularly since the Medieval epoch, caused by deforestation and changes in land uses in the watershed (Valero Garcés et al., 2009).

Silt-size particles transported by wind are likely an input in karst lakes located in areas with strong prevalent winds. However, the identification of such particles in the lacustrine or marine sediments by optical or chemical techniques is not easy (Moreno et al., 2002). One of the areas where aeolian processes are very significant is the Central Ebro Valley, where a number of saline lakes occur (Valero-Garcés et al., 2000; Moreno et al., 2004). The origin of these lakes includes dissolution and karstification of the Miocene carbonates and evaporites and also aeolian activity (Gutiérrez et al., 2013). In this setting, aeolian processes are responsible for the erosion and transport of large amounts of sediments: erosion of Holocene terraces suggests a specific rate of $30\text{m}^3/\text{ha}/\text{yr}$ (Gutiérrez et al., 2013). However, most Iberian karst lakes are located in relatively well-vegetated areas where wind erosion and deflation processes are likely to be secondary.

5.2. Controls on carbonate deposition in karst lakes

Iberian karst lake sequences are characterized by a large variability of facies and depositional environments with abrupt lateral and vertical changes. As in most lakes (Valero-Garcés and Kelts, 1995; Renaut and Gierlowski-Kordesch, 2010),

the main factor controlling sedimentation in Iberian karstic lakes is hydrology, influencing intensity of karstic processes, hydrochemistry, thermal regime and sediment delivery to the lake. We have identified five main parameters on facies distribution and carbonate deposition, all closely connected with hydrologic dynamics: intensity of karstification processes, basin morphology, hydrological balance and surface hydrology, watershed, and climate. Figure 16 summarizes some of the main physical and chemical characteristics of the selected lakes.

Karstification processes are not only responsible for the origin of the basins through collapse, subsidence, and dissolution of carbonate and evaporite formations, but they also effectively control the morphology of the basin and the dynamics of the hydrogeological system. Initial lake formation is directly related to mechanical and dissolution processes that generate the accommodation space for the lake (Kindinger et al., 1999; Gutiérrez et al., 2008). According to Kindinger et al.'s (1999) classification of progressive developmental phases of sinkhole lake formation, the Lake Banyoles with at least thirteen active springs and formation of dolines as recent as 1978 on the western shore (Brusi et al., 1990; Julià, 1980; Moner et al., 1987) could be considered the early stages of karst lake development. Most of the Iberian karst lakes are in a mature phase, when the sinkhole has been plugged with sediments. In larger karstic systems (Lakes Banyoles and Estanya), a number of small sinkholes have been filled with sediments and have reached the “polje” stage. Dry sinkholes with steep walls occur in several karst areas in the Iberian Range.

A different category for lake basin origin are the tufa – dammed lakes like Taravilla, and barrage lake systems in the Guadiana River (Tablas de Daimiel

National Park (Pedley et al., 1996), and in the Piedra River (Arenas et al., 2013). They have steep margins associated with the barrage, with gradual margins closer to the river inflow and flat bottoms. They are hydrologically open and connected to the fluvial drainage. The regional hydrology and the topographic location in the landscape control the formation of tufa-barrage lakes (above the regional springline) or sinkhole lakes (below the regional springline).

Lake morphology. Once karst lakes have reached a mature stage, the morphology and bathymetry of the lake is only altered by sedimentary infilling and lake level changes. Lake morphology determines the surface extent of depositional environments, - particularly littoral versus distal – and the maximum depth. Three main categories occur:

- a) Elongated, relatively steep margins and flat bottoms in tufa – barrage lakes (Lake Taravilla)
- b) Circularly- shaped, sinkhole, funnel-shaped morphologies. Lakes develop in single sinkholes with steep margins and very restricted littoral areas (Lakes El Tejo, La Cruz, La Parra, Montcortès).
- c) Sinkholes with a relatively large littoral realm (Lakes Arreo, Tobar, Enol, Marboré, Basa de la Mora); or double sinkholes (Lake Estanya) or more complex sinkhole coalescence patterns (Lakes Zoñar and Banyoles).

Changes in basin morphology occur during the lake history. For example, currently, Lake Montcortès does not have a large platform bench; however, during the mid-Holocene, carbonate littoral facies developed most likely as a result of lake level changes induced by hydrologic fluctuations that altered the shape of the littoral and facilitated carbonate formation.

In all lakes, carbonates are more abundant in the littoral and transitional areas than the deepest areas. Carbonate particle size decreases from shore to distal areas, as a reflection of distance to the main sources of carbonate: detrital component from the watershed and endogenic component from the littoral zone. The organic matter content decreases from the platform towards the slope of the depression, but increases again in the distal areas. The main organic components in the sediments are different in proximal and distal environments: a mixture of terrestrial, submerged macrophytes and algal material in the littoral zone, and a higher contribution of algal sources in the distal facies. In all cases, distal organic matter has lower TOC/TN ratios suggestive of a higher contribution of algal sources (Meyers, 1997).

Hydrologic balance and surface hydrology. The hydrological balance and lake level fluctuations are main factors controlling facies distribution and composition. Lake water input occurs through rainfall, surface drainage, surface runoff and groundwaters; outputs are through evaporation, surface drainage and groundwater fluxes. Changes in lake level occur more rapidly in hydrologically – closed lakes as a response to moisture fluctuations (Hardie et al., 1978). The presence of an active inlet determines the intensity of the clastic input to the lakes (Renaut and Gierlowski-Kordesch, 2010). Although conceptual hydrogeological models and some water balances indicate a large groundwater contribution for all lakes (except Lake Taravilla), few of the studied lakes (Lakes Banyoles and Zoñar) have a quantified water balance. In spite that surface hydrology is not a sure indication of the global water balance, open or closed surface drainage is a key characteristic of the lake basin, defining the water and sediment input to the lake. The variety of

surface drainage in Iberian karstic lakes includes: i) no inlets, only seasonal surface run-off and no surface outlets (Lakes Tejo and La Cruz) ii), only ephemeral inlets and no surface outlets (Parra, Estanya, Zoñar), iii) seasonal outlets and inlets (Lakes Tobar, Basa and Arreo), iv) permanent outlets (Lakes Montcortès, Enol, and Marboré), v) permanent inlets and outlets (Lakes Banyoles and Taravilla). We grouped them into the following types (Fig. 16C and 17):

1. Open (through – flowing) lakes, with a permanent surface outlet. It includes lakes with tufa dam like Lake Banyoles and Taravilla, and lakes without tufa dams, developed in glaciated terrains (Lakes Enol and Marboré) and non-glaciated terrains (Lakes Montcortès and El Tobar)
2. Intermediate, with an ephemeral or seasonal outlet: Examples are Lake Arreo and Basa de la Mora.
3. Closed lakes, with no surface outlet: Lakes El Tejo, La Cruz, La Parra, Estanya and Zoñar.

Watershed. The geology of the watershed influences the clastic supply to the lake and the hydrochemistry of the waters. Lakes developed in terrains with more evaporite formations (Lakes Arreo, El Tobar and Estanya) have both, more saline waters and more abundant gypsum formation. The size of the catchment also exerts a large influence in the sediment delivered to the lake: lakes with larger catchments (Lakes Banyoles, Tobar, Zoñar and Taravilla) have higher carbonate and siliciclastic contribution and higher sedimentation rates during the last millennia. Figure 17 illustrates a lake classification based on surface hydrology and sediment supply indicated by the watershed surface area.

Changes in main watershed characteristics strongly control lake sedimentation. Lakes in glaciated areas during the Late Quaternary (Lakes Marboré, Enol and Basa de la Mora) show clear depositional changes associated to the dynamics of the glaciers in the watershed and changes in the vegetation cover, both controlled by climate and hydrologic changes during the Pleistocene and Holocene (Moreno et al., 2010). Deforestation and increased population in the watersheds during Medieval times resulted in higher sedimentation rates and increased frequency of turbidity processes in most lakes. A flood frequency increase during the Little Ice Age is detected in Lake Taravilla, and it has been associated with higher rainfall (Moreno et al., 2008). The impact in sedimentation rate caused by decreasing human pressure in rural Spain after the 1950s has also been recorded in several lakes: Lake Zoñar; (Martín-Puertas et al., 2008); Lake Arreo (Corella et al., 2011a) and Lake Montcortès (Corella, 2011b).).

Climate. Iberian karst lakes are located in regions with Mediterranean, continental and high altitude climates. Precipitation ultimately controls the lake hydrology so its influence is paramount in lake dynamics. Aquifers and groundwater dynamics may introduce a time lag in the response of the lake system although observations during the last decades demonstrate a rapid transfer to rainfall signal to lake dynamics: Lake Zoñar (Valero-Garcés et al., 2006), Cañada del Hoyo lakes (López-Blanco et al., 2011), and Lake Banyoles (Brusi et al., 1990).

At millennial time-scales, the long-term trends reflect the large changes associated to glacial/interglacial dynamics. Deglaciation onset is commonly marked by an increase in organic and carbonate productivity in the lakes (Lakes Enol and Estanya, Figs. 11, 13). Some abrupt climate changes within these glacial periods

are clearly identified as changes in texture and lamination of littoral facies in Lake Banyoles (Valero-Garcés et al., 1998; Höbig et al., 2012) and as an increase in carbonate content in the Enol sequence (Moreno et al., 2010). Carbonate and gypsum content also mark the rapid hydrological response of Lake Estanya during deglaciation (Fig. 11; (Morellón et al., 2009b)). Changes in the drainage and provenance area caused by glacier retreat could be responsible of some of the observed changes as documented in the Alps (L'Annecy (Nicoud and Manalt, 2001)) and the Pyrenees (Lake Tramacastilla, (García-Ruiz and Valero Garcés, 1998)).

Large Holocene moisture changes in the Iberian Peninsula have also been reflected in facies variability in the lake sequences. For example, Lakes Basa de la Mora (Fig. 7) and Estanya (Fig. 11) show more profundal facies during the relatively more humid early Holocene and more littoral and carbonate-rich littoral facies during the more arid mid-Holocene. Another clear example of climatically-driven evolution is the coherent trends identified in Iberian lakes during the more arid Medieval Climate Anomaly and the more humid Little Ice Age (Valero Garcés et al., 2009; Morellón et al., 2012; Moreno et al., 2012).

Climatic factors cannot be disregarded on the generation processes of clastic facies as an increase in storminess is directly related to higher run-off. Increasing aridity during some intervals (e.g., Medieval Climate Anomaly) may also play an important role in the sediment availability as dry soils are more easily eroded. In Lake Montcortès, the more abundant presence of clastic laminae in the varved intervals is linked to more precipitation events per year (Corella et al 2012). Temperature has a strong influence in biological activity, and in the timing of

carbonate precipitation related to planktonic blooms. In both, Lake Zoñar and Lake Montcortès changes in water temperature have resulted in different textures of the calcite precipitated in the lakes during the Little Ice Age and the Iberian – Roman Humid Period (Martín-Puertas et al., 2008; Corella et al., 2012). Recent endogenic calcite formation (calcite lamina thickness) is correlated with winter rainfall in Lake La Cruz (Romero-Viana et al., 2008) and a similar mechanism could be in place in other Spanish lakes. Temperature also influences the solubility of carbonate (Deocampo, 2010) and may be the main reason for the low carbonate content of proglacial Enol and recent Marboré sedimentary sequences (Moreno et al., 2011). The preservation potential of carbonates in cold-water lakes could be diminished by preferential dissolution of fine-grained calcite (glacier flour) in these settings.

Water chemistry. Hydrological balance and watershed geology greatly determine water chemical composition. In our data set, lakes with functional outlets have lower TDS and conductivities and lakes with no surface outlets have the highest salinities (Fig. 16C, D). Sulfate is the most common anion in lakes developed over a dominantly evaporite substrate (Lakes Arreo and El Tobar) while carbonate- bicarbonate dominates in carbonate terrains. Ca^{2+} is the dominant cation in evaporite-rich watersheds while the dominance of Ca^{2+} or Mg^{2+} in carbonate-terrain lakes depends on the relative abundance of dolomite. El Tobar is a unique case because of the presence a hypersaline monimolimnion related to the input of saline springs (Vicente et al., 1993).

In hydrologically – open basins with short residence times as in Lakes Taravilla and Banyoles, chemical compositions of lake waters and surface waters is similar and do not greatly change through time. If lake level drops below the

outlet, hydrologically – open lakes may become closed, evaporative basins, as it is the case in Lake Zoñar during several dry periods in the past (Martín-Puertas et al., 2008b) or Lake Taravilla in recent droughts in the 1990s (Valero-Garcés et al., 2008). Increased evaporation changes water salinity as reflected in deposition of other carbonate phases (aragonite, HMC) and sulfates (gypsum) (Fig. 9). Lakes depending almost exclusively on groundwater fluxes as those in Cañada del Hoyo react even more rapidly to water balances with changes in littoral and distal facies distribution (Romero-Viana et al., 2009; López-Blanco et al., 2011); however, changes in salinity are not large enough to precipitate more saline minerals. Saline water input plays a definitive role in Lakes El Tobar and Arreo water chemistry and mineral deposition because the presence of saline groundwaters.

Water chemistry is also key to carbonate preservation since hypolimnetic conditions may dissolve small calcite grains before deposition (Dean et al., 2007). Even lakes with high littoral and epilimnetic carbonate productivity and well-developed carbonate benches may have little accumulation of carbonates in the distal, deeper zones dominated by organic deposition (Gilbert and Lesak, 1981). Dissolution of carbonates in hypolimnetic waters is favored by lower temperatures and lower alkalinity and pH caused by organic matter decay (Deocampo, 2010). These processes could also contribute to the absence of some calcite layers in varve facies (Montcortès, Zoñar).

Dissolution also affects clastic carbonates. Rhythmites in Lakes Enol, Marboré and Basa de la Mora have very low carbonate content, in spite of the dominant carbonate composition of the rock formations and the main moraines in the catchment. Low temperatures during glacial times (Lake Enol) would have

1138 been conducive to dissolution of fine clastic carbonates delivered into the lake by
1139 the proglacial streams. In Lakes Marboré and Basa de la Mora, the Late glacial
1140 and Holocene moraines in the catchment are carbonate - rich (up to 11 % TIC)
1141 while the laminated sediments are carbonate-free (< 0.1 % TIC). Cold, low salinity
1142 surface waters in these high altitude settings (> 2500 m asl) could have dissolved
1143 some of the fine carbonate particles before entering the lake. Groundwater seeping
1144 into the lake could also contribute to further dissolve fine-grained carbonates.

1145 Water stratification. Lake depth is the main parameter controlling thermal
1146 stratification and development of anoxic conditions (Wetzel, 2001) The
1147 development of seasonal or permanent thermal stratification requires a minimum
1148 water depth of 6 m (Shaw et al., 2002) and so thermal stratification occurs in all
1149 Iberian karst lakes, except Lake La Basa de la Mora (4 m max. water depth) (Fig.
1150 16B, D). Lake Montcortès is the only lake with permanent anoxic waters due to
1151 thermal stratification. Recent monitoring and paleolimnological reconstructions
1152 based on short cores point to dominant meromictic conditions in Lake Arreo prior to
1153 the 1960s (Corella et al., 2011a), disturbed by water removal for irrigation. In Lake
1154 El Tobar the input of highly saline waters is responsible for hypersaline
1155 hypolimnion and permanent meromixis (Vicente et al., 1993).

1156 The sedimentary sequences of the studied Spanish lakes show examples
1157 where changes from predominantly stratified to mixed water column conditions in
1158 distal areas are reflected in the alternation of banded/ massive and laminated
1159 facies (Lakes Arreo, Estanya and Zoñar).

1160 Lake evolution. Iberian karstic lakes are small compared with large karstic
1161 lacustrine basins, but their facies associations and sequences serve as analogs for

both Quaternary and pre-Quaternary lake basins. Our review shows a large spatial facies and depositional environment variability: in a < 100 m long transect we find littoral facies with evidences of subaerial exposure and profundal finely facies (Lakes Montcortés and Zoñar). Similar large variability occurs if time is considered. Only in Lake Montcortés, varved facies have been deposited during at least the last 5000 years in the deepest part of the lake basin. Most profundal areas have experienced rather abrupt changes in depositional environment conditions (depth, oxygen content) at a decadal or centennial scale.

Hydrology is clearly the paramount control on facies and depositional environment patterns distribution and evolution. The lake's hydrological budget is greatly governed by the basin morphology and the water balance but at short time scales as those illustrated in this review (millennial scale), factors as tectonics, subsidence and karstic activity intensity are secondary to climate. Most Iberian karstic lake facies sequences indicative of lake level changes are reflecting the local and regional climate variability during the last millennia.

In large lakes, allogenic factors creating accommodation space (tectonics, karstic activity, volcanism, fluvial damming) and controlling water budgets (climate) determine sedimentation patterns (Bohacs et al., 2000). As we have shown in previous sections, the smaller Iberian karstic lakes, show similar interactions of the four main variables: sediment supply, water supply, basin-sill height (spill point), and basin floor depth. Our classification of Iberian Karstic lake basins based on surface hydrology and watershed surface area (Fig. 17) respond to the two main criteria in Bohacs et al (2000) classification of lake sequence stratigraphic patterns: overfilled, balanced-fill and underfilled. Overfilled basins are characterized by water

and sediment supply higher than accommodation rates, stable lake level, open hydrology and dilute waters. Lakes with open surface hydrology as Banyoles, Montcortès and Taravilla could be included in this type. Examples of underfilled basins, with smaller water and sediment supply than accommodation rates and closed surface hydrology are Lakes Zoñar and Estanya. Aggradations pattern are more common in Iberian lake sequences, and only in those cases with stronger fluvial influence (Lakes Taravilla and Banyoles), some progradational patterns occur in littoral sequences.

Our review illustrates how lake types evolve through the existence of a lake basin at centennial or even smaller time scales. In the absence of strong tectonic control, accommodation space and basin morphology are determined by surface (karstic and hydrologic) processes. Karstic processes generating new accommodation space are only active in Lake Banyoles; in most Iberian lake sequences, hydrology has determined changes in the surface drainage evolution. Lacustrine sedimentation may start much later than the basin was formed by karstic processes. This is clearly documented in Lake La Parra by the occurrence of an alteration zone between the substrate and the lake sediments. Profundal facies occurred early in the lake evolution (Lakes La Parra and Estanya), as an indication of the rapid flooding of the basin. At a Holocene- scale, the lake basins have not deepened much due to karstification processes, so the sedimentation pattern is comparable to volcanic crater lakes or dammed-lakes, starting with deep facies and gradually filled with sediment. Human activities have been a significant contributor to sediment input during the last millennium.

The variability of lake facies, depositional patterns and evolutionary trends envisaged in the geologic record of lake basins (Gierlowski-Kordesch and Kelts, 2000) is even more evident in extant and Quaternary lakes. Facies analyses from modern lake systems help to refine facies models for larger lake basins and to understand their dynamics at different temporal and spatial scales.

6. Conclusions

Iberian karst lakes are small lake systems with a large variability of facies and depositional environments. Carbonates are the main sediment components because of the geology of the watershed and the lake water hydrochemistry. Shallow littoral depositional environments with massive and banded facies are dominant because of the relatively small size of the lakes. Distal profundal facies are more spatially restricted but show a large variability of laminated facies. Slump and gravitational processes are more common in the lake basins with higher depth/surface ratio. Endogenic carbonate formation occurs in diverse settings and, although dominated by littoral processes, also includes epilimnetic calcite precipitation. Except in glaciated terrains, detrital carbonate input is a major contribution to the carbonate budget in these lakes. In spite of the complexity of the facies evolution in the Iberian karst lakes, the sedimentary sequences provide examples of evolutionary trends related to paleohydrological variability and also to changes in the watershed. Hydrology is the main control on facies and depositional environment patterns distribution and evolution. The large changes associated with glacial/interglacial cycles and the inherent large humidity variability of the Holocene Mediterranean climates have exerted a great control on lake deposition and

climate has been an important forcing behind the hydrological/ depositional response of the lakes.

Human impact in the watersheds and the lakes has been documented since Iberian-Roman times and it has increased since the Medieval Ages with a much higher sediment delivery to the lakes. Several examples of the impact of historical land-use changes as the Medieval deforestation, the periods of higher population in the Pyrenees during the 19th century, or the abandonment of the rural areas in the mid-20th century are documented in several lake sequences.

Detailed sedimentary facies of these systems provide the framework to interpret past environmental changes and contribute to reconstruction of past hydrological and climate variability.

Acknowledgements

The research on Iberian Karst systems has been supported by LIMNOCAL (CGL2006-13327-C04-01), Consolider GRACCIE (CSD2007-00067) and GLOBALKARST CGL2009-08415 funded by the Spanish Ministry of Economy and Competitiveness. We also wish to thank Tim Horscroft and Brian Jones for inviting us to submit a review article on this topic to Sedimentary Geology. We are grateful to the in-depth reviews by Elizabeth Gierlowski-Kordesch and Bill Last.

References

- 1256 Alagöz, C.A., 1967. Sivas Çevresi ve Doğusunda Jips Karstı Olayları - Les
1257 phénomènes karstiques du gypse aux environs et à l'est de Sivas. Ankara
1258 Üniversitesi Dil ve Tarih-Coğrafya Fakültesi Yayını, Ankara, 126 pp.
- 1259 Alonso-Zarza, A.M., Dorado-Valiño, M., Valdeolmillos-Rodríguez, A., Ruiz-Zapata,
1260 M.B., 2006. A recent analogue for palustrine carbonate environments: The
1261 Quaternary deposits of Las Tablas de Daimiel wetlands, Ciudad Real,
1262 Spain. SPECIAL PAPERS-GEOLOGICAL SOCIETY OF AMERICA, 416:
1263 153.
- 1264 Alonso-Zarza, A.M., Wright, V.P., 2010. Palustrine Carbonates. In: A.M. Alonso-
1265 Zarza, L.H. Tanner (Eds.), Carbonates in Continental Settings: Processes,
1266 Facies, and Application. Developments in Sedimentology. Elsevier,
1267 Amsterdam (The Netherlands), pp. 103-131.
- 1268 Alonso, M., 1998. Las lagunas de la España peninsular. Limnetica, 15: 1-176.
- 1269 Anselmetti, F.S., Ariztegui, D., Hodell, D.A., Hillesheim, M.B., Brenner, M., Gilli, A.,
1270 McKenzie, J.A., Mueller, A.D., 2006. Late Quaternary climate-induced lake
1271 level variations in Lake Peten Itza, Guatemala, inferred from seismic
1272 stratigraphic analysis. Palaeogeography, Palaeoclimatology, Palaeoecology,
1273 230(1-2): 52-69.
- 1274 Arenas, C., Vázquez-Urbez, M., Auqué, L., Sancho, C., Osácar, C., Pardo, G.,
1275 2013. Intrinsic and extrinsic controls of spatial and temporal variations in
1276 modern fluvial tufa sedimentation: A thirteen-year record from a semi-arid
1277 environment. Sedimentology: n/a-n/a.
- 1278 Ariztegui, D., Anselmetti, F.S., Robbiani, J.M., Bernasconi, S.M., Brati, E., Gilli, A.,
1279 Lehmann, M.F., 2010. Natural and human-induced environmental change in

1280 southern Albania for the last 300 years -- Constraints from the Lake Butrint
 1281 sedimentary record. *Global and Planetary Change*, 71(3-4): 183-192.

1282 Barreiro-Lostres, F., Moreno, A., Giralt, S., Valero-Garcés, B.L., 2013. Evolución
 1283 sedimentaria del lago kárstico de La Parra (Cuenca) durante los últimos
 1284 1600 años: paleohidrología, clima e impacto humano. *Cuadernos de*
 1285 *Investigación Geográfica*, 39(1): 179-193.

1286 Barreiro-Lostres, F., Moreno, A., Valero-Garcés, B., 2011. Depositional facies of La
 1287 Parra karstic lake (Cuenca) during the last 1600 cal. years BP. *Geogaceta*,
 1288 50(2): 109-112.

1289 Bischoff, J.L., Julià, R., Shanks, W.C., Rosenbauer, R.J., 1994. Karstification
 1290 without carbonic acid: Bedrock dissolution by gypsum-driven
 1291 dedolomitization. *Geology*, 22(11): 995-998.

1292 Bohacs, K.M., Carroll, A.R., Neal, J.E., Mankiewicz, P.J., 2000. Lake-basin type,
 1293 source potential, and hydrocarbon character: an integrated sequence-
 1294 stratigraphic-geochemical framework. In: E.H. Gierlowski-Kordesch, K. Kelts
 1295 (Eds.), *Lake basins through space and time*. AAPG Studies in Geology, pp.
 1296 3-34.

1297 Brauer, A., 2004. Annually laminated lake sediments y their palaeoclimatic
 1298 relevance. In: H. Fischer, e. al. (Eds.), *The climate in historical times*
 1299 *towards a synthesis of Holocene proxy data and climate Models*. Springer,
 1300 Berlin, pp. 109-128.

1301 Brauer, A., Casanova, J., 2001. Chronology and depositional processes of the
 1302 laminated sediment record from Lac d'Annecy, French Alps. *Journal of*
 1303 *Paleolimnology*, 25(2): 163-177.

- 1304 Brauer, A., Haug, G.H., Dulski, P., Sigman, D.M., Negendank, J.F.W., 2008. An
1305 abrupt wind shift in western Europe at the onset of the Younger Dryas cold
1306 period. *Nature Geosciences*, 1(8): 520-523.
- 1307 Brusi, D., Bach, J., Sanz, M., 1990. Itinerari geològic de Banyoles. Descoberta del
1308 funcionament del sistema lacustre. Apunts 22. Eumo Editorial, Barcelona,
1309 124 pp.
- 1310 Camps, J., Gonzalvo, I., Güell, J., López, P., Tejero, A., Toldrà, X., Vallespinos, F.,
1311 Vicens, M., 1976. El lago de Montcortès, descripció de un ciclo anual.
1312 *Oecologia aquatica*, 2: 99-110.
- 1313 Canals, M., Got, H., Julia, R., Serra, J., 1990. Solution-collapse depressions and
1314 suspensates in the limnogenic lake of Banyoles (NE Spain). *Earth Surface*
1315 *Processes and Landforms*, 15(3): 243-254.
- 1316 Capel Molina, J.J., 1981. Los climas de España. *Ciencias geográficas*, 8. Oikos-
1317 Tau, 429 pp.
- 1318 Catalán, J., Camarero, L., Felip, M., Pla, S., Ventura, M., Buchaca, T., Bartumeus,
1319 F., de Mendoza, G., Miró, A., Casamayor, E.O., Medina-Sánchez, J.M.,
1320 Bacardit, M., Altuna, M., Bartrons, M., Díaz de Quijano, D., 2006. High
1321 mountain lakes: extreme habitats and witnesses of environmental changes.
1322 *Limnetica*, 25((1-2)): 551-584.
- 1323 Cohen, A.S., 2003. *Paleolimnology: The history and evolution of lake systems*.
1324 Oxford University Press, New York, 500 pp.
- 1325 Colomer, J., Serra, T., Soler, M., Casamitjana, X., 2002. Sediment fluidization
1326 events in a lake caused by large monthly rainfalls. *Geophys. Res. Lett.*,
1327 29(8): 1260.

1328 Corella, J., Amrani, A., Sigró, J., Morellón, M., Rico, E., Valero-Garcés, B., 2011a.
1329 Recent evolution of Lake Arreo, northern Spain: influences of land use
1330 change and climate. *Journal of Paleolimnology*, 46(3): 469-485.

1331 Corella, J.P., Valero-Garcés, B., Moreno, A., Morellón, M., Rull, V., Giralt, S., Rico,
1332 M.T., Pérez-Sanz, A., 2011b. Climate and human impact on a meromictic
1333 lake during the last 6000 years (Montcortès Lake, Central Pyrenees, Spain)
1334 *Journal of Paleolimnology*, 46(3): 351-367.

1335 Corella, J.P., Brauer, A., Mangili, C., Rull, V., Vegas-Vilarrúbia, T., Morellón, M.,
1336 Valero-Garcés, B.L., 2012. The 1.5-ka varved record of Lake Montcortès
1337 (southern Pyrenees, NE Spain). *Quaternary Research*, 78(2): 323-332.

1338 Corella, J.P., Stefanova, V., El Anjoumi, A., Rico, E., Giralt, S., Moreno, A., Plata-
1339 Montero, A., Valero-Garcés, B.L., 2013a. A 2500-year multi-proxy
1340 reconstruction of climate change and human activities in northern Spain:
1341 The Lake Arreo record. *Palaeogeography, Palaeoclimatology,*
1342 *Palaeoecology*, 386(0): 555-568.

1343 Corella, J.P., Arantegui, A., Loizeau, J.L., DelSontro, T., le Dantec, N., Stark, N.,
1344 Anselmetti, F.S., Girardclos, S., 2013b. Sediment dynamics in the
1345 subaquatic channel of the Rhone delta (Lake Geneva, France/Switzerland).
1346 *Aquatic Sciences*, in press. DOI: 10.1007/s00027-013-0309-4

1347 Cvijic, J., 1981. The dolines: translation of geography. In: M.M. Sweeting (Ed.),
1348 *Karst geomorphology*. Hutchinson, Pennsylvania, pp. 225-276.

1349 Chapron, E., Van Rensbergen, P., De Batist, M., Beck, C., Henriot, J.P., 2004.
1350 Fluid-escape features as a precursor of a large sublacustrine sediment slide
1351 in Lake Le Bourget, NW Alps, France. *Terra Nova*, 16(5): 305-311.

- 1352 D'Amore, G., 1983. Il problema della conservazione e della utilizzazione del lago di
1353 Pergusa. Atti del Convegno: La protezione dei laghi e delle zone umide in
1354 Italia, XXXVIII** Memorie Soc. Geogr. Italiana.
- 1355 Dasi, M.J., Miracle, M.R., 1991. Distribución vertical y variación estacional del
1356 fitoplancton de una laguna Karstica meromictica, La Laguna de la Cruz
1357 (Cuenca, España). Limnetica, 7: 37-59.
- 1358 Davaud, E., Girardclos, S., 2001. Recent Freshwater Ooids and Oncoids from
1359 Western Lake Geneva (Switzerland): Indications of a Common Organically
1360 Mediated Origin. Journal of Sedimentary Research, 71(3): 423-429.
- 1361 Dean, W.E., 1981. Carbonate minerals and organic matter in sediments of modern
1362 north-temperate hard-water lakes. In: E.G. Ethridge, R.M. Flores (Eds.),
1363 Recent and ancient nonmarine depositional environments: models for
1364 exploration. SEPM Special Publication, pp. 213-231.
- 1365 Dean, W.E., Forester, R.M., Bright, J., Anderson, R.Y., 2007. Influence of the
1366 diversion of Bear River into Bear Lake (Utah and Idaho) on the environment
1367 of deposition of carbonate minerals. Limnology and Oceanography, 52(3):
1368 1094-1111.
- 1369 Dean, W.E., Fouch, T.D., 1983. Lacustrine environment. In: R.A. Schole, D.G.
1370 Bebout, C.H. Moore (Eds.), Carbonate depositional environments. AAPG
1371 Memoir pp. 96-130.
- 1372 Dean, W.E., Gorham, E., 1998. Magnitude and significance of carbon burial in
1373 lakes, reservoirs, and peatlands. Geology, 26(6): 535-538.

- 1374 Deocampo, D.M., 2010. Chapter 1 The Geochemistry of Continental Carbonates.
 1375 In: A.M. Alonso-Zarza, L.H. Tanner (Eds.), *Developments in Sedimentology*.
 1376 Elsevier, Amsterdam (The Netherlands), pp. 1-59.
- 1377 Eugster, H.P., Kelts, K.R., 1983. Lacustrine chemical sediments. In: A.S. Goudie,
 1378 K. Pye (Eds.), *Chemical sediments and Geomorphology*. Academic Press,
 1379 London, pp. 321-368.
- 1380 García-Ruiz, J.M., Valero Garcés, B.L., 1998. Historical Geomorphic Processes
 1381 and Human Activities in the Central Spanish Pyrenees. *Mountain Research*
 1382 *and Development*, 18(4): 309-320.
- 1383 Garrote Ruíz, A., Muñoz Jiménez, J.I.E.V., 2001. Mapa geológico del País Vasco
 1384 Escala 1:25.000. In: d.I.E. (EVE) (Ed.), Bilbao (Spain).
- 1385 Gibbons, W., Moreno, T., 2002. *The Geology of Spain*. The Geological Society of
 1386 London, 649 pp.
- 1387 Gierlowski-Kordesch, E.H., 2010. Lacustrine carbonates. In: A.M. Alonso-Zarza,
 1388 L.H. Tanner (Eds.), *Carbonates in Continental Settings: Processes, Facies,*
 1389 *and Application*. *Developments in Sedimentology*. Elsevier, Amsterdam, pp.
 1390 1-101.
- 1391 Gierlowski-Kordesch, E.H., Kelts, K.R., 1994. Global geological record of lake
 1392 basins. *World and regional geology series 4, 1*. Cambridge University
 1393 Press, Cambridge (United Kingdom), 427 pp.
- 1394 Gierlowski-Kordesch, E.H., Kelts, K.R., 2000. *Lake basins through space and time*.
 1395 *Studies in Geology 46*. The American Association of Petroleum Geologists,
 1396 Tulsa, OK, 648 pp.

1397 Gilbert, R., Lesak, S.M., 1981. Factors affecting marl deposition in Knowlton Lake,
1398 southeastern Ontario. *Journal of Great Lakes Research*, 7: 286-289.

1399 Girardclos, S., Schmidt, O.T., Sturm, M., Ariztegui, D., Pugin, A., Anselmetti, F.S.,
1400 2007. The 1996 AD delta collapse and large turbidite in Lake Brienz. *Marine*
1401 *Geology*, 241(1-4): 137-154.

1402 Glenn, C.R., Kelts, K., 1991. Sedimentary rhythms in lake deposits. In: G. Einsele,
1403 W. Ricken, A. Seilacher (Eds.), *Cycles and Events in Stratigraphy*. Springer,
1404 Berlín (Germany), pp. 188–221.

1405 Gutiérrez-Elorza, M., 2001. *Geomorfología climática*. Editorial Omega, Barcelona,
1406 642 pp.

1407 Gutierrez-Elorza , M., Peña Monne, J.L., 1979. El karst de Villar del Cobo (Sierra
1408 de Albarracín). *Estudios Geológicos IGME*, 35: 651-654.

1409 Gutiérrez, F., Calaforra, J., Cardona, F., Ortí, F., Durán, J., Garay, P., 2008.
1410 Geological and environmental implications of the evaporite karst in Spain.
1411 *Environmental Geology*, 53(5): 951-965.

1412 Gutiérrez, F., Valero-Garcés, B., Desir, G., González-Sampériz, P., Gutiérrez, M.,
1413 Linares, R., Zarroca, M., Moreno, A., Guerrero, J., Roqué, C., Arnold, L.J.,
1414 Demuro, M., 2013. Late Holocene evolution of playa lakes in the central
1415 Ebro depression based on geophysical surveys and morpho-stratigraphic
1416 analysis of lacustrine terraces. *Geomorphology*, in press.

1417 Hampton, M.A., Lee H.J., Locat, J., 1996. Submarine landslides. *Reviews of*
1418 *Geophysics* 34: 33-59.

1419 Hardie, L.A., Smoot, J.P., Eugster, H.P., 1978. Saline lakes and their deposits: a
1420 sedimentological approach. In: A. Matter, M.E. Tucker (Eds.), *Modern and*

1421 Ancient Lake Sediments. International Association of Sedimentologists
1422 Special Publication, pp. 7-42.

1423 Höbig, N., Weber, M.E., Kehl, M., Weniger, G.-C., Julià, R., Melles, M., Fülöp, R.-
1424 H., Vogel, H., Reicherter, K., 2012. Lake Banyoles (northeastern Spain): A
1425 Last Glacial to Holocene multi-proxy study with regard to environmental
1426 variability and human occupation. *Quaternary International*, 274(0): 205-
1427 218.

1428 Hodell, D.A., Anselmetti, F.S., Ariztegui, D., Brenner, M., Curtis, J.H., Gilli, A.,
1429 Grzesik, D.A., Guilderson, T.J., Müller, A.D., Bush, M.B., Correa-Metrio, A.,
1430 Escobar, J., Kutterolf, S., 2008. An 85-ka record of climate change in
1431 lowland Central America. *Quaternary Science Reviews*, 27(11-12): 1152-
1432 1165.

1433 Hsü, K.J., Kelts, K., 1985. Swiss lakes as a geological laboratory.
1434 *Naturwissenschaften*, 72(6): 315-321.

1435 Imboden, D.M., 2007. The Motion of Lake Waters, *The Lakes Handbook*, Volume
1436 1. Blackwell Science Ltd, pp. 115-152.

1437 Jones, M.D., Roberts, C.N., Leng, M.J., Turkes, M., 2006. A high-resolution late
1438 Holocene lake isotope record from Turkey and links to North Atlantic and
1439 monsoon climate. *Geology*, 34(5): 361-364.

1440 Julià, R., 1980. La conca lacustre de Banyoles *Centro d'Estudios Comarcals de*
1441 *Banyoles*, Besalu, 188 pp.

1442 Julià, R., Burjachs, F., Dasí, M.J., Mezquita, F., Miracle, M.R., Roca, J.R., Seret,
1443 G., Vicente, E., 1998. Meromixis origin and recent trophic evolution in the

1444 Spanish mountain lake La Cruz. *Aquatic Sciences - Research Across*
 1445 *Boundaries*, 60(4): 279-299.
 1446 Kelts, K., 1978. Geological and sedimentary evolution of Lakes Zurich and Zug,
 1447 ETH-Zürich, 250 pp.
 1448 Kelts, K., Hsü, K.J., 1978. Freshwater carbonate sedimentation. In: A. Lerman
 1449 (Ed.), *Lakes: chemistry, geology and physics*. Springer-Verlag, New York,
 1450 pp. 295-323.
 1451 Kindinger, J.L., Davis, J.B., Flocks, J.G., 1999. Geology and evolution of lakes in
 1452 north-central Florida. *Environmental Geology*, 38(4): 301-321.
 1453 Koschel, R., 1997. Structure and function of pelagic calcite precipitation in lake
 1454 ecosystems. *Verhandlungen der Internationalen Vereinigung Limnologie*,
 1455 26: 343-349.
 1456 Last, F.M., Last, W.M., 2012. Lacustrine carbonates of the northern Great Plains of
 1457 Canada. *Sedimentary Geology*, 277–278(0): 1-31.
 1458 Last, W.M., 1990. Paleochemistry and paleohydrology of Ceylon Lake, a salt-
 1459 dominated playa basin in the northern Great Plains, Canada. *Journal of*
 1460 *Paleolimnology*, 4(3): 219-238.
 1461 Last, W.M., 1994. Deep-water evaporite mineral formation in lakes of western
 1462 Canada. In: R.W. Renault, Last W.M. (Ed.), *Sedimentology and*
 1463 *Geochemistry of Modern and Ancient saline lakes*. Society for Sedimentary
 1464 Geology, Tulsa, pp. 51-59.
 1465 Last, W.M., Smol, J.P., 2001. *Tracking Environmental Change using Lake*
 1466 *Sediments*. *Developments in Paleoenvironmental Research Series*. Kluwer
 1467 Academic Publishers, Norwell, MA, U.S.A.

- 1468 Leonard, E.M., Reasoner, M.A., 1999. A Continuous Holocene Glacial Record
1469 Inferred from Proglacial Lake Sediments in Banff National Park, Alberta,
1470 Canada. *Quaternary Research*, 51(1): 1-13.
- 1471 Lézine, A.M., von Grafenstein, U., Andersen, N., Belmecheri, S., Bordon, A.,
1472 Caron, B., Cazet, J.P., Erlenkeuser, H., Fouache, E., Grenier, C.,
1473 Huntsman-Mapila, P., Hureau-Mazaudier, D., Manelli, D., Mazaud, A.,
1474 Robert, C., Sulpizio, R., Tiercelin, J.J., Zanchetta, G., Zeqollari, Z., 2010.
1475 Lake Ohrid, Albania, provides an exceptional multi-proxy record of
1476 environmental changes during the last glacial-interglacial cycle.
1477 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 287(1-4): 116-127.
- 1478 Lindhorst, K., Vogel, H., Krastel, S., Wagner, B., Hilgers, A., Zander, A., Schwenk,
1479 T., Wessels, M., Daut, G., 2010. Stratigraphic analysis of lake level
1480 fluctuations in Lake Ohrid: an integration of high resolution hydro-acoustic
1481 data and sediment cores. *Biogeosciences*, 7(11): 3531-3548.
- 1482 Locat, J., Lee, H.J., 2002. Submarine landslides: advances and challenges.
1483 *Canadian Geotechnical Journal*, 39(1): 193-212.
- 1484 López-Blanco, C., Gaillard, M.-J., Miracle, M.R., Vicente, E., 2011. Lake-level
1485 changes and fire history at Lagunillo del Tejo (Spain) during the last
1486 millennium: Climate or human impact? *The Holocene*.
- 1487 López-Vicente, M., Navas, A., Machín, J., 2009. Geomorphic mapping in endorheic
1488 catchments in the Spanish Pyrenees: An integrated GIS analysis of karstic
1489 features. *Geomorphology*, 111(1-2): 38-47.
- 1490 Lotter, A.F., Lemcke, G., 1999. Methods for preparing and counting biochemical
1491 varves. *Boreas*, 28(2): 243-252.

- 1492 Magny, M., de Beaulieu, J.-L., Drescher-Schneider, R., Vannière, B., Walter-
1493 Simonnet, A.-V., Millet, L., Bossuet, G., Peyron, O., 2006. Climatic
1494 oscillations in central Italy during the Last Glacial-Holocene transition: the
1495 record from Lake Accesa. *Journal of Quaternary Science*, 21(4): 311-320.
- 1496 Magny, M., de Beaulieu, J.-L., Drescher-Schneider, R., Vanniere, B., Walter-
1497 Simonnet, A.-V., Miras, Y., Millet, L., Bossuet, G., Peyron, O., Brugiapaglia,
1498 E., Leroux, A., 2007. Holocene climate changes in the central
1499 Mediterranean as recorded by lake-level fluctuations at Lake Accesa
1500 (Tuscany, Italy). *Quaternary Science Reviews*, 26(13-14): 1736-1758.
- 1501 Mangili, C., Brauer, A., Plessen, B., Moscariello, A., 2007. Centennial-scale
1502 oscillations in oxygen and carbon isotopes of endogenic calcite from a
1503 15,500 varve year record of the Pianico interglacial. *Quaternary Science*
1504 *Reviews*, 26(13-14): 1725-1735.
- 1505 Marquínez, J., Adrados, L., 2000. La geología y el relieve de los Picos de Europa.
1506 *Naturalia Cantabricae*, 1: 3-19.
- 1507 Martín-Puertas, C., Valero-Garcés, B.L., Brauer, A., Mata, M.P., Delgado-Huertas,
1508 A., Dulski, P., 2009. The Iberian-Roman Humid Period (2600-1600 cal yr
1509 BP) in the Zoñar Lake varve record (Andalucía, southern Spain). *Quaternary*
1510 *Research*, 71(2): 108-120.
- 1511 Martín-Puertas, C., Valero-Garcés, B.L., Mata, M.P., González-Sampériz, P., Bao,
1512 R., Moreno, A., Stefanova, V., 2008. Arid and humid phases in southern
1513 Spain during the last 4000 years: the Zoñar Lake record, Cordoba. *The*
1514 *Holocene*, 18(6): 907-921.

1515 Martín-Rubio, M., Rodríguez-Lazaro, J., Anadón, P., Robles, F., Utrilla, R.,
 1516 Vázquez, A., 2005. Factors affecting the distribution of recent lacustrine
 1517 ostracoda from the Caicedo de Yuso-Arreo Lake (Western Ebro Basin,
 1518 Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 225(1-4):
 1519 118-133.

1520 Meyers, P.A., 1997. Organic geochemical proxies of paleoceanographic,
 1521 paleolimnologic and paleoclimatic processes. *Organic Geochemistry*,
 1522 97(5/6): 213-250.

1523 Miracle, M.R., Vicente, E., Pedrós-Alió, C., 1992. Biological studies of spanish
 1524 meromictic and stratified karstic lakes. *Limnetica*, 8: 59-77.

1525 Modamio, X., Pérez, V., Samarra, F., 1988. Lirnnologia del lago de Montcortés
 1526 (ciclo 1978-79) (Pallars Jussá, Lleida). *Oecologia aquatica*, 9: 9-17.

1527 Moner, J., Riera, J., Abellá, C., Garrofé, T., 1987. Banyoles, a l'ombra dels JJ.OO.
 1528 92. Edicions Periódiques de les Comarques, S.A., Girona (Spain).

1529 Morellón, M., Anselmetti, F.S., Valero-Garcés, B., Giralt, S., Ariztegui, D., Sáez, A.,
 1530 Mata, M.P., Barreiro-Lostres, F., Rico, M., Moreno, A., in review. The
 1531 influence of groundwater in lacustrine sedimentation: origin and
 1532 paleoenvironmental significance of homogenites in karstic Lake Banyoles
 1533 (NE Spain). *Sedimentology*.

1534 Morellón, M., Pérez-Sanz, A., Corella, J.P., Büntgen, U., Catalán, J., González-
 1535 Sampériz, P., González-Trueba, J.J., López-Sáez, J.A., Moreno, A., Pla-
 1536 Rabes, S., Saz-Sánchez, M.Á., Scussolini, P., Serrano, E., Steinhilber, F.,
 1537 Stefanova, V., Vegas-Vilarrúbia, T., Valero-Garcés, B., 2012. A multi-proxy

1538 perspective on millennium-long climate variability in the Southern Pyrenees.
1539 Clim. Past, 8(2): 683-700.

1540 Morellón, M., Valero-Garcés, B., Anselmetti, F., Ariztegui, D., Schnellmann, M.,
1541 Moreno, A., Mata, P., Rico, M., Corella, J.P., 2009a. Late Quaternary
1542 deposition and facies model for karstic Lake Estanya (North-eastern Spain).
1543 Sedimentology, 56(5): 1505-1534.

1544 Morellón, M., Valero-Garcés, B., González-Sampériz, P., Vegas-Vilarrúbia, T.,
1545 Rubio, E., Rieradevall, M., Delgado-Huertas, A., Mata, P., Romero, Ó.,
1546 Engstrom, D., López-Vicente, M., Navas, A., Soto, J., 2011. Climate
1547 changes and human activities recorded in the sediments of Lake Estanya
1548 (NE Spain) during the Medieval Warm Period and Little Ice Age. Journal of
1549 Paleolimnology, 46(3): 423-452.

1550 Morellón, M., Valero-Garcés, B., Moreno, A., González-Sampériz, P., Mata, P.,
1551 Romero, O., Maestro, M., Navas, A., 2008. Holocene palaeohydrology and
1552 climate variability in Northeastern Spain: The sedimentary record of Lake
1553 Estanya (Pre-Pyrenean range). Quaternary International, 181: 15-31.

1554 Morellón, M., Valero-Garcés, B., Vegas-Vilarrúbia, T., González-Sampériz, P.,
1555 Romero, Ó., Delgado-Huertas, A., Mata, P., Moreno, A., Rico, M., Corella,
1556 J.P., 2009b. Lateglacial and Holocene palaeohydrology in the western
1557 Mediterranean region: The Lake Estanya record (NE Spain). Quaternary
1558 Science Reviews, 28(25–26): 2582-2599.

1559 Moreno, A., Cacho, I., Canals, M., Prins, M.A., Sanchez-Goni, M.-F., Grimalt, J.O.,
1560 Weltje, G.J., 2002. Saharan Dust Transport and High-Latitude Glacial

1561 Climatic Variability: The Alboran Sea Record. *Quaternary Research*, 58(3):
1562 318-328.

1563 Moreno, A., López-Merino, L., Leira, M., Marco-Barba, J., González-Sampériz, P.,
1564 Valero-Garcés, B., López-Sáez, J., Santos, L., Mata, P., Ito, E., 2011.
1565 Revealing the last 13,500 years of environmental history from the multiproxy
1566 record of a mountain lake (Lago Enol, northern Iberian Peninsula). *Journal*
1567 *of Paleolimnology*, 46(3): 327-349.

1568 Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., Martrat,
1569 B., González-Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J.P.,
1570 Belmonte, Á., Sancho, C., Cacho, I., Herrera, G., Canals, M., Grimalt, J.O.,
1571 Jiménez-Espejo, F., Martínez-Ruiz, F., Vegas-Vilarrúbia, T., Valero-Garcés,
1572 B.L., 2012. The Medieval Climate Anomaly in the Iberian Peninsula
1573 reconstructed from marine and lake records. *Quaternary Science Reviews*,
1574 43(0): 16-32.

1575 Moreno, A., Valero-Garcés, B., González-Sampériz, P., Navas, A., Machín, J.,
1576 Delgado-Huertas, A., 2004. El registro paleoambiental y paleoclimático de
1577 las saladas de la Playa y la Salineta (Zona Central de la Depresión del
1578 Ebro). *Geotemas*, 6(5): 137-140.

1579 Moreno, A., Valero-Garcés, B., González-Sampériz, P., Rico, M., 2008. Flood
1580 response to rainfall variability during the last 2000 years inferred from the
1581 Taravilla Lake record (Central Iberian Range, Spain). *Journal of*
1582 *Paleolimnology*, 40(3): 943-961.

1583 Moreno, A., Valero-Garcés, B.L., Jiménez-Sánchez, M., Domínguez-Cuesta, M.J.,
1584 Mata, M.P., Navas, A., González-Sampériz, P., Stoll, H., Farias, P.,

1585 Morellón, M., Corella, J.P., Rico, M., 2010. The last deglaciation in the Picos
 1586 de Europa National Park (Cantabrian Mountains, northern Spain). *Journal of*
 1587 *Quaternary Science*, 25(7): 1076-1091.

1588 Murphy, D.H., Wilkinson, B.H., 1980. Carbonate deposition and facies distribution
 1589 in a central Michigan marl lake. *Sedimentology*, 27(2): 123-135.

1590 Negendank, J.F.W., Zolitschka, B., 1993. *Paleolimnology of European Maar Lakes*.
 1591 Springer-Verlag, New York.

1592 Nicod, J., 1999. Phénomènes karstiques et mouvements de terrain récents dans le
 1593 Dépt. du Var; Risques naturels (Avignon, 1995). In: CTHS (Ed.), Paris, pp.
 1594 115-130.

1595 Nicod, J., 2006. Lakes in gypsum karst: some examples in alpine and
 1596 mediterranean countries. *Acta Carsologica*, 35(1): 69-78.

1597 Nicoud, G., Manalt, F., 2001. The lacustrine depression at Annecy (France),
 1598 geological setting and Quaternary evolution. *Journal of Paleolimnology*,
 1599 25(2): 137-147.

1600 O'Sullivan, P.E., 1983. Annually-laminated lake sediments and the study of
 1601 Quaternary environmental changes — a review. *Quaternary Science*
 1602 *Reviews*, 1(4): 245-313.

1603 Ohlendorf, C., Sturm, M., 2001. Precipitation and Dissolution of Calcite in a Swiss
 1604 High Alpine Lake. *Arctic, antarctic, and alpine research*, 33(4): 410-417.

1605 Ohlendorf, C., Sturm, M., Hausmann, S., 2003. Natural environmental changes
 1606 and human impact reflected in sediments of a high alpine lake in
 1607 Switzerland. *Journal of Paleolimnology*, 30(3): 297-306.

1608 Palmquist, R., 1979. Geologic controls on doline characteristics in mantled karst.
1609 Zeitschrift fuer Geomorphologie Suppl Bd, 32: 90-106.

1610 Pedley, M., Andrews, J., Ordonez, S., del Cura, M.A.G., Martin, J.-A.G., Taylor, D.,
1611 1996. Does climate control the morphological fabric of freshwater
1612 carbonates? A comparative study of Holocene barrage tufas from Spain and
1613 Britain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 121(3-4):
1614 239-257.

1615 Pérez-Bielsa, C., Lambán, L., Plata, J., Rubio, F., Soto, R., 2012. Characterization
1616 of a karstic aquifer using magnetic resonance sounding and electrical
1617 resistivity tomography: a case-study of Estaña Lakes (northern Spain).
1618 *Hydrogeology Journal*, 20(6): 1045-1059.

1619 Pérez-Obiol, R., Julià, R., 1994. Climatic Change on the Iberian Peninsula
1620 Recorded in a 30,000-Yr Pollen Record from Lake Banyoles. *Quaternary*
1621 *Research*, 41(1): 91-98.

1622 Pérez-Sanz, A., González-Sampériz, P., Moreno, A., Valero-Garcés, B., Gil-
1623 Romera, G., Rieradevall, M., Tarrats, P., Lasheras-Álvarez, L., Morellón, M.,
1624 Belmonte, Á., Sancho, C., Sevilla-Callejo, M., Navas, A., 2013. Holocene
1625 climate variability, vegetation dynamics and fire regime in the Central
1626 Pyrenees: The Basa de la Mora sequence (NE Spain). *Quaternary Science*
1627 *Reviews*, in press.

1628 Platt, N.H., Wright, V.T., 1991. Lacustrine carbonates: facies models, facies
1629 distribution and hydrocarbon aspects. In: P. Anadón, L. Cabrera, K. Kelts
1630 (Eds.), *Lacustrine facies analysis*. International Association of
1631 *Sedimentologists Special Publication*, pp. 57-74.

- 1632 Prat, N., Rieradevall, M., 1995. Life cycle and production of Chironomidae (Diptera)
1633 from Lake Banyoles (NE Spain). *Freshwater Biology*, 33(3): 511-524.
- 1634 Pueyo, J.J., Sáez, A., Giralt, S., Valero-Garcés, B.L., Moreno, A., Bao, R.,
1635 Schwalb, A., Herrera, C., Klosowska, B., Taberner, C., 2011. Carbonate and
1636 organic matter sedimentation and isotopic signatures in Lake Chungará,
1637 Chilean Altiplano, during the last 12.3 kyr. *Palaeogeography*,
1638 *Palaeoclimatology*, *Palaeoecology*, 307(1-4): 339-355.
- 1639 Pulido-Bosch, A., 1989. Les gypses triasiques de Fuente Camacho, Réunion
1640 franco-espagnole sur les karsts d 'Andalousie, pp. 65-82.
- 1641 Renault, R.W., Last, W.M., 1994. Sedimentology and geochemistry of modern and
1642 ancient saline lakes, 50. Society for Sedimentary Geology Special
1643 Publication, Tulsa.
- 1644 Renaut, R.W., Gierlowski-Kordesch, E.H., 2010. Lakes. In: R. Dalrymple, N. James
1645 (Eds.), *Facies Models*. Geological Association of Canada, Toronto
1646 (Canada), pp. 541-575.
- 1647 Reynolds, C.S., Walsby, A.E., 1975. Water-blooms. *Biological Reviews*, 50(4):
1648 437-481.
- 1649 Rico, E., Chicote, A., González, M.E., Montes, C., 1995. Batimetría y análisis
1650 morfométrico del lago de Arreo (N. España). *Limnética*, 1(11): 55-58.
- 1651 Rodrigo, M.A., Vicente, E., Miracle, M.R., 1993. Short-term calcite precipitation in
1652 the karstic meromictic lake La Cruz (Cuenca, Spain). *Verh. Internat. Verein.*
1653 *Limnol.*, 25: 711-719.

1654 Romero-Viana, L., Julià, R., Camacho, A., Vicente, E., Miracle, M., 2008. Climate
1655 signal in varve thickness: Lake La Cruz (Spain), a case study. *Journal of*
1656 *Paleolimnology*, 40(2): 703-714.

1657 Romero-Viana, L., Julià, R., Schimmel, M., Camacho, A., Vicente, E., Miracle,
1658 M.R., 2011. Reconstruction of annual winter rainfall since A.D.1579 in
1659 central-eastern Spain based on calcite laminated sediment from Lake La
1660 Cruz. *Climatic Change*, 107(3-4): 343-361.

1661 Romero-Viana, L., Miracle, M.R., López-Blanco, C., Cuna, E., Vilaclara, G., Garcia-
1662 Orellana, J., Keely, B., Camacho, A., Vicente, E., 2009. Sedimentary
1663 multiproxy response to hydroclimatic variability in Lagunillo del Tejo (Spain).
1664 *Hydrobiologia*, 631(1): 231-245.

1665 Romero, L., Camacho, A., Vicente, E., Miracle, M., 2006. Sedimentation patterns
1666 of photosynthetic bacteria based on pigment markers in meromictic Lake La
1667 Cruz (Spain): paleolimnological Implications. *Journal of Paleolimnology*,
1668 35(1): 167-177.

1669 Romero Viana, L., 2007. Paleolimnological study in two karstic lakes: climate
1670 signal in varved sediment and phototrophic organisms variability, Universitat
1671 de València, València (España), 174 pp.

1672 Sadori, L., Narcisi, B., 2001. The Postglacial record of environmental history from
1673 Lago di Pergusa, Sicily. *The Holocene*, 11(6): 655-671.

1674 Sánchez, M., Fernández-Delgado, C., F.J., S.-P., 1992. Nuevos datos acerca de la
1675 morfometría y batimetría de la Laguna de Zoñar (Aguilar de la Frontera,
1676 Córdoba). *Oxyura*, 6(73-77).

1677 Sancho-Marcén, C., 1988. El Polje de Saganta (Sierras Exteriores pirenaicas,
1678 prov. de Huesca). Cuaternario y Geomorfología, 2(1-4): 107-113.

1679 Sanz-Montero, M.E., Rodríguez-Aranda, J.P., García del Cura, M.A., 2008.
1680 Dolomite-silica stromatolites in Miocene lacustrine deposits from the Duero
1681 Basin, Spain: the role of organotemplates in the precipitation of dolomite.
1682 Sedimentology, 55(4): 729-750.

1683 Sanz, M., 1981. El sistema hidrogeológico de Banyoles-La Garrotxa. Unpublished
1684 PhD thesis, Universitat Autònoma de Barcelona, Barcelona, Spain.

1685 Scussolini, P., Vegas-Vilarrúbia, T., Rull, V., Corella, J., Valero-Garcés, B., Gomà,
1686 J., 2011. Middle and late Holocene climate change and human impact
1687 inferred from diatoms, algae and aquatic macrophyte pollen in sediments
1688 from Lake Montcortès (NE Iberian Peninsula). Journal of Paleolimnology,
1689 46(3): 369-385.

1690 Schmidt, R., Müller, J., Drescher-Schneider, R., Krisai, R., Szeroczynska, K., Baric,
1691 A., 2000. Changes in lake level and trophy at Lake Vrana, a large karstic
1692 lake on the Island of Cres (Croatia), with respect to palaeoclimate and
1693 anthropogenic impacts during the last approx. 16,000 years. Journal of
1694 Limnology, 59(2): 113-130.

1695 Schrag, D.P., Higgins, J.A., Macdonald, F.A., Johnston, D.T., 2013. Authigenic
1696 Carbonate and the History of the Global Carbon Cycle. Science, 339(6119):
1697 540-543.

1698 Schreiber, B.C., Tabakh, M.E., 2000. Deposition and early alteration of evaporites.
1699 Sedimentology, 47(s1): 215-238.

1700 Serra, T., Colomer, J., Gacia, E., Soler, M., Casamitjana, X., 2002. Effects of a
 1701 turbid hydrothermal plume on the sedimentation rates in a karstic lake.
 1702 Geophys. Res. Lett., 29(21): 2029.

1703 Serra, T., Soler, M., Julià, R., Casamitjana, X., Colomer, J., 2005. Behaviour and
 1704 dynamics of a hydrothermal plume in Lake Banyoles, Catalonia, NE Spain.
 1705 Sedimentology, 52(4): 795-808.

1706 Shaw, B., Klessig, L., Mechenich, C., 2002. Understanding Lake Data, G3582.
 1707 University of Wisconsin Cooperative Extension., Madison (Wisconsin).

1708 Smoot, J.P., Lowenstein, T., 1991. Depositional environments of non-marine
 1709 evaporites. In: J. Melvin (Ed.), Evaporites, petroleum and mineral resources.
 1710 Developments in Sedimentology. Elsevier, Amsterdam, pp. 189-348.

1711 Sondi, I., Jura Ci, M., 2009. Whiting events and the formation of aragonite in
 1712 Mediterranean Karstic Marine Lakes: new evidence on its biologically
 1713 induced inorganic origin. Sedimentology, 57(1): 85-95.

1714 Stabel, H., 1986. Calcite precipitation in Lake Constance: chemical equilibrium,
 1715 sedimentation, and nucleation by algae. Limnol. Oceanogr., 31: 1081-1093.

1716 Strasser, M., Stegmann, S., Bussmann, F., Anselmetti, F.S., Rick, B., Kopf, A.,
 1717 2007. Quantifying subaqueous slope stability during seismic shaking: Lake
 1718 Lucerne as model for ocean margins. Marine Geology, 240(1-4): 77-97.

1719 Sturm, M., Matter, A., 1978. Turbidites and varves in Lake Brienz (Switzerland):
 1720 deposition of clastic detritus by density currents. Special Publications of the
 1721 International Association of Sedimentologists, 2: 147-168.

1722 Sumner, G., Homar, V., Ramis, C., 2001. Precipitation seasonality in eastern and
 1723 southern coastal Spain. International Journal of Climatology, 21(2): 219-247.

- 1724 Talbot, M.R., Allen, P.A., 1996. Lakes In: H.G. Reading (Ed.), *Sedimentary*
 1725 *Environments: processes, facies and stratigraphy*. Blackwell Science,
 1726 Oxford, pp. 83-124.
- 1727 Teranes, J.L., McKenzie, J.A., Bernasconi, S.M., Lotter, A.F., Sturm, M., 1999a. A
 1728 study of oxygen isotopic fractionation during bio-induced calcite precipitation
 1729 in eutrophic Baldeggersee, Switzerland. *Geochimica et Cosmochimica Acta*,
 1730 63(13–14): 1981-1989.
- 1731 Teranes, J.L., McKenzie, J.A., Lotter, A.F., Sturm, M., 1999b. Stable isotope
 1732 response to lake eutrophication: calibration of a high-resolution lacustrine
 1733 sequence from Baldeggersee, Switzerland. *Limnology and Oceanography*,
 1734 44: 320–333.
- 1735 Treese, K.L., Wilkinson, B.H., 1982. Peat-marl deposition in a Holocene paludal-
 1736 lacustrine basin—Sucker Lake, Michigan. *Sedimentology*, 29(3): 375-390.
- 1737 Valero-Garcés, B., Delgado-Huertas, A., Navas, A., Machín, J., González-
 1738 Sampériz, P., Kelts, K., 2000. Quaternary palaeohydrological evolution of a
 1739 playa lake: Salada Mediana, central Ebro Basin, Spain. *Sedimentology*, 47:
 1740 1135-1156.
- 1741 Valero-Garcés, B., González-Sampériz, P., Navas, A., Machín, J., Mata, P.,
 1742 Delgado-Huertas, A., Bao, R., Moreno, A., Carrión, J.S., Schwalb, A.,
 1743 González-Barrios, A., 2006. Human impact since medieval times and recent
 1744 ecological restoration in a Mediterranean lake: the Laguna Zoñar, southern
 1745 Spain. *Journal of Paleolimnology*, 35: 441-465.
- 1746 Valero-Garcés, B., Navas, A., Mata, P., Delgado-Huertas, A., Machín, J.,
 1747 González-Sampériz, P., Moreno, A., Schwalb, A., Ariztegui, D.,

1748 Schnellmann, M., Bao, R., González-Barrios, A., 2003. Sedimentary facies
 1749 analyses in lacustrine cores: from initial core descriptions to detailed
 1750 palaeoenvironmental reconstructions. A case study from Zoñar Lake
 1751 (Cordoba province, Spain). In: B. Valero-Garcés (Ed.), *Limnogeology in*
 1752 *Spain: a tribute to Kerry Kelts*. Biblioteca de Ciencias. C.S.I.C., Madrid, pp.
 1753 385-414.

1754 Valero-Garcés, B., Zeroual, E., Kelts, K., 1998. Arid phases in the Western
 1755 Mediterranean Region during the last glacial cycle reconstructed from
 1756 lacustrine records. In: G. Benito, Baker, V.R., Gregory, K.J. (Ed.),
 1757 *Palaeohydrology and Environmental Change*. John Wiley & Sons Ltd., pp.
 1758 67-80.

1759 Valero-Garcés, B.L., Kelts, K.R., 1995. A sedimentary facies model for perennial
 1760 and meromictic saline lakes: Holocene Medicine Lake Basin, South Dakota,
 1761 USA. *Journal of Paleolimnology*, 14(2): 123-149.

1762 Valero-Garcés, B.L., Moreno, A., Navas, A., Mata, P., Machín, J., Delgado-
 1763 Huertas, A., González-Sampériz, P., Schwalb, A., Morellón, M., Cheng, H.,
 1764 Edwards, R.L., 2008. The Taravilla lake and tufa deposits (Central Iberian
 1765 Range, Spain) as palaeohydrological and palaeoclimatic indicators.
 1766 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 259(2-3): 136-156.

1767 Valero Garcés, B., Morellón, M., Martín Puertas, C., Moreno, A., Corella, P.,
 1768 González Sampériz, P., Rico, M., Mata, P., Navas, A., 2009. A
 1769 paleolimnological perspective of three karstic lake (Taravilla, Zoñar and
 1770 Estanya): Sedimentological and hydrogeological evolution climate and

1771 human impact and implications for management and restoration policies.
 1772 Boletín Geológico y Minero, 120(3): 479-496.
 1773 Valero Garcés, B.L., Moreno, A., Navas, A., Mata, P., Machín, J., Delgado
 1774 Huertas, A., González Sampériz, P., Schwalb, A., Morellón, M., Cheng, H.,
 1775 Edwards, R.L., 2008. The Taravilla lake and tufa deposits (Central Iberian
 1776 Range, Spain) as palaeohydrological and palaeoclimatic indicators.
 1777 Palaeogeography, Palaeoclimatology, Palaeoecology, 259(2-3): 136-156.
 1778 Vicente, E., Camacho, A., Rodrigo, M.A., 1993. Morphometry and physico-
 1779 chemistry of the crenogenic meromictic Lake 'El Tobar' (Spain). Verh. Int.
 1780 Ver. Limnol. , 25: 698-704.
 1781 Vicente, E., Miracle, M.R., 1987. Physicochemical and microbial stratification in a
 1782 meromictic karstic lake of Spain. Verh. Internat. Verein. Limnol., 23: 522-
 1783 529.
 1784 Vogel, H., Wagner, B., Zanchetta, G., Sulpizio, R., Rosén, P., 2010. A paleoclimate
 1785 record with tephrochronological age control for the last glacial-interglacial
 1786 cycle from Lake Ohrid, Albania and Macedonia. Journal of Paleolimnology,
 1787 44(1): 295-310.
 1788 Wetzel, R.G., 2001. Limnology: lake and river ecosystems. Academic Press, San
 1789 Diego (USA).
 1790 Wilhelm, B., Arnaud, F., Sabatier, P., Crouzet, C., Brisset, E., Chaumillon, E.,
 1791 Disnar, J.-R., Guiter, F., Malet, E., Reyss, J.-L., Tachikawa, K., Bard, E.,
 1792 Delannoy, J.-J., 2012. 1400 years of extreme precipitation patterns over the
 1793 Mediterranean French Alps and possible forcing mechanisms. Quaternary
 1794 Research, 78(1): 1-12.

1795 Wright, V.P., 1990. Lacustrine carbonates. In: M.E. Tucker, Wright, V. P. (Ed.),
1796 Carbonate Sedimentology. Blackwell Scientific Publications, Oxford, pp.
1797 164-190.

1798 Zanchetta, G., Borghini, A., Fallick, A., Bonadonna, F., Leone, G., 2007. Late
1799 Quaternary palaeohydrology of Lake Pergusa (Sicily, southern Italy) as
1800 inferred by stable isotopes of lacustrine carbonates. Journal of
1801 Paleolimnology, 38(2): 227-239.

1802 Zolitschka, B., 2007. Varved lake sediments. In: S.A. Elias (Ed.), Encyclopedia of
1803 Quaternary Science. Elsevier, Amsterdam, pp. 3105-3114.

1804

1805

1806

Tables

Table 1. Main characteristics of selected Iberian karst lakes, including physical, limnological and hydrochemical parameters and main literature references.

Table 2. Facies and Facies Associations in Iberian karst lakes, including sedimentology, depositional processes, and environments

Figure captions

Figure 1. Simplified geological map of Spain (after Gibbons and Moreno, 2002) and location of the Iberian karst lakes sequences discussed (black dots) and reviewed (white dots) in this paper.

Figure 2. The diversity of karstic lakes in the Iberian Peninsula. Lake basins occur in fluvial drainages dammed by tufa buildups - Banyoles (A) and Taravilla (B) - in glaciated terrains - Enol (C) and Marboré (D) -and as simple – El Tejo (F), Montcortès (H) and Arreo (I) (Photo: Eugenio Rico) or multiple sinkholes – Estanya (E) and Zoñar (G). According to surface hydrology, lakes are open (through-flowing) as Banyoles (A), Taravilla (B), Enol (C), Marboré (D) and Montcortès (H); closed with no surface outlet as Estanya (E) and El Tejo (F) or with a seasonal outlet in the recent past as Zoñar (G) and Arreo (I).

Figure 3. Core photographs of some diagnostic facies: A. Taravilla littoral facies association: Coarse clastic (LC) and Banded carbonate-rich silts and muds (LF) B. Zoñar varves (DLV(c) grading upwards to massive carbonate silts (LF) C. Varves with calcite laminae (DLV (c) with intercalated graded (DG) in Zoñar. D. Arreo Littoral Facies LF: Banded carbonate silts and muds. E. Arreo Distal Facies: Varves with calcite laminae DLV(c) (black), gypsum-rich laminites DLL(g) and massive DM distal facies. F. Arreo Distal Facies: Varves with calcite laminae DLV(c) (black) and graded DG distal facies. G. Montcortés distal facies: Varves with calcite laminae DLV(c) with intercalated graded DG distal facies. H. Varves with calcite laminae DLV(c) with intercalated graded DG distal facies. I. Organic varves DLV(o) in Montcorès. J. Enol Rhythmites (DLR facies). Scale bars: 10 cm

Figure 4. Littoral and palustrine facies in the sedimentary sequence of Lake Arreo: (modified from Corella et al., 2011a, 2013). From left to right, sequence includes depth, sedimentological units, core photograph, facies, magnetic susceptibility, TIC (Total Inorganic Carbon), TOC (Total Organic Carbon), TOC/TN ratio and selected mineralogy (Cl=clay minerals, Gy=Gypsum, Qz+Pl=Quartz + Plagioclases, Cc=calcite) and age model. Also shown, bathymetric map and location of the sediment core.

Figure 5. Clastic dominated littoral facies and massive/graded turbidites in the sedimentary sequence of Lake Taravilla (modified from Moreno et al., 2008). From left to right, sequences include depth, sedimentological units, core photograph,

1854 facies, magnetic susceptibility, lightness (core 2A) and also TIC (Total Inorganic
1855 Carbon), TOC (Total Organic Carbon) and TOC/TN ratio (core 1A) and age model.
1856 Also shown, bathymetric map and location of the sediment cores.

1857

1858 Figure 6. Littoral facies in Lake La Parra (modified from Barreiro-Lostres, 2012).
1859 From left to right, sequence includes depth, sedimentological units, core
1860 photograph, facies, TIC (Total Inorganic Carbon), TOC (Total Organic Carbon),
1861 TOC/TN ratio and mineralogy (Dol+Qz+Pl= dolomite + quartz + plagioclase;
1862 Cc+Qz+CMg = calcite + aragonite + magnesium-rich calcite; Cl = clay minerals)
1863 and age model. Also shown, bathymetric map and location of the sediment core.

1864

1865 Figure 7. Littoral and distal facies in the sedimentary sequence of Lake La Basa de
1866 la Mora (modified from (Pérez-Sanz et al., 2013)). From left to right, sequence
1867 includes depth, sedimentological units, core photograph, facies, magnetic
1868 susceptibility, TIC (Total Inorganic Carbon), TOC (Total Organic Carbon) and
1869 TOC/TN ratio and age model. Also shown contour map and location of the
1870 sediment cores. Maximum water depth is 4 m.

1871

1872 Figure 8. Distal facies (varves, laminites and massive/graded) in Lake Montcortés
1873 (modified from Corella et al., 2011). From left to right, the sequence include depth,
1874 sedimentological units, core photograph, facies, magnetic susceptibility, TIC (Total
1875 Inorganic Carbon), TOC (Total Organic Carbon), TOC/TN ratio and mineralogy
1876 (Qz=quartz, Cc=calcite; Cl=clay minerals; Gy=gypsum) and age model. Also
1877 shown bathymetric map and location of the sediment cores.

1878

1879 Figure 9. Distal and littoral facies in Lake Zoñar (modified from Martín-Puertas et
1880 al., 2008). From left to right, both sequences include depth, sedimentological units,
1881 core photograph, facies, magnetic susceptibility, TIC (Total Inorganic Carbon),
1882 TOC (Total Organic Carbon), TOC/TN ratio and mineralogy (Qz=quartz,
1883 Cc=calcite; Ar= aragonite, Gy=gypsum; Cl=clay minerals) and age model. Also
1884 shown bathymetric map and location of the sediment cores.

1885

1886 Figure 10. Microphotographs of laminated carbonates. A. Varve sequences in
1887 Montcortès (XN light) and Zoñar (PN light). B. SEM photographs of endogenic and
1888 clastic carbonates. C. Fining-upward and coarsening-upward textures in calcite
1889 laminae from Zoñar varves.

1890

1891 Figure 11. Distal facies with gypsum-rich sediments (laminites) and littoral facies in
1892 Lake Estanya (modified from Morellón et al., 2010). From left to right, sequence
1893 includes depth, sedimentological units, core photograph, facies, magnetic
1894 susceptibility, TIC (Total Inorganic Carbon), TOC (Total Organic Carbon) and
1895 TOC/TN ratio and mineralogy and age model. Also shown bathymetric maps and
1896 location of the sediment cores.

1897

1898 Figure 12. Distal facies (laminites, varves, massive and graded) in Lake Arreo
1899 (modified from Corella et al., 2011). From left to right, sequence includes depth,
1900 sedimentological units, core photograph, facies, magnetic susceptibility, TIC (Total
1901 Inorganic Carbon), TOC (Total Organic Carbon), TOC/TN ratio and mineralogy

1902 (Qz=quartz, Cl=clay minerals, Cc=calcite;Dol= dolomite; Gy=gypsum) and age
1903 model. Also shown bathymetric maps and location of the sediment core.

1904

1905 Figure 13. Rhythmites and laminated facies in the Lake Enol sequence (modified
1906 from Moreno et al., 2010). From left to right, sequences include depth,
1907 sedimentological units, core photograph, facies, magnetic susceptibility, lightness,
1908 bulk density, TIC (Total Inorganic Carbon) and TOC (Total Organic Carbon) and
1909 age model. Also shown contour map and location of the sediment cores.

1910

1911 Figure 14. Distal facies (laminites and homogeneous) in Lake Banyoles. From left
1912 to right, sequence includes depth, sedimentological units, core photograph, facies,
1913 Lightness, TIC (Total Inorganic Carbon) and TOC (Total Organic Carbon) and age
1914 model. Also included bathymetric map and location of the sediment cores.

1915

1916 Figure 15. A. A 3-D model illustrating the hydrologic settings of Iberian Karstic
1917 lakes. The lines with numbers correspond with the possible location of the cross-
1918 sections on B. B. Depositional environments in Iberian karst lakes: palustrine
1919 environments are well developed in Taravilla (higher alluvial influence) and Arreo
1920 (lower alluvial influence). Littoral carbonates are well developed in Arreo, Estanya,
1921 and Banyoles and during the mid Holocene in Montcortès. Distal-profunda facies
1922 occur in most lakes, although significant development of varves only in Montcortès
1923 and Zoñar. Sinkholes with active groundwater input only occur in Banyoles.

1924

Figure 16. Relationships among selected morphometric (lake and watershed surface area, maximum depth, lake surface/depth ratio), surface hydrology and salinity (electric conductivity) parameters of Iberian Karst lakes. A. Lake and watershed surface plot identifies single sinkholes (Tejo and La Parra) and complex large basins (Banyoles and Zoñar) at both ends of the spectrum; most lakes with smaller surface area have larger watersheds. B. Except for complex basin (> 30 ha lake surface), smaller lakes are usually deeper for both categories: lakes < 5 ha, and between 5-20 ha. C. Surface hydrology (1. Permanent outlet, 2. Ephemeral outlet; 3. No outlet) versus Electric Conductivity. High mountain lakes (Enol and Marboré) have the lowest EC values and lakes with no surface outlet (Estanya and Zoñar) the highest. However, high EC values occur in all hydrologic categories, and low EC values occur in lakes with no outlet but with strong groundwater input (La Parra, Tejo). D. Surface/Depth - EC plot does not show trends between morphometry and lake salinity although data are grouped in three categories: smaller and deeper lakes (low S/D ratios) have lower EC; lakes with intermediate values show a negative trend between S/D and EC; higher EC values occur in larger lakes.

Figure 17. A classification of Iberian Karstic lakes based on surface hydrology and sediment input.

Table 1. Main characteristics of selected Iberian karst lakes, including physical, limnological and hydrochemical parameters and main literature references. Hydrology classification: 1. Open (through-flowing) lake (Water input >> Water output), permanent surface outflow; 2. Open lake (Water input > water output), seasonal outflow; 3. Closed lake (Water input < water output); no surface outflow. Geologic setting: 1. Glaciated terrain; 2. Fluvial environment with tufa dams; 3. Sinkhole in dominant carbonate formations; 4. Sinkhole in carbonate/gypsum formations

Lake	Geologic setting	Location	Altitude (m asl)	Surface Area (ha)	Maximum depth (m)	Watershed surface area (ha)	Hydrochemistry	Thermal regime	Surface Hydrology	Main References
Banyoles	3	42°1'N; 2°4'E	173	118	40 (80)	1142	Alkalinity: 3.87 meq L ⁻¹ pH = 7-8.2; [SO ₄ ²⁻] > [HCO ₃ ²⁻] > [Ca ²⁺] EC = 900-2000 µS cm-1	Monomictic with meromictic sub-basins	1: Open; permanent inlet/outlet	Juliá, (1980) Valero-Garcés et al., (1998) Höbig et al. (2012)
El Tejo	3	39°N, 1°52'W	1000	1,5	27	4	Alkalinity= 3.05 meq L ⁻¹ pH = 8.7; [HCO ₃ ²⁻] > [Ca ²⁺] EC = 540 µS cm-1	Monomictic to meromictic	3: Closed; Endorheic	This paper
El Tobar	4	40°32'N; 3°56'W	1200	16.2 ha	Holomictic: 12 (Meromictic: 20)	129	Alkalinity= 2.12 meq L ⁻¹ Mixolimnion: ([HCO ₃ ²⁻] > [Ca ²⁺]-[Mg ²⁺]) pH = 8; EC=600 µS cm-1 Monimolimnion: Alkalinity= 4.33 meq L ⁻¹ [NaCl ¹⁻] > [Ca ²⁺] > [Mg ²⁺] > [SO ₄ ²⁻] EC = 2000 µS cm-1	Meromictic	2:Open; permanent inlet/outlet	Vicente et al. (1993)
La Parra	3	39°59'N, 1°52'W	1000	1.13	17.5	20	Alkalinity= 6.0 meq L ⁻¹ pH = 8.3; [HCO ₃ ²⁻] > [Ca ²⁺] EC=335 µS cm-1	Holomictic	3: Closed; Endorheic	Barreiro-Lostres et al. (2011, 2013)
Taravilla	2	40°39' N 1°58' W	1100	2.11	12	550	Alkalinity = 2.92 meq L ⁻¹ pH = 7.8; [Ca ²⁺] > [HCO ₃ ²⁻] > [SO ₄ ²⁻] EC =550 µS cm-1	Holomictic	1: Open; permanent inlet/outlet	Valero-Garcés et al. (2008)
Enol	1	43°11' N 4°09'W	1070	12.2	22	150	Alkalinity = 2.4 meq l-1; pH = 7.7–8.2 ([HCO ₃ ²⁻] > [Ca ²⁺] > [SO ₄ ²⁻]) EC: 202 µS cm-1	Monomictic	Open; permanent inlet/outlet	Moreno et al., (2011)
Marboré	1	00°23' E, 42°41' N	2592	14.3	27	137	Alkalinity = 0.8 meq L ⁻¹ ; pH = 7- 8.5 EC: 54.5 – 72.0 µS/cm;	Cool monomictic	1: Open; permanent inlet/outlet	Salabarnada, (2011)
Arreo	4	42°46' N, 2°59' W	657	6.57	24.8	287	Alkalinity = 4.18 meq L ⁻¹ ; pH= 7.6 – 8.2 [Ca ²⁺] >[Mg ²⁺]>[Na ⁺]>[SO ₄ ²⁻] EC: 703-1727 µS/cm	Monomictic	2: Open; seasonal inlet/outlet	Corella, (2011)a, (2013)

Montcortés	3	42° 19´ N, 0° 59´ E	1027	9.3	30	300	Alkalinity = 2.5- 3.5 meq L ⁻¹ ; pH = 7-8.5 [HCO ₃ ²⁻]>[Ca ²⁺]>[SO ₄ ²⁻] EC = 372 µS/cm	Meromictic	1: Open; permanent inlet/outlet	Corella et al., (2011b; 2012)
Estanya	3	42°02' N, 0°32' E	670	18.83	20	106	Alkalinity = 2 -3.5 meq L ⁻¹ ; pH = 7.4 – 7.6 [SO ₄ ²⁻]>[Ca ²⁺]> [Mg ²⁺] EC = 3440 µS/cm	Monomictic	3: Closed; Endorheic	Morellón et al., (2009)
Zoñar	3	37°29'00"N, 4°41'22" W	300	37	14.5	876	Alkalinity = 3 – 5 meq L ⁻¹ pH = 7.1 -8.4 [Cl ⁻]>[HCO ₃ ²⁻]> [SO ₄ ²⁻]>[Na ⁺] EC = 2500-3000 µS cm-1	Monomictic	3: Closed; Endorheic	Martín- Puertas et al. (2008)
Basa de la Mora	1	42°32' N 0°19' E	1914	5.5	3.5	462	Alkalinity= 0.6 meq L ⁻¹ pH= 8.6 [Ca ²⁺]>[SO ₄ ²⁻]> [Mg ²⁺] EC = 215 µS cm-1	Holomictic	2: Open; seasonal inlet/outlet	Pérez-Sanz et al. (2013)

Table 2. Facies and Facies Associations in Iberian karst lakes, including sedimentology and depositional processes and environments

<i>FACIES</i>	<i>Sedimentological features</i>	<i>Depositional Processes and Environments</i>
PALUSTRINE FACIES ASSOCIATION (P)		
P.C. Coarse clastic.	Up to 25 cm thick gravel beds organized in fining-upwards sequences, composed of bedrock and/or tufa clasts embedded within a carbonate and quartz sandy matrix with abundant plant remains	Wetland, strong alluvial influence
P.F. Fine clastic.	Cm to dm thick carbonate-rich coarse silts with edaphic textures, bioturbation and evidences of subaerial exposure	Wetland, low alluvial influence
P.O. Organic- rich.	Cm to dm thick, massive organic-rich silts with abundant plant remains	Wetland
LITTORAL FACIES ASSOCIATION (L)		
L.C. Coarse clastic.	Meter thick fining upward sequences composed of carbonate-rich silty-sand to coarse sand; massive to faintly laminated carbonate silt and massive to faintly laminated, dark grey carbonate silt and mud. Occasionally centimetre-thick silt layers with abundant organic rests are also deposited.	Littoral, high energy, variable alluvial influence
L.F. Massive to banded carbonate-rich silts and muds.	Dm- to cm thick layers with variable amounts of silicates and carbonates, and significant content of biogenic carbonate (<i>Chara</i> , gastropods) and plant remains. Subfacies defined by banded textures with biogenic carbonate content	Littoral, low energy, low alluvial influence
L.F (s). Massive to banded carbonate-rich silts and muds with mottling and evaporite nodules..	Gypsum nodules and bioturbation features (root traces, coarse plant remains, mottling, mixed sediment textures) are common. Abundant gastropods and large mm to cm-size terrestrial plant remains	Littoral with subaerial exposure and saline stages
DISTAL FACIES ASSOCIATION (D)		
D.B. Banded to laminated carbonate silts.	Dm- thick layers with silty fraction composed of carbonates, quartz and abundant clay-rich matrix. Minor amounts of feldspars, high-magnesium calcite (HMC) and gypsum. Common biogenic components as aggregates of amorphous lacustrine organic matter, macrophyte remains and diatoms	Distal, variable bottom conditions (bioturbation, oxygenation)
D.L. Finely laminated D.L.V. Varves		
D.L.V (c). With calcite laminae.	Sets of dark-brown laminae (lacustrine organic matter, diatoms), yellowish mm-thick laminae (authigenic carbonates (calcite, aragonite, dolomite)), and rare grey carbonate silt laminae.	Distal, moderately deep, meromictic conditions
D.L.V (o). Without calcite laminae	Sets of mm-thick, brown to greenish, organic-rich laminae and mm-thick terrigenous grey silt layers.	Distal, Moderately deep, anoxic meromictic conditions
D.L.L. Laminites		
D.L.L(g). Gypsum- rich.	Sets of cm-thick alternating yellowish bands of gypsum with reworked biogenic carbonates, dark brown diatom ooze with lacustrine organic matter and lenticular gypsum and grey massive clays.	Distal, deep saline with anoxic bottom conditions
D.L.L (o). Organic-rich.	Organic laminites are composed of amorphous lacustrine organic matter, diatoms and some macrophyte remains, with minor amounts of clay minerals, calcite, dolomite and quartz. In some cases (Zofíar, Estanya) includes laminated microbial mats.	Distal, shallow saline
D.L.R. Rhythmites.	Carbonate-poor, dm- to meter thick layers composed of laminae defined by grain size, from clay to coarse silt and sand.	Distal, periglacial to pro-glacial environments
D.M/G/H. Massive to Graded.	Cm- to dm- thick layers with high magnetic susceptibility, graded textures and mixed carbonate and silicate composition. Presence of biogenic components and common plant remains	Distal, floods and turbidite currents
D.G. Graded.	They occur as: dm-thick, laminated to banded intervals with regular, sharp contacts, and ii) cm to mm-thick laminae with diffuse and irregular contacts, constituted by massive, graded sediments.	
D.M. Massive	They occur as cm- thick, massive layers with sharp and erosive basal contacts.	Distal, floods and mass wasting processes
D.H. Homogeneites.	Cm to dm thick, grey, massive layers composed by fine-grained carbonate-rich silts	Distal, Resuspension of sediments from active spring discharge in sinkholes
GRAVITATIONAL FACIES (G)		
- G.F. Convoluted.	Thin intervals of laminated facies with convoluted, folded and distorted textures	Distal, slumps and gravitational deposits
- G.C. Chaotic.	Heterometric coarse facies with gravel size-clasts and heterometric sandy matrix

Palustrine facies

- PC: Coarse clastic
- PO: Organic-rich

Littoral facies

- LC: Coarse clastic
- LF: Massive to banded carbonate-rich silts and muds

Littoral facies

- LC: Coarse clastic
- LF: Massive to banded carbonate-rich silts and muds

Distal facies

- DL: Finely laminated

Littoral facies

- LC: Coarse clastic
- LF: Massive to banded carbonate-rich silts and muds

Distal facies

- DLV (c): With calcite laminae

Littoral facies

- LF: Massive to banded carbonate-rich silts and muds
- LF (s): Massive to banded carbonate rich silts and muds with evaporites

Distal facies

- DB: Banded to laminated carbonate silts
- DL: Finely laminated

Littoral facies

- LF: Massive to banded carbonate-rich silts and muds
- LF (s): Massive to banded carbonate-rich silts and muds with evaporites

Distal facies

- DLV: Varves
- DLV (c): With calcite laminae
- DLL: Laminites
- DLL (g): Gypsum-rich
- DLL (o): Organic-rich

Littoral facies

- LF: Massive to banded carbonate-rich silts and muds
- LF (s): Massive to banded carbonate-rich silts and muds with evaporites

Distal facies

- DL: Finely laminated
- DLL (g): Gypsum- rich
- DLL (o): Organic-rich
- DG: Graded

Littoral facies

- LC: Coarse clastic
- LF: Massive to banded carbonate-rich silts and muds

Distal facies

- DL: Finely laminated
- DLR: Rhythmites

Distal facies

- DB: Banded to laminated carbonate silts
- DL: Finely laminated
- DG: Graded
- DH: Homogeneities

LAKES

FACIES

PALUSTRINE

Palustrine (PC & PF fcs)

LITTORAL

Coarse, clastics (LC fcs)

Massive banded carbonate-rich silts and muds (LF & LF(s) fcs)

DISTAL

Banded to laminated carbonate-silts (DB fcs)

Finely laminated (DL fcs)

Homogeinites (DH fcs)

Turbidites (DG fcs)

WATERSHED SURFACE

LARGE

INTERMEDIATE

SMALL

OPEN

INTERMEDIATE

CLOSED

BANYOLES

TARAVILLA

MONTCORTÈS

ENOL

MARBORÉ

INTERMEDIATE

ARREO

TOBAR

BASA MORA

CLOSED

ZOÑAR

ESTANYA

LA PARRA

EL TEJO /
LA CRUZ