Influence of the hydrodynamic conditions on the
accessibility of *Aristeus antennatus* and other
demersal species to the deep water trawl fishery
off the Balearic Islands (western Mediterranean)

Angel Amores^{*1}, Lucía Rueda², Sebastià Monserrat^{1,3}, Beatriz
Guijarro², Catalina Pasqual¹, and Enric Massutí²

¹Instituto Mediterráneo de Estudios Avanzados, IMEDEA
(UIB-CSIC). Palma de Mallorca, Spain

²Instituto Español de Oceanografía. Centre Oceanogràfic de les
Balears. Moll de Ponent s/n, 07015 Palma de Mallorca, Spain

³Departament de Física. Universitat de les Illes Balears (UIB).
Palma de Mallorca, Spain

November 20, 2013

^{*}aamores@imedea.uib-csic.es

Abstract

Monthly catches per unit of effort (CPUE) of adult red shrimp (*Aristeus antennatus*), reported in the deep water bottom trawl fishery developed on the Sóller fishing ground off northern Mallorca (Western Mediterranean), and the mean ocean surface vorticity in the surrounding areas are compared between 2000 and 2010. A good correlation is found between the rises in the surrounding surface vorticity and the drops in the CPUE of the adult red shrimp. This correlation could be explained by assuming that most of the surface vorticity episodes could reach the bottom, increasing the seabed velocities and producing sediment resuspension, which could affect the near bottom water turbidity. *A. antennatus* would respond to this increased turbidity disappearing from the fishing grounds, probably moving downwards to the deeper waters. This massive displacement of red shrimp specimens away from the fishing grounds would consequently decrease their accessibility to fishing exploitation. Similar although more intense responses, have been observed during the downslope shelf dense water current episodes that occurred in a submarine canyon, northeast of the Iberian peninsula. The proposed mechanism suggesting how the surface vorticity observed can affect the bottom sediments is investigated using a year-long moored near-bottom current meter and a sediment trap moored near the fishing grounds.

The relationship between vorticity and catches is also explored for fish species (*Galeus melastomus*, *Micromesistius poutassou*, *Phycis blennoides*) and other crustacean (*Geryon longipes* and *Nephrops norvegicus*), considered as by-catch of the deep water fishery in the area. Results appear to support the suggestion that the water turbidity generated by the vorticity episodes is significant enough to affect the dynamics of the demersal species.

27 1 Introduction

28 The decapod crustacean red shrimp, *Aristeus antennatus* (Risso, 1816), a demersal
29 species distributed throughout the Mediterranean and the north-eastern Atlantic,
30 from Portugal to the Cabo Verde Islands [Arrobas and Ribeiro-Cascalho, 1987],
31 mainly occurs in the muddy bottoms of the slope, between 400 and at least 3300
32 m [Sardà et al., 2004]. This species is one of the most valuable deep-water fishing
33 resources in the western and central basins of the Mediterranean, remaining at a
34 low level of exploitation in the eastern basin [Papaconstantinou and Kapisiris, 2001]
35 and revealing important bathymetric migrations [Relini et al., 2000]. However,
36 despite its wide bathymetric distribution, it is mainly exploited between 400 and
37 800 m depth, and is the target species of the well-developed deep water bottom
38 trawl fishery on the western basin slope [Sardà et al., 2003].

39 The trawl fleet operating off the Balearic Islands (western Mediterranean) is
40 characterized by its versatility, which is determined by the specific dynamics of
41 the resources, among other factors (e.g. sea conditions and fish market). Bottom
42 trawlers not only target different species, they also change the fishing location at
43 a given time of the year, as well as the fishing tactics during the same fishing
44 trip. Palmer et al. [2009] defined four fishing tactics in this fishery, related to the
45 exploitation of different bathymetric strata and target species.

46 The annual catches of the red shrimp in the Balearic Islands are estimated to be
47 around 100–200 t, which represents 10% of the landings and 40% of the earnings
48 in the trawl fishery [Guijarro et al., 2012]. Sóller, one of the most important
49 fishing grounds for red shrimp around the Balearic Islands, is situated North
50 of Mallorca (solid black line area in Fig. 1), where an important part of the
51 island fleet is concentrated during the summer months [Moranta et al., 2008],
52 when catches of large specimens occur. The red shrimp population in this fishing
53 ground shows important seasonal variations throughout the year (such as the high
54 abundance of juveniles recruiting to the fishing grounds in autumn-winter and the
55 high abundance of large spawning females during the summer), compared with
56 the other nearby fishing grounds, south of Mallorca [Guijarro et al., 2008].

57 The Sóller fishing ground is located on the island slope, in a well known very
58 active area, with numerous eddies normally generated by some instabilities of
59 the Northern current or the Balearic current (Fig. 1), particularly more intense
60 during winter (October-March; Amores et al. [2013]). These eddies, clearly visible
61 on satellite images, have been known to reach the deeper waters, and their effects

are usually felt down to the seabed, where their velocities may increase to several times those of the mean currents measured in the zone [Amores et al., 2013]. These strong bottom currents of the order of 25 cm/s are known to produce sediment resuspension which, in turn, may generate additional cross slope turbidity currents [Thomson et al., 2010].

In the western Mediterranean, the red shrimp distribution, and its accessibility to fishing exploitation, has been shown to be mainly influenced by geomorphology [Sardà et al., 1994, 1997] and hydrodynamics [Bombace, 1975; Demestre and Martín, 1993; Ghidalia and Bourgois, 1961; Guijarro et al., 2008; Relini and Relini, 1987; Sardà et al., 2009]. These last factors are probably linked to regional and large-scale climatic patterns [Carbonell et al., 1999; Massutí et al., 2008; Maynou, 2008]. In a recent study, Company et al. [2008] revealed that the downslope shelf dense water current events into submarine canyons, along the whole northern Catalan margin, strongly affected the red shrimp landings. This downslope shelf dense water current events is one of the main processes contributing to the shelf-deep ocean exchange [Ivanov et al., 2004], enhancing organic-matter flux and deposition, increasing suspended particulate matter concentrations and transport of organic matter from coastal zones to the deep ocean [Bosley et al., 2004; Canals et al., 2006; Company et al., 2008]. In the northern Catalan margin, it exerts a negative effect on the catches of red shrimp and a positive effect for recruitment, due to the transportation of the particulate organic matter. The increase of suspended particulate matter also appears to be related to the abundance of other crustacean species such as pandalids and penaeid [Lin et al., 1992; Puig et al., 2001] and to an enhance of benthic productivity and biodiversity inside canyon habitats [Rowe et al., 1982; Schlacher et al., 2007; Vetter et al., 2010]. In addition to downslope shelf dense water current, mesoscale eddies have also been reported to be responsible of transport of shelf sediments to the deep ocean, resuspension of bottom sediments creating turbidity layers and formation of sediment plumes around their periphery [Washburn et al., 1993]. The influence of vorticity (as indicator of eddy development) on catchability of marine species has been mostly addressed for pelagic organisms such as tuna fisheries [Hyder et al., 2009; Kai and Marsac, 2010; Ramos et al., 1996; Zainuddin et al., 2006]. However, the effect of such physical processes has also been explored for benthic species, which are also linked to variables that describe water column properties and structures [Beentjesa and Renwick, 2001; Palamara et al., 2012].

97 The objective of this work is to analyze the possible links between the presence
98 of eddies (which will be quantified by their associated surface vorticity) affecting
99 the Sóller fishing ground and the red shrimp yields of the deep water trawl fishery
100 developed in the area. This relationship is also explored for other demersal species
101 frequently caught by the deep water bottom trawl fishery developed in the area
102 [Guijarro and Massutí, 2006], which consist of three fishes (*Galeus melastomus*,
103 *Micromesistius poutassou* and *Phycis blennoides*) and two decapod crustaceans
104 (*Geryon longipes* and *Nephrops norvegicus*), with the objective of discussing their
105 different responses in relation to their living habits.

106 A year-long near-bottom current meter and a sediment trap moored near the
107 fishing grounds are used to infer the mechanism to explain how the surface vorticity
108 observed can affect the bottom sediments and, in turn, the red shrimp yields.

109 2 Data and Methods

110 2.1 Catches

111 Daily time series of the landings from the bottom trawl fleet have been obtained
112 from the official sale bills of OP Mallorca Mar, the fishery producer organization
113 of Mallorca, between 2000 and 2010 (both years included). Each daily sale bill
114 was assigned to one fishing tactic (FT) or a combination of them following the
115 methodology described by Palmer et al. [2009]. Landings were standardized to
116 CPUEs (catches per unit of effort), referred to as kilograms caught per day and
117 boat. For *A. antennatus*, only catches obtained from the middle slope fishing
118 tactic, developed between 600 and 800 m depth, have been considered, because
119 this is the target species for this FT. Moreover, the daily sale bills distinguished
120 red shrimp catches into two size categories (small and large) up to year 2004,
121 and three categories (small, medium and large) from 2004 to the present day.
122 According to Guijarro et al. [2008], two different categories were defined in order
123 to homogenize the available data, small (including individuals with a carapace
124 length $<32\text{mm}$) and medium-large (adults, with a carapace length $\geq 32\text{mm}$). For
125 this analysis, only those of the medium-large sized category, mainly adult females,
126 were considered. Juveniles are not taken into account for two reasons:

- 127 1. The fishing fleet mainly targets large individuals (adults) due to their higher
128 commercial value. This fact would surely provoke a bias when trying to
129 relate juvenile catches with abundances
- 130 2. Adult and juvenile red shrimps present a clear different bathymetric distri-
131 bution. Adult individuals are mainly located at the 500-800 m range, where
132 the fishing fleet is developed. But the highest concentrations of juveniles are
133 situated deeper than 1000 m Sardà et al. [2003], where the bottom trawl
134 fishery is forbidden. So juvenile catches do not properly reflect the juvenile
135 population abundances.

136 From the entire fleet that currently operates in Mallorca, only five boats regu-
137 larly fish in the zone of interest (Sóller) throughout the year (other boats fishing in
138 this area only in summer are not considered). Among these five boats, exclusively
139 two devote most of their efforts to red shrimp fishery along the middle slope and
140 they were the only ones finally considered for the analysis.

141 Finally, as a direct response to hydrodynamics changes in a daily basis is not
 142 expected, a monthly average was calculated according to the daily time series.
 143 We intentionally filtered out the high frequency variations in order to compute an
 144 integrated response in a longer time scale and take into account for the gaps in
 145 the data (that occurred, for example, on weekends or bad weather days).

146 Regarding the other species considered in this work (*G. melastomus*, *M. poutas-*
 147 *sou*, *P. blennoides*, *G. longipes* and *N. norvegicus*), all the boats and slope fishing
 148 tactics have been considered because, unlike *A. antennatus*, they are the ‘by-
 149 catch’ species of the deep water trawl fishery, and therefore, their abundance in
 150 daily landings is not as frequent as that of the red shrimp. The final time series
 151 for each species were also averaged monthly as the CPUEs in terms of kg of catch
 152 per day per boat.

153 2.2 Hydrodynamic Data

154 2.2.1 Satellite images

155 We estimated the relative vorticity ζ (from now on referred to as only vorticity)
 156 from the daily Sea Surface Height (SSH) satellite images with a map spacing
 157 of $1/8^\circ \times 1/8^\circ$, obtained from the merged satellite AVISO products available at
 158 <http://www.aviso.oceanobs.com>. The vorticity is calculated as the curl of the
 159 velocity field, but we only retain the third component as it represents the vorticity
 160 of a horizontal field. By considering the hydrostatic and homogeneous fluid, the
 161 final expression of ζ is:

$$\zeta = \frac{g}{f} \cdot \nabla^2 SSH \quad (1)$$

162 where g is the gravity acceleration, f the Coriolis parameter and ∇^2 the hori-
 163 zontal Laplacian.

164 After computing the daily vorticity fields, their absolute value was taken be-
 165 cause both, cyclonic ($\zeta > 0$) and anticyclonic episodes ($\zeta < 0$), were expected to
 166 have the same effect on the seabed velocities and sediment resuspension. Next,
 167 we computed the spatial average in the dashed rectangle, as shown in Fig. 1. The
 168 choice of the area is somewhat arbitrary. The size should be significantly greater
 169 than the fishing ground dimension because the eddy sizes are significantly greater,
 170 and their horizontal influence is not known. The best results are found when the
 171 area is selected to be large enough to include the Northern and the Balearic cur-

rents, potentially the eddy generators. Finally, the daily time series was averaged on a monthly basis in order to have the same time step as the time series of the catches.

2.2.2 Moorings

A mooring was deployed north-west of Mallorca ($39^{\circ}49.682'$ N – $02^{\circ}12.778'$ E; star in Fig. 1) between November 2009 and February 2011. Located around 900 m depth in the Mallorca slope, it had four CTD Seabird 37 (300, 500, 700 and 900 m) and two current meters Nortek Aquadopp (500 and 900 m). Moreover, the mooring had also a sediment trap placed 30 m above the bottom. The CTD sampling rate was 10 minutes, while the current meters recorded one value every 30 minutes. The sediment trap had a sampling interval of 10 days and it had 12 bottles. The combination of sampling rate and number of bottles made necessary a maintenance of the mooring every four months.

All the instruments operated perfectly during the entire period, except for the 500 and 900 m CTDs, which ran out of batteries in mid-December 2010 and mid-January 2011, respectively. The sediment trap worked well too, but the unavailability of boat lead to no recorded data between July 5 and September 21, 2010.

Sediment trap samples were wet-sieved through a 1 mm nylon mesh in order to retain the largest organisms. Swimmers smaller than 1 mm were manually removed under a dissecting microscope using fine tweezers. Finally, the sample was freeze-dried and weighed to calculate the Total Mass Flux (TMF).

Despite we did not have any direct measurement of turbidity, the Nortek Aquadopp current meters give us an estimation of it throughout the backscattering of the particles used to compute the velocity (as suggested by Lohrmann [2001]). Acoustic backscattering has been used as turbidity surrogate in different references such as Thomson et al. [2010].

2.3 Statistical Analysis

Quantification of the similarity between surface vorticity and time series of catches was performed with the correlation function. If we define V and C as the monthly anomalies (time series after subtracting the mean value) for vorticity and catches, respectively, the correlation between these two series is calculated as:

$$\rho_{VC}(lag) = \frac{1}{(N-1-|lag|) \cdot \sqrt{\sigma_{VV} \cdot \sigma_{CC}}} \begin{cases} \sum_{i=|lag|+1}^N V_i \cdot C_{i-|lag|} & \text{if } lag < 0 \\ \sum_{i=|lag|+1}^N V_{i-|lag|} \cdot C_i & \text{if } lag \geq 0 \end{cases} \quad (2)$$

204 with N being the series length, σ_{VV} the covariance of V and σ_{CC} the covariance
205 of C .

206 The significance level may be obtained with:

$$sig(lag) = \frac{T_q(0.99, N - |lag| - 2)}{\sqrt{N - |lag| - 2 + [T_q(0.99, N - |lag| - 2)]^2}} \quad (3)$$

207 where $T_q(0.99, D)$ is the t-student distribution with a significance of 99% and
208 D degrees of freedom.

209 3 Results and Discussion

210 A visual inspection of the monthly average vorticity and *A. antennatus* CPUE
211 time series appears to suggest that any increase in vorticity generally causes a
212 decrease in the CPUEs, although a decrease in vorticity does not cause an increase
213 in CPUEs (Fig. 2a). In fact, a negative correlation (i.e. both series are in
214 antiphase) between time series has been found (Fig. 2b).

215 In the following we will give several evidences supporting the fact that surface
216 vorticity affects red shrimp availability by modifying near bottom turbidity in the
217 fishing grounds. If we assume that any increase in vorticity affects the CPUE of the
218 red shrimp by producing a near bottom turbidity, which in turn would decrease
219 the resource availability, we could not expect that the opposite, a decrease in
220 vorticity, would immediately produce an increase in the CPUEs. Water turbidity
221 would persist for some period of time in the area after the end of any vorticity
222 episode and then the red shrimp response would be delayed. Therefore, the two
223 series have been modified, trying to take into account the different, although
224 expected, CPUE response to the increase and decrease in vorticity. Vorticity and
225 CPUE time derivatives have been computed. As we want to highlight the part
226 when the vorticity increases (positive derivative) and when the CPUE decreases
227 (negative derivative), the negative vorticity and positive CPUE derivatives were
228 artificially set at zero. Therefore, only the increases in vorticity and decreases in
229 CPUE are considered. Furthermore, the sign of the CPUE derivatives has also
230 been reversed for better visualization (Fig. 3a).

231 Vorticity and *A. antennatus* CPUE's derivative series show quite a similar
232 pattern. The zones where the series have been forced to read zero coincide, and
233 almost any increase in the vorticity derivative corresponds with an increase in the
234 reversed catches derivative. The correlation between both series at lag zero, now
235 positive due to the sign change in the CPUE's derivatives mentioned, is slightly
236 larger than before, as expected, reaching a value of 0.48 (Fig. 3b). This result
237 strongly supports a relationship between the increases in the surrounding absolute
238 surface vorticity and decreases in the adult *A. antennatus* availability in the fishing
239 grounds.

240 The suggested mechanism explaining the relationship observed is now sup-
241 ported analyzing some surface vorticity episodes recorded when the mooring was
242 deployed in the area. During 2010, at least three of these episodes produced some
243 footprint in the instruments deployed in the mooring line.

244 The increase in the absolute value of the surface vorticity is commonly caused
 245 by the presence of an eddy, such as the one shown in Fig. 4. This particular eddy
 246 remained in the area between mid-November and mid-December 2010 and was
 247 studied in detail by Amores et al. [2013]. This eddy was clearly reflected in the
 248 currents registered at 500 and 900 m depth. A significant velocity increase was
 249 measured at both depths (episode 3 in Fig. 5a and 5b). Velocities showed spikes
 250 reaching up to 26 cm/s at 900 m, where the mean current during the whole year
 251 was computed to be around 5 cm/s. This eddy also affected the current direction,
 252 causing a complete reversal in the currents at 500 m (Fig. 5c) and a down slope
 253 gyre at 900 m (Fig. 5d).

254 This particular eddy clearly reached down to the bottom, and the recorded
 255 gyres and velocity increases could easily have caused the material resuspension.
 256 This hypothesis is supported by three indirect measurements:

- 257 1. the increase in the total flux mass (TFM) recorded by the moored sediment
 258 trap at the time of the eddy (Fig. 6)
- 259 2. the increase of the acoustic backscattering during the episode (Fig. 7a)
- 260 3. the clear down slope gyre at 900 m which could be related to a near bottom
 261 turbidity current (Fig 5d).

262 The eddy shown in Fig. 4 was not the only one recorded when the mooring was
 263 deployed. Another eddy, which occurred between mid-January and March 2010,
 264 was also measured by the mooring (episode 1 in Fig. 5). This eddy too reached
 265 the bottom, although with a weaker footprint in velocity. However, the TFM still
 266 reached similar values to those observed during the December 2010 eddy and an
 267 increase in backscattering is also observed.

268 Still another gyre was observed between June and July 2010; however, it was
 269 only noticeable at 500 m (episode 2 in Fig. 5). Its effect at 900 m was weak.
 270 Even so, a TFM peak was also measured by the sediment trap, although much
 271 weaker than during the other two eddies (Fig. 6). Backscattering did not show
 272 any significant increase during this episode. This could be explained by the steep
 273 slope of the area. Even if the eddy was not energetic enough to reach down to
 274 900 m depth, where the mooring was deployed, it could still affect the bottom
 275 at shallower depths and the resuspension of material could have reached deeper
 276 waters, causing the increased TFM that is recorded in the sediment trap.

277 From the above described episodes, it has been observed that the surface eddies
 278 exert some degree of influence on the seabed dynamics and that might increase
 279 the water turbidity near the bottom and affect the availability of *A. antennatus*
 280 in the fishing grounds. This mechanism can be better visualized by restricting the
 281 time interval to a shorter period of time of some of the data shown in Fig. 2, for
 282 the period, 2006-2010 (Fig. 8). During these years, the vorticity episodes are time
 283 spaced enough to allow an almost complete recovery towards a normal situation
 284 after any episode, before the arrival of the next one. The mechanism suggested
 285 are then better observed in the data. A vorticity increase (dark gray bands in
 286 Fig. 8) would trigger an increase in the re-suspended material and would force
 287 *A. antennatus* to move away from the fishing ground, probably towards greater
 288 depths, leading to a decrease in the catches of this species. This scenario would
 289 remain unchanged until the eddy effects disappear and the sediments once again
 290 precipitate to the sea floor (soft gray bands in Fig. 8). Once all the sediments
 291 are completely settled down, the water conditions would become suitable to allow
 292 the individuals to return to the depths where they can be caught (white bands in
 293 Fig. 8). Four episodes appear to follow one after another in the period shown.

294 Company et al. [2008] described a similar but stronger phenomenon in the
 295 submarine canyon system off the north-eastern Iberian Peninsula. They found a
 296 correlation between the strong currents associated with intense downslope shelf
 297 dense water current events and the disappearance of *A. antennatus* from its fish-
 298 ing grounds, exerting a negative effect on the catches reporting and a temporary
 299 collapse of its fishery. An increase in the mortality rated after exposure to high
 300 turbidity has also been detected for Penaeid shrimps at juvenile and adult stages
 301 [Lin et al., 1992]. Both downslope shelf dense water current events and mesoscale
 302 eddies have been reported to enhance organic-matter flux and deposition, increas-
 303 ing suspending particulate matter concentrations with the transport of organic
 304 matter from coastal zones to the deep ocean or by resuspension of bottom sed-
 305 iments [Bosley et al., 2004; Canals et al., 2006; Washburn et al., 1993]. Life
 306 forms as diverse as phytoplankton, protozoans, crustaceans, fish, sea snakes, ma-
 307 rine mammals and birds are found to alter their distributions in the presence of
 308 such flow patterns [Owen, 1981], which can be responsible for enhancing benthic
 309 productivity and biodiversity inside canyon habitats [Rowe et al., 1982; Schlacher
 310 et al., 2007; Vetter et al., 2010]. The presence of a significant amount of suspended
 311 sediment has also been related to the higher occurrence of juveniles and females

312 of the deep-water pandalid shrimp species, genus *Plesionika* [Puig et al., 2001]
 313 and the regions where the intermediate nepheloid layers detach from the seabed
 314 have been defined as potential deep-water nursery habitats for these species. The
 315 transportation of the particulate organic matter associated with the downslope
 316 shelf dense water current appears to be positive for recruitment of red shrimp
 317 in the north-eastern Iberian Peninsula [Company et al., 2008], with a positive
 318 increase in the landings 3-5 years after these events, preceded by an increase in
 319 the number of juveniles. However, this last effect has not been detected off the
 320 Balearic Islands, probably because of the slight difference in the feeding strate-
 321 gies of the species between the north-eastern Iberian Peninsula and the Balearic
 322 Islands. *A. antennatus* has a highly varied diet, being among the mega-benthic
 323 species mainly preying on the benthos in the deep Mediterranean [Cartes, 1994;
 324 Cartes and Carrassón, 2004]. However, benthic preys are particularly significant
 325 off the north-eastern Iberian Peninsula, where the submarine canyons enhance
 326 such types of food availability. Conversely, the trophic webs off the Balearic Is-
 327 lands show an impoverishment of the benthos biomass and depend more directly
 328 on food of planktonic origin, enhancing the consumption of micro-nektonic preys
 329 [Cartes et al., 2008; Maynou and Cartes, 2000]. In this sense, the positive effects
 330 of downslope shelf dense water current and sediment resuspension in the long
 331 term should be also more marked off north-eastern Iberian Peninsula than off the
 332 Balearic Islands.

333 Although the mechanism suggested is somehow speculative because we have to
 334 rely on indirect data (backscattering) to deduce bottom turbidity, we have shown
 335 several evidences that the presence of enough energetic eddies may cause bottom
 336 turbidity increases in the fishing ground. Still another indirect evidence supporting
 337 the mechanism suggested comes from the analysis of other demersal species caught
 338 in the same region that also appear to be related to vorticity changes in the area.
 339 The correlations found for each species (Fig. 9) are different from those observed
 340 for *A. antennatus* but consistent with the suggested mechanism where vorticity
 341 affects the seabed by increasing the bottom water turbidity.

342 Other decapod crustaceans, *Geryon longipes* and *Nephrops norvegicus*, the
 343 more sedentary and benthic species, show a significant positive correlation to the
 344 vorticity events at around 0.4 and 0.5, respectively. These two species are closely
 345 connected to the bottom, as reflected by their feeding behavior and biological char-
 346 acteristics. *G. longipes* preys on a broad range of benthic invertebrates [Cartes,

1993] and *N. norvegicus* shows a scavenging activity [Cristo and Cartes, 1998] and has been related to the sediment characteristics [Maynou and Sardà, 1997]. Unlike *A. antennatus*, it is likely that these two species may take some advantage of the re-suspended matter.

Galeus melastomus, which has more mobility than the previously considered epi-benthic species, feeds on the mesopelagic preys with the occasional occurrence of benthic feeding activity and scavenging in the adult phase [Fanelli et al., 2009]. This species showed a lower correlation (0.33) with the vorticity time series.

Finally, *Micromesistius poutassou* and *Phycis blennoides*, the two benthopelagic teleosts with greater capacity of movement above the bottom, are expected to be less affected by bottom water turbidity. In fact, no significant correlations between the CPUEs and vorticity have been found.

359 4 Summary and Conclusions

360 A reasonable good negative correlation is noted between the monthly CPUE of
361 the adult *A. antennatus* bottom trawl yields in the fishing grounds off northern
362 Mallorca and the mean surface vorticity in the surrounding area. We have shown
363 that the eddies causing the vorticity events may reach the bottom, increasing
364 the current velocities, which in turn would trigger sediment resuspension and
365 increased bottom water turbidity. Such a change in the water conditions would
366 force adult *A. antennatus* individuals to move away from the fishing ground,
367 probably downwards, to greater depths. This proposed mechanism is similar,
368 although lower in magnitude, to the one suggested by Company et al. [2008] in
369 the downslope shelf dense water current events of the submarine canyons in the
370 northern Catalan margin, off the north-eastern Iberian Peninsula. In the Balearic
371 Islands, where these geomorphological structures do not exist and where there is
372 no river runoff, the eddies would be the triggering factor.

373 Other deep water demersal species found along with the catch of the red shrimp
374 fishery, possessing different behavior and feeding habits, exhibit different responses
375 to these events, but all of them are consistent with the eddy generation near
376 bottom velocities increase and bottom water turbidity.

377 A final hypothesis could also be suggested from the results obtained. The
378 seasonal migration of most of the fishing fleet of Mallorca, targeting the red shrimp
379 in the Sóller fishing grounds during the summer has been explained by the highest
380 abundance of large spawning females in this area during this season [Guijarro
381 et al., 2008], similar to other areas off the north-eastern Iberian Peninsula [Sardà
382 et al., 1994, 1997]. In light of this, the absence of these large aggregations in
383 the Sóller fishing grounds during the rest of the year could be related to the
384 particular behavior of the species. However, it is worth noting that, according to
385 Amores et al. [2013], the vorticity episodes are much more intense off northern
386 Mallorca during the winter time (October to March) than in the summer. This
387 fact could be an additional factor explaining the decrease in the availability of
388 the red shrimp to fishing exploitation during these months. However, off southern
389 Mallorca, yields from the bottom trawl fishery targeting to red shrimp remain
390 more stable [Guijarro et al., 2008].

391 Acknowledgments

392 This research has been partially sponsored by the IDEADOS project (CMT2008-
393 04489-C03-01 and 03). The work of A. Amores has been funded by a JAE-PreDoc
394 grant from Consejo Superior de Investigaciones Científicas (CSIC) and co-funded
395 by Programa Operativo FSE 2007-2013. The authors are grateful to the fishermen
396 associations in Palma and the Autonomous Government of the Balearic Islands
397 for providing landing data. The authors also wish to express their gratitude to
398 the IDEADOS team for their collaboration in the data acquisition process and
399 the crews of the R/V Odón de Buen and F/V Punta des Vent for their valuable
400 assistance during the mooring deployment.

401 References

- 402 Amores, A., Monserrat, S., Marcos, M., 2013. Vertical structure and temporal
403 evolution of an anticyclonic eddy in the balearic sea (western mediterranean).
404 J. Geophys. Res. Oceans 118–4, 2097–2106.
- 405 Arrobas, I., Ribeiro-Cascalho, A., 1987. On the biology and fishery of *aristeus*
406 *antennatus* (risso, 1816) in the south portuguese coast. Inv. Pesq. 51 (suppl. 1),
407 233–243.
- 408 Beentjesa, M. P., Renwick, J. A., 2001. The relationship between red cod, *pseu-*
409 *dophycis bachus*, recruitment and environmental variables in new zealand. En-
410 vironmental Biology of Fishes 61 (3), 315–328.
- 411 Bombace, G., 1975. Considerazioni sulla distribuzione delle popolazioni di livello
412 batiale con particolare riferimento a quelle bentoniche. Pubbl. Staz. Zool. Napoli
413 39 (suppl 1), 7–21.
- 414 Bosley, K. L., Lavelle, J. W., Brodeur, R. D., Wakefield, W. W., Emmett, R. L.,
415 Baker, E. T., Rehmke, K. M., 2004. Biological and physical processes in and
416 around astoria submarine canyon, oregon, {USA}. Journal of Marine Systems
417 50 (1–2), 21 – 37.
- 418 Canals, M., Puig, P., de Madron, X. D., Heussner, S., Palanques, A., Fabres, J.,
419 2006. Flushing submarine canyons. Nature 444 (7117), 354–357.
- 420 Carbonell, A., Carbonell, M., Demestre, M., Grau, A., Monserrat, S., 1999. The
421 red shrimp *aristeus antennatus* (risso, 1816) fishery and biology in the balearic
422 islands, western mediterranean. Fisheries Research 44, 1–13.
- 423 Cartes, J. E., 1993. Diets of deep-sea brachyuran crabs in the western mediter-
424 ranean sea. Marine Biology 117, 449–457.
- 425 Cartes, J. E., 1994. Influence of depth and season on the diet of the deep-water
426 aristeid *aristeus antennatus* along the continental slope (400 to 2300 m) in the
427 catalan sea (western mediterranean). Marine Biology 120, 639–648.
- 428 Cartes, J. E., Carrassón, M., 2004. Influence of trophic variables on the depth-
429 range distributions and zonation rates of deep-sea megafauna: the case of the
430 western mediterranean assemblages. Deep Sea Research Part I: Oceanographic
431 Research Papers 51, 263–279.

- 432 Cartes, J. E., Papiol, V., Guijarro, B., 2008. The feeding and diet of the deep-sea
433 shrimp *aristeus antennatus* off the balearic islands (western mediterranean):
434 Influence of environmental factors and relationship with the biological cycle.
435 Progress in Oceanography 79, 37–54.
- 436 Company, J. B., Puig, P., Sardà, F., Palanques, A., Latasa, M., Scharek, R., 2008.
437 Climate influence on deep sea populations. PLoS ONE 3, e1431.
- 438 Cristo, M., Cartes, J. E., 1998. A comparative study of the feeding ecology of
439 nephrops norvegicus (l.), (decapoda: Nephropidae) in the bathyal mediter-
440 ranean and the adjacent atlantic. Scientia Marina 62 (Suppl. 1), 81–90.
- 441 Demestre, M., Martín, P., 1993. Optimum exploitation of a demersal resource
442 in the western mediterranean: the fishery of the deep-water shrimp *aristeus*
443 *antennatus* (risso, 1816). Scientia Marina 57 (2-3), 175–182.
- 444 Fanelli, E., Rey, J., Torres, P., de Sola, L. G., 2009. Feeding habits of black-
445 mouth catshark *galeus melastomus rafinesque*, 1810 and velvet belly lantern
446 shark *etmopterus spinax* (linnaeus, 1758) in the western mediterranean. J. Appl.
447 Ichthyol. 25 (Suppl. 1), 83–93.
- 448 Ghidalia, W., Bourgois, F., 1961. Influence of temperature and light on the dis-
449 tribution of shrimps in medium and great depths. Gfcm Studies and Reviews
450 16, 49.
- 451 Guijarro, B., Fanelli, E., Moranta, J., Cartes, J. E., Massutí, E., 2012. Small-scale
452 differences in the distribution and population dynamics of pandalid shrimps
453 in the western mediterranean in relation to environmental factors. Fisheries
454 Research 119-120, 33–47.
- 455 Guijarro, B., Massutí, E., 2006. Selectivity of diamond- and square-mesh co-
456 dends in the deepwater crustacean trawl fishery off the balearic islands (western
457 mediterranean). ICES J. Mar. Sci. 63 (1), 52–67.
- 458 Guijarro, B., Massutí, E., Moranta, J., Díaz, P., 2008. Population dynamics of the
459 red shrimp *aristeus antennatus* in the balearic islands (western mediterranean):
460 Short spatio-temporal differences and influence of environmental factors. Jour-
461 nal of Marine Systems 71 (3-4), 385–402.

- Hyder, P., Bigelow, K., Brainard, R., Seki, M., Firing, J., Flament, P., 2009. Migration and abundance of bigeye tuna (*Thunnus obesus*), and other pelagic species, inferred from catch rates and their relation to variations in the ocean environment. SOEST 09-02.
- Ivanov, V., Shapiro, G., Huthnance, J., Aleynik, D., Golovin, P., 2004. Cascades of dense water around the world ocean. Progress in Oceanography 60 (1), 47 – 98.
- Kai, E. T., Marsac, F., 2010. Influence of mesoscale eddies on spatial structuring of top predators' communities in the mozambique channel. Progress in Oceanography 86 (1-2), 214 – 223.
- Lin, H.-P., Charmantier, G., Thuét, P., Trilles, J.-P., 1992. Effects of turbidity on survival, osmoregulation and gill Na^+ - K^+ atpase in juvenile shrimp *Penaeus japonicus*. Marine Ecology Progress Series 90, 31–37.
URL <http://www.int-res.com/articles/meps/90/m090p031.pdf>
- Lohrmann, A., 2001. Monitoring sediment concentration with acoustic backscattering instruments. Nortek Tech. notes No. N4000–712.
- Massutí, E., Monserrat, S., Oliver, P., Moranta, J., López-Jurado, J. L., Marcos, M., Hidalgo, J. M., Guijarro, B., Carbonell, A., Pereda, P., 2008. The influence of oceanographic scenarios on the population dynamics of demersal resources in the western mediterranean: Hypothesis for hake and red shrimp off balearic islands. Journal of Marine Systems 71 (3-4), 421–438.
- Maynou, F., 2008. Environmental causes of the fluctuations of red shrimp (*aristeus antennatus*) landings in the catalan sea. Journal of Marine Systems 71 (3-4), 294–302.
- Maynou, F., Cartes, J. E., 2000. Community structure of bathyal decapod crustaceans off south-west balearic islands (western mediterranean): seasonality and regional patterns in zonation. Journal of the Marine Biological Association of the United Kingdom 80, 789–798.
- Maynou, F., Sardà, F., 1997. Nephrops norvegicus population and morphometrical characteristics in relation to substrate heterogeneity. Fisheries Research 30 (1-2), 139–149.

493 Moranta, J., Quetglas, A., Massutí, E., Guijarro, B., Hidalgo, M., Diaz, P., 2008.
494 Spatio-temporal variations in deep-sea demersal communities off the balearic
495 islands (western mediterranean). *Journal of Marine Systems* 71 (3-4), 346–366.

496 Owen, R. W., 1981. Fronts and eddies in the sea: mechanisms, interactions and
497 biological effects. *Analysis of marine ecosystems*, 197–233.
498 URL <http://swfsc.noaa.gov/publications/cr/1981/8149.pdf>

499 Palamara, L., Manderson, J., Kohut, J., Oliver, M. J., Gray, S., Goff, J., 2012.
500 Improving habitat models by incorporating pelagic measurements from coastal
501 ocean observatories. *Marine Ecology Progress Series* 447, 15–30.

502 Palmer, M., Quetglas, A., Guijarro, B., J.Moranta, Ordines, F., Massutí, E., 2009.
503 Performance of artificial neural networks and discriminant analysis in predicting
504 fishing tactics from multispecific fisheries. *Canadian Journal of Fisheries and*
505 *Aquatic Sciences* 66, 224–237.

506 Papaconstantinou, C., Kapisir, K., 2001. Distribution and population structure of
507 the red shrimp (*aristeus antennatus*) on an unexploited fishing ground in the
508 greek ionian sea. *Aquatic Living Resources* 14, 303–312.

509 Puig, P., Company, J. B., Sardà, F., Palanques, A., 2001. Responses of deep-water
510 shrimp populations to intermediate nepheloid layer detachments on the north-
511 western mediterranean continental margin. *Deep Sea Research Part I: Oceano-*
512 *graphic Research Papers* 48, 2195–2207.

513 Ramos, A. G., Santiago, J., Sangra, P., Canton, M., 1996. An application
514 of satellite-derived sea surface temperature data to the skipjack (*katsuwonus*
515 *pelamis linnaeus*, 1758) and albacore tuna (*thunnus alalunga bonaterre*, 1788)
516 fisheries in the north-east atlantic. *International Journal of Remote Sensing*
517 17 (4), 749–759.

518 Relini, G., Relini, L. O., 1987. The decline of red-shrimp stocks in the gulf of
519 genoa. *Investigacion Pesquera* 51, 245–260.

520 Relini, M., Maiorano, P., D’Onghia, G., Relini, L. O., Tursi, A., Panza, M., 2000.
521 A pilot experiment of tagging the deep shrimp *aristeus antennatus* (risso, 1816).
522 *Scientia Marina* 64, 357–361.

523 Rowe, G. T., Polloni, P. T., Haedrich, R. L., 1982. The deep-sea macrobenthos
524 on the continental margin of the northwest atlantic ocean. Deep Sea Research
525 Part A. Oceanographic Research Papers 29 (2), 257 – 278.

526 Sardà, F., Cartes, J. E., Norbis, W., 1994. Spatia-temporal structure of the deep-
527 water shrimp *aristeus antennatus* (decapoda: Aristeidae) population in the
528 western mediterranean. Fishery Bulletin 92, 599–607.

529 Sardà, F., Company, J. B., Bahamón, N., Rotllant, G., Flexas, M. M., Sánchez,
530 J. D., niga, D. Z., Coenjaerts, J., Orellana, D., Jordà, G., Puigdefàbregas,
531 J., Sánchez-Vidal, A., Calafat, A., Martín, D., Espino, M., 2009. Relationship
532 between environment and the occurrence of the deep-water rose shrimp *aristeus*
533 *antennatus* (risso, 1816) in the blanes submarine canyon (nw mediterranean).
534 Progress in Oceanography 82, 227–238.

535 Sardà, F., Company, J. B., Maynou, F., 2003. Deep-sea shrimp (*aristeus anten-*
536 *natus* risso 1816) in the catalan sea: a review and perspectives. J. Northw. Atl.
537 Fish. Sci 31, 117–136.

538 Sardà, F., D’Onghia, G., Politou, C. Y., Company, J. B., Maiorano, P., Kapisir,
539 K., 2004. Deep-sea distribution, biological and ecological aspects of *aristeus*
540 *antennatus* (risso, 1816) in the western and central mediterranean sea. Scientia
541 Marina 68, 117–127.

542 Sardà, F., Maynou, F., Talló, L., 1997. Seasonal and spatial mobility patterns
543 of rose shrimp *aristeus antennatus* in the western mediterranean: results of a
544 long-term study. Marine Ecology Progress Series 159, 133–141.

545 Schlacher, T. A., Schlacher-Hoenlinger, M. A., Williams, A., Althaus, F., Hooper,
546 J. N. A., Kloser, R., 2007. Richness and distribution of sponge megabenthos in
547 continental margin canyons off southeastern australia. Marine Ecology Progress
548 Series 340, 73–88.

549 Thomson, R. E., Davis, E. E., Heesemann, M., Villinger, H., 2010. Observations of
550 long-duration episodic bottom currents in the middle america trench: Evidence
551 for tidally initiated turbidity flow. J. Geophys. Res. 115, C10020.

552 Vetter, E. W., Smith, C. R., de Leo, F. C., 2010. Hawaiian hotspots: enhanced
553 megafaunal abundance and diversity in submarine canyons on the oceanic is-
554 lands of hawaii. Marine Ecology 31 (1), 183–199.

- 555 Washburn, L., Swenson, M. S., Largier, J. L., Kosro, P. M., Ramp, S. R., 1993.
556 Cross-shelf sediment transport by an anticyclonic eddy off northern california.
557 Science 261 (5128), 1560–1564.
- 558 Zainuddin, M., Kiyofuji, H., Saitoh, K., Saitoh, S.-I., 2006. Using multi-sensor
559 satellite remote sensing and catch data to detect ocean hot spots for albacore
560 (thunnus alalunga) in the northwestern north pacific. Deep Sea Research Part
561 II: Topical Studies in Oceanography 53 (3–4), 419 – 431.

Figure Captions

Fig. 1. Map of the studied area in the western Mediterranean. The unbroken line encloses Sóller fishing grounds where *Aristeus antennatus* is exploited and the broken line corresponds to the zone where the time series of vorticity has been calculated. Mooring location is indicated by a star.

Fig. 2. a) Monthly averaged time series of vorticity (black) and *Aristeus antennatus* CPUE's (green). b) Correlation between these two series, showing the maximum correlation (-0.35) around lag 0. Black line represents the 95% confidence level.

Fig. 3. a) Derivative time series of vorticity (black) and *Aristeus antennatus* CPUE's (green). In the series of the vorticity derivative, the negative values have been fixed at 0, while the positive values of the derivative CPUE's series have been set at 0. Notice that the last one has suffered a change of sign. b) Correlation between these two series show the maximum value (0.48) at lag 0. Black line represents the 95% confidence level.

Fig. 4. Sea Surface Height (SSH) image from December 1, 2010. It shows an eddy in the region analyzed. The star shows the mooring position.

Fig. 5. 24h low-pass filtered speed series of 500 (a) and 900 (b) m depth current meters of the mooring for the whole recorded period. Blue indicates the low speed values degrading to red, which indicates the high values. (c) and (d) are the progressive vector diagrams for 500 and 900 m depth, respectively. The different colors coincide temporally with the speed time series. Enclosed areas represent moments where an eddy is present in the zone. Ellipse number 1 highlight an eddy which is strongly present at 500 m depth, but weakly present at 900 m; number 2 ellipse shows an eddy which reached to 500 m depth, although not right up to 900 m; and number 3 ellipse illustrates an eddy which arrived strongly to the 500 and 900 m depths. Note that in the PVDs the ratio between the scales of x and y axis is 2:1.

Fig. 6. Total Flux Mass (TFM) collected by the sediment trap during the whole

597 sampling time. The gap in the data is due to the unavailability of ship for carrying
598 out the mooring maintenance. The dashed ellipses show the increment of TFM
599 due to the eddies reported in Fig. 5.

600

601 Fig. 7. Acoustic backscattering (a) and speed (b) measured by the 900 m current
602 meter during the third episode.

603

604 Fig. 8. Zoom of the Fig. 2, where the effect of the vorticity (blue) on the *Aristeus*
605 *antennatus* CPUE's (green) can be seen. The colored bands indicate the amount
606 of particles that would be re-suspended.

607

608 Fig. 9. Time series of CPUE's for five demersal species (by catch) from the deep
609 water trawl fishery and its correlation with the absolute value of surface vorticity.

610 Figures

Figure 1: Map of the studied area in the western Mediterranean. The unbroken line encloses Sóller fishing grounds where *Aristeus antennatus* is exploited and the broken line corresponds to the zone where the time series of vorticity has been calculated. Mooring location is indicated by a star.

Figure 2: a) Monthly averaged time series of vorticity (black) and *Aristeus antennatus* CPUE's (green). b) Correlation between these two series, showing the maximum correlation (-0.35) around lag 0. Black line represents the 95% confidence level.

Figure 3: a) Derivative time series of vorticity (black) and *Aristeus antennatus* CPUE's (green). In the series of the vorticity derivative, the negative values have been fixed at 0, while the positive values of the derivative CPUE's series have been set at 0. Notice that the last one has suffered a change of sign. b) Correlation between these two series show the maximum value (0.48) at lag 0. Black line represents the 95% confidence level.

Figure 4: Sea Surface Height (SSH) image from December 1, 2010. It shows an eddy in the region analyzed. The star shows the mooring position.

Figure 5: 24h low-pass filtered speed series of 500 (a) and 900 (b) m depth current meters of the mooring for the whole recorded period. Blue indicates the low speed values degrading to red, which indicates the high values. (c) and (d) are the progressive vector diagrams for 500 and 900 m depth, respectively. The different colors coincide temporally with the speed time series. Enclosed areas represent moments where an eddy is present in the zone. Ellipse number 1 highlight an eddy which is strongly present at 500 m depth, but weakly present at 900 m; number 2 ellipse shows an eddy which reached to 500 m depth, although not right up to 900 m; and number 3 ellipse illustrates an eddy which arrived strongly to the 500 and 900 m depths. Note that in the PVDs the ratio between the scales of x and y axis is 2:1.

Figure 6: Total Flux Mass (TFM) collected by the sediment trap during the whole sampling time. The gap in the data is due to the unavailability of ship for carrying out the mooring maintenance. The dashed ellipses show the increment of TFM due to the eddies reported in Fig. 5.

Figure 7: Acoustic backscattering (a) and speed (b) measured by the 900 m current meter during the third episode.

Figure 8: Zoom of the Fig. 2, where the effect of the vorticity (blue) on the *Aristeus antennatus* CPUE's (green) can be seen. The colored bands indicate the amount of particles that would be re-suspended.

Figure 9: Time series of CPUE's for five demersal species (by catch) from the deep water trawl fishery and its correlation with the absolute value of surface vorticity.