1 Date of preparation: October 4, 2001 (1st manuscript)

2 July 2, 2002 (revised manuscript)

3 December 12, 2002 (2nd revision)

4 January 20, 2003 (Accepted)

5 Number of text pages: 28 (including references)

6 Number of tables: 7

7 Number of figures: 2

9 Phosphorus fractions and dynamics in surface earthworm casts under
10 native and improved grasslands in a Colombian savanna Oxisol[¶]

11
12 Juan J. Jiménez^{a*}, Alex Cepeda^a, Thibaud Decaëns^b, Astrid Oberson^c, Dennis K. Friesen^d

13
14 ^a *Soil and Plant Nutrition Unit. CIAT. P.O. Box 6713, Cali, Colombia*

15 ^b *Laboratoire d'Ecologie. UPRES-EA 1293. UFR Sciences. Université de Rouen. F-76821 Mont Saint Aignan*
16 *Cedex, France*

17 ^c *ETH, Institute of Plant Sciences. Eschikon, 33. CH-8315. Lindau, Switzerland*

18 ^d *CIMMYT-IFDC, P.O. Box 25171, Nairobi, Kenya*

19
20 * Author for correspondence

21
22 Current address (until May 2003):

23 Land and Water Development Division (AGLL), Room B-701

24 FAO

25 Viale Terme di Caracalla – 00100 Rome, Italy.

26 E-mail: juan.jimenez@fao.org

27
28 [¶] This paper is dedicated to Paulo César Cepeda Virviescas *in memoriam*.

29 **Abstract**

30 The objective of this study was to assess the effect of a native anecic species on phosphorus
31 availability in an Oxisol characterised by a low chemical fertility. Experiments were carried
32 out at Carimagua research station in a representative site of the isohyperthermic savannas on

the Colombian Orinoco basin. One field study and two laboratory/incubation studies were performed in a natural herbaceous savanna and a Brachiaria decumbens and Pueraria phaseoloides pasture. In the laboratory, experiment pots were prepared containing soil collected from the respective field paddock's topsoil. Total P content was higher in earthworm casts than in the surrounding soil in field samples, 50% in native savanna soil and more than 100% in pasture soil. In casts produced under laboratory conditions this increase was relatively low (10–20%). Under field conditions, almost without exception, all P fractions were increased in casts relative to the original soil (corresponding to the increase in total P content), being relatively greater in the labile inorganic P fractions. In addition, samples from the natural savanna showed that pH of casts was higher (5.2) than that of soil (4.6) in both field and laboratory samples. Except in the native savanna under field conditions, the phosphatase activity was reduced in casts by 16.7 to 44%. From our results we conclude that earthworms in the field incorporate P from litter or other organic sources (i.e. undecomposed plant and root material, earthworm faeces) which is not normally measured in the analysis of bulk soil.

Keywords: Earthworms; Phosphorus; Earthworm casts; Ecosystem engineers; Savanna; Oxisols; Grasslands.

1. Introduction

At present there is increasing evidence that soil macroinvertebrates improve soil fertility due to their role in soil organic matter transformations and nutrient dynamics at different spatial and temporal scales, which may improve nutrient uptake by plants (Lavelle, 1997). A few of these invertebrates have been defined as the “soil ecosystem engineers” (Stork and Eggleton, 1992; Jones et al., 1994; Lavelle, 1997). By definition, ecosystem engineers are “those

organisms that directly or indirectly modulate the availability of resources to other species by causing physical state changes in biotic or abiotic materials” (Jones et al., 1994). This means that they are capable of regulating the trophic (organic matter) and spatial distribution (habitat availability) of resources in the soil through the production of physical bio-structures (e.g. casts, galleries and nests). Soil invertebrates are major determinants of soil processes, especially in tropical ecosystems (Lavelle, 2000).

Earthworms belong to this functional group because through their burrowing activities, mixing soil with litter and egesting casts inside the soil or at the soil surface, they affect the physical properties of soils, nutrient cycling and plant dynamics (Lal, 1991; Thompson et al., 1993; Lavelle, 1997). To assess the contribution of these organisms to soil processes and ecosystem function, it is necessary to describe the phenomena that occur in the casts (Martin and Marinissen, 1993). Yet the impact on nutrient cycling has not been investigated in detail in tropical anecic earthworms (Decaëns et al., 1999b), even though they produce a substantial amount of casts in the soil surface. For example, some studies have revealed higher contents of available phosphorous (P) (that can be taken up by plants) in earthworm casts than in the control soil (Lunt and Jacobson, 1944; Nye, 1955; Lal, 1974; Krishnamoorthy, 1990; Guggenberger et al., 1996). Effects of earthworms on P are especially interesting since part of the pool, which is normally adsorbed onto the soil solid phase, may be desorbed after gut transit (López-Hernández et al., 1993; Brossard et al., 1996). These organisms have a marked impact on mineralisation of P^{δ} (Sharpley and Syers, 1976; James, 1991; López-Hernández et al., 1993; Chapuis and Brossard, 1995; Brossard et al., 1996), and are able to increase its availability for plants in their casts. This process has been widely documented for both tropical and temperate species (Sharpley and Syers, 1976;

^δ Phosphorus mineralization is an enzymatic process and a group of phosphatases are involved in the catalysis and release of phosphate from organic P compounds to the soil solution (Mullen 1998).

Barois et al., 1987; Lavelle and Martin, 1992; Lavelle et al., 1992; López-Hernández et al., 1993; Scheu, 1987). It is the result of their highly efficient digestive system while they excrete intestinal and cutaneous mucus that leaves nutrients in excess. As a consequence, earthworms have an important role in nutrient availability and cycling in natural and agricultural ecosystems (Coleman et al., 1994; Buse, 1990; Marinissen and de Ruiter, 1993; Bohlen et al., 1997; Decaëns et al., 1999b).

About 75% of soils in Neotropical savannas are strongly weathered, acidic and infertile belonging to the order of Oxisols (USDA Soil Taxonomy 1978). Low total and available P contents and high P fixation capacity due to high contents of Fe and Al oxides characterise these soils. Only 20% of total fertilizer P applied to soils in these agroecosystems is recovered by the crop to which it is applied (Friesen et al., 1997). The remainder is gradually rendered less available to succeeding crops by processes that slowly move P into more stable inorganic and organic pools in the soil. One strategy to increase the productivity and sustainability of production in such agroecosystems is to increase P recovery from these less accessible forms using crop and forage cultivars that are more efficient in acquiring P and cycling it into pools more available to crops. In addition, the role of soil macroinvertebrates in P cycling must be considered for their potential and availability of soil P for plant uptake, since the activities of soil organisms may preserve nutrients in the biostructures they produce and hence reduce the availability to plants in short temporal scales.

The potential of the Colombian savannas, an isohyperthermic ecosystem dominated by Oxisols, for both crop and livestock production systems, is limited by the lack of available P for primary production (Thomas et al., 1995). The conversion of native savanna into intensive pastures generally leads to a huge increase in earthworm biomass (Decaëns et al., 1994, Jiménez et al., 1998b). Martiodrilus carimaguensis (Glossoscolecidae) is a large anecic

native savanna earthworm from the Colombian “Llanos” (Jiménez et al., 1998a, b) which has been shown to affect soil processes (Decaëns, 2000; Decaëns et al., 1999a, b) and there is potential to take advantage of its activities in tropical agroecosystems (Mariani, 2001).

Casts of large anecic M. carimaguensis are enriched in labile organic P, which suggests that this species improves the supply of P in soil under pastures by creating an available organic P pool (Guggenberger et al., 1996). Oberson et al. (1999) presented evidence from soil P fractionation analyses which indicated that a high level of P cycling in intensive pastures contributed to their sustainability with very low P inputs. Based on these observations, we hypothesized that M. carimaguensis, through greatly increased population density and their effect on soil P dynamics, is a major contributor to the sustainable productivity of intensive pastures on Colombian savannas Oxisols. The objective of this study was to determine the temporal dynamics of P fractions in casts and quantify P availability and enhanced P cycling in an Oxisol of the Colombian Llanos. The study was carried out in both laboratory and field experiments to assess the impact of this species of high casting surface activity (Jiménez et al., 1998a, b; Decaëns et al., 1999b) in nutrient cycling. Since earthworms can accelerate the mineralisation of organic matter by reducing the size of residues to particles more available to microbes (Swift et al., 1979; James, 1991) one laboratory experiment examined the effect of incorporating finely ground vegetative material on P mineralization.

2. Materials and methods

2.1. Study site

This study was conducted during May-September 1994 at the CIAT-CORPOICA experimental station at Carimagua (4°30'N; 71°19'W; 150 m above sea level) located 320 km east of Villavicencio in the Eastern Plains of Colombia. The site is representative of the

well-drained isohyperthermic savanna ecosystem. Mean annual rainfall and temperature are 2,240 mm and 26.5 °C respectively. Rainfall distribution is characterized by a 4-month sharp dry season from December to March. Native vegetation varies with topography accordingly: open herbaceous savannas in the uplands (“altos”), and gallery forests or flooding savannas in the low-lying areas (“bajos”). Samplings were done in a well-drained silt-clay Oxisol (typic haplustox, fine kaolinitic, isohyperthermic). They are characterised by favourable physical properties, e.g. porosity and water retention, but high Al saturation (>80%) and low chemical fertility (Table 1).

2.2. Background ecology of M. carimaguensis

Martiodrilus carimaguensis is a large native dorsally pigmented earthworm quite common in the savannas of Carimagua (Colombia). The dimensions for adults are 9.3 mm in diameter, 194 mm in length and weighs 9.2 g (in 4% formalin) on average (Jiménez et al., 1998a). Its morphology and life history traits are like most anecic species (*sensu* Bouché, 1977); however, it only eats litter in an opportunistic way and its feeding regime is mostly based on small casts egested by other earthworm species (Mariani et al., 2001). The establishment of intensive pastures results in a spectacular increase of density compared to the native savanna, 18.2 and 0.2 ind m⁻², respectively (Jiménez et al., 1998b). The main aspects of their biology, ecology and adaptive strategies can be consulted in Jiménez and Decaëns (2000) and Jiménez et al. (1999; 2000).

2.3. Casts of M. carimaguensis

Martiodrilus carimaguensis produces a substantial amount of tower-like casts at the soil surface. Their size range from 3 to 6 cm in diameter, 2 to 10 cm in height and 25 g dry weight on average (Decaëns, 2000). As they are deposited the casts are a pasty structure, and within a period of days, the final structure is a combination of dry and fresh material at the bottom and the top of the cast, respectively. Whether the earthworm leaves its semi-permanent burrow or descends deep in the soil to begin diapause (Jiménez et al., 2000) the cast dries completely, and may remain at the soil surface for more than one year (Decaëns, 2000). The disappearance of casts from the soil surface are both of environmental and anthropogenic origin, i.e. rainfall impacts, fire, vegetation cover, cattle trampling and the burrowing by small invertebrates (Decaëns, 2000; Mariani et al., in press).

2.4. Experimental design

Three experiments were performed during the rainy season of 1994: one field study and two laboratory/incubation studies using soils from the same plots (paddocks) where the fieldwork was conducted.

2.4.1. Field experiment

The field experiment was carried out from May to August in two nearby land use systems (paddocks). They were a 2.3 ha. herbaceous native savanna dominated by Andropogon bicornis L. and Trachypogon vestitus Anders. that was burnt the previous year, and a 18 yr-old 2-ha. grazed pasture of Brachiaria decumbens cv Basilisk and Pueraria phaseoloides (Roxb.) Benth. CIAT 9900 (Kudzu). The pasture was grazed by 1 cattle ha⁻¹ during the dry season and 2 cattle ha⁻¹ during the rainy period.

1 A randomized block-design was used. In each paddock, nine areas of 4x4 and 1x1 m²,
2 respectively, were randomly allocated and, in the case of the pasture, these were protected
3 with wire exclusion cages to avoid cattle trampling and physical disturbance of surface casts.
4 The difference in the size of these areas was due to the fact that outstanding differences
5 regarding earthworm numbers existed and therefore in the number of surface casts (Jiménez
6 et al., 1998a). In each land use system (paddock) the nine sampling areas were randomly
7 grouped in clusters containing three areas. The surface casts were sampled from each area
8 and included within each group, so that there were three replicates for each sampling date.
9 Hence, a replicate was considered as the sum of casts collected from the three randomly
10 selected areas.

11 The surface casts that were already present in the sampling areas were removed the day
12 before the experiment started, and on the following day, fresh surface casts recently
13 deposited were identified, and displaced aside the earthworm gallery. Casts were tagged with
14 a plastic peg to identify the time of *in situ* incubation that corresponded to six different ages,
15 i.e. 1, 4, 8, 16, 32 and 64 days after they were egested in the soil surface. In order to ensure
16 enough cast material for a given incubation time, casts were picked up, placed in an ice chest
17 and carried to the laboratory. 70 g of fresh casts were collected approximately for each
18 group, of which 40 g were dried under forced draft in an oven at 40 °C and 30 g were stored
19 in a refrigerator at 4 °C prior to analysis as described below.

20 Control soil samples were taken in each paddock by splitting it into four areas where five 0-
21 15 cm soil cores were taken per area and mixed. Samples were further prepared for analyses
22 by crushing and sieving through a 2-mm mesh. Samples for microbial P determinations were
23 maintained fresh under refrigeration as mentioned above.

2.4.2. Laboratory experiments

Two laboratory experiments were conducted. In the first (LE1), no additional treatment other than the soil from the paddocks was applied. In the second (LE2), two additional treatments were applied in factorial combination with the two field treatments, e.g. green vegetative material collected from the same land use systems was added to one of the treatments. In order to test the effect of plant material addition on P dynamics the vegetative material was oven-dried, finely ground and mixed with the corresponding soil at a rate of 20 g kg⁻¹ soil. A first set of twelve pots were prepared using 6 kg of soil (air-dried and sieved to 2 mm) collected from the top 15 cm of the two land use systems where the fieldwork was conducted, although not in the same replicate areas used in the field study. These twelve pots corresponded to the land use systems studied, savanna and introduced pasture, two treatments (with and without organic amendments) and three replicates. The moisture content of the soil was adjusted to pF 2 (25% H₂O v/v) in each pot 5 days prior to earthworm introduction. Twelve adults of M. carimaguensis were collected in the field and placed in each pot for a 6-day period of conditioning to void field-ingested material. Afterwards, earthworms were transferred directly to another set of pots [12 units that corresponded to land use system (2), addition of plant material (2) and three replicates] containing 2-kg of the same sieved and similarly preconditioned soil. These pots were used only to collect cast material. After 1 day, these second pots were examined, casts retrieved and earthworms from each pot were moved to another set of pots prepared in an identical manner to ensure that casts belonged to the same experimental unit. This procedure was repeated six times to complete the temporal lag, i.e. 1, 4, 8, 16, 32, 64 days of cast incubation and to ensure the same conditions for cast sorting. Surface cast material from the pots was placed in Petri dishes containing moistened filter paper in order to maintain the humidity of casts. These

were incubated at ambient temperature (24 ± 3 °C) in the laboratory for the same time periods, after which they were broken up, mixed and separated into two other samples (one air dried, and the other moist) as described above for the field experiment. Control (non-ingested) soil was sampled from pots when earthworms were introduced and treated in the same manner as the incubated casts.

2.5. Determination of soil P fractions in casts and controls

Phosphorous in soil and earthworm casts were fractionated according to a modified method of Hedley et al. (1982) using successively the following increasingly aggressive extractants: H₂O with anion exchange resin in HCO₃⁻ form, 0.5 M NaHCO₃, 0.1 M NaOH, 1 M HCl, and hot concentrated HCl (Tiessen and Moir, 1993). Inorganic P in extracts was determined by the molybdate-ascorbic acid method (Murphy and Riley, 1962). Total P in the H₂O, NaHCO₃ and NaOH was measured after digestion with K₂S₂O₈ (Bowman, 1988). Total soil P was determined by perchloric acid digestion (Olsen and Sommers, 1982). Controls and one day-old casts were fractionated according to the full method while a reduced method using the first three extractants was applied to the remaining samples.

Inorganic P removed by anion exchange resin comes either from solution or is desorbed from the Al and Fe oxyhydroxide colloid surfaces in the soil (Mattlingly, 1975). Sodium bicarbonate (0.5 M at pH 8.5) also extracts weakly adsorbed P_i (Hedley et al., 1982). Together these two P_i fractions constitute a highly available P_i pool in soil. More slowly available P_i (also called “secondary P_i”) is extracted by 0.5 M NaOH and is associated with amorphous and crystalline Fe and Al oxyhydroxides. Highly labile soluble organic P compounds are found in the water phase of the resin-H₂O extractant. The weakly alkaline NaHCO₃ extractant also removes easily hydrolyzable organic P (P_o) compounds such as

ribonucleic acids and glycerophosphate (Bowman and Cole, 1978) while the more strongly alkaline NaOH solution extracts less labile P_o associated with fulvic and humic acids. Dilute HCl (0.1 M) extracts P_i from apatite or octocalcium phosphate (Frossard et al., 1995), neither of which are likely to be present in Oxisols unless they have been fertilized with phosphate rocks. Hot concentrated HCl extracts more stable pools of P_i and P_o , some of which may be associated with particulate organic matter (Tiessen and Moir, 1993). The P remaining in the sample (residual P_t) contains very recalcitrant P_i and P_o forms that likely participate in P cycling processes only at long term.

Phosphorus in the microbial biomass and acid soil phosphatase activity were estimated on moist samples using procedures described by Oberson et al. (1999) and Tabatabai (1982), respectively. This procedure was used since air drying of cast and soil samples may cause loss of inorganic P from microbial biomass and give erroneous results (Sparling et al., 1985; James, 1991).

Total carbon and nitrogen were analyzed on previously 2 mm sieved subsamples. A LECO CR-12 furnace with CO_2 infrared detection was used to determine total C, and the standard Kjeldahl digestion to measure total N contents (Krom, 1980). A titration method was used to extract exchangeable Al and H using 1 M KCl. Cations were extracted with 1 M NH_4 -acetate and determined by atomic absorption spectrometry using standard methods.

2.6. Statistical analysis

The Kolmogorov-Smirnov test (Lilliefors, 1967) was used to test the normality of data distribution, and transform data before analysis if necessary. A two-way analysis of variance (ANOVA) was performed with land use (savanna / pasture) and sample origin (cast vs. soil) as the fixed main effects for phosphatase activity, microbial P, total P, total C, Bray-II P and

pH (water, 1:1). To analyse the differences on the P fractions extracted with H₂O/resin, bicarbonate and NaOH solutions for each sampling date a two-way ANOVA with again land use and cast age as the fixed main effects was used. The Bonferroni procedure for multiple nested tests was applied (Cooper, 1968) to adjust the significance probability levels. The adjusted 0.05, 0.01 and 0.001 levels were, respectively: 0.004 [= 0.05 / (2 x 6)], 0.0008 [= 0.01 / (2 x 6)], and 8.3×10^{-5} [= 0.001 / (2 x 6)]. Additional comparisons of means were performed with the Tukey HSD test for all variables when the F-test was significant ($P < 0.05$). The software package Statistica 5.1 for Windows (© Statsoft Inc. 1996) was used in all statistical analyses.

3. Results

3.1. Phosphatase activity, microbial P, total P, and other chemical properties

Phosphatase activity in soils and casts ranged from 120 to 313 mg-nitrophenol kg⁻¹ h⁻¹ in laboratory experiment samples, and from 249 to 312 mg-nitrophenol kg⁻¹ h⁻¹ in the *in-situ* (field collected) pasture and savanna samples, respectively. Except for *in-situ* samples from the native savanna phosphatase activity was significantly lower in casts than in the control soil (Table 2), probably due to the manipulation of soil or because enzymes were partly degraded during gut transit or because phosphates production was decreased due to higher availability of P in casts. In field samples, phosphatase activity was higher in earthworm casts than in soil in the native savanna system but lower in the intensive pasture, although these differences were not significant the interaction of fixed main effects was significant (ANOVA, $F = 54.72$; $P < 0.001^{***}$) (Table 3). Microbial biomass P was significantly higher in 1-day-old casts than in the corresponding control soil from the savanna and from the

pasture (Table 3). Values of pH were significantly higher in earthworm casts than in the control soil in both systems and for both field and laboratory produced samples.

Total P content was significantly higher in 1-day old earthworm casts than in control soil for both laboratory incubated and field aged samples (Tables 2 and 3). Total P calculated from the sum of P fractions did not differ significantly from that determined directly by perchloric acid digestion for any particular treatment (not shown). In the laboratory experiments, casts contained 10-20 mg-P kg⁻¹ soil or 5-10% more total P than the soil from which they were produced. Casts produced in the field had approximately 1.5 and 2 fold more P than the bulk soil in the native savanna and the introduced pasture, respectively (Table 2).

Bray-II P and total C concentrations were significantly higher in 1-day-old casts than in the corresponding control soil for field samples in both systems (Table 3), while no significant differences were observed for the laboratory samples (Table 2).

3.2. P dynamics

The effect of M. carimaguensis on movement of P among soil P pools is considered in two phases, the first being the immediate effect of transit through the earthworm's gut and the second being the subsequent temporal changes in P concentration in pools as the deposited cast is aged *in situ* in the field or incubated in the laboratory.

The immediate effects of casting on the concentration of P in soil P pools in the field and laboratory are shown in Tables 4 and 5, respectively. Under field conditions, almost without exception P fractions were larger in casts than in the bulk soil (corresponding to the increase in total P content). Increases ranged from 0% (HClhc-P_o) to 875% (Resin- P_i) in the egested savanna soil and 46% (Residue Pt) to 814% (Resin- P_i) in the egested pasture soil, and were relatively greater in the labile P_i fractions (resin- P_i and NaHCO₃- P_i) than in the other P

fractions. In both savanna and pasture derived casts, about 60% of the increased P content was found in P_i fractions, 30% in P_o fractions and 10% in the residual P_t fraction. Most of the total P increase in cast over soil was found in secondary (NaOH) P_i and P_o pools in both savanna and pasture derived casts, with significant amounts entering stable P pools as well, especially in the pasture casts.

Under laboratory conditions where the increase in total P content due to casting (11-17%) was much smaller than in the field, increases in the sizes of P fractions ranged from 0% to 344% (Table 5). In the savanna soil casts, most of the added P was found in the secondary P pools (20% NaOH- P_i and 64% NaOH- P_o) whereas substantial amounts were also found in the stable P and residual P_t pools in the pasture soil casts. The addition of green material residues had no significant effect on either P_i or P_o fractions in the laboratory experiment, neither in the soil nor in the casts.

The dynamics of labile pools of P in ageing casts is shown in Figures 1 and 2 for laboratory and field incubation experiments, respectively. The statistical significance of the fixed main effects and their interaction is shown in tables 6 and 7 for casts incubated in the field and in the laboratory, respectively. In most cases the interaction of cast age and land use was not significant, except for the P_i extracted by NaOH and bicarbonate in casts incubated in the field. P_i extracted by resin was increased strongly in fresh casts but then slowly declined to the levels in soil over the following 64 days of incubation. In contrast, organic P extracted by bicarbonate and hydroxide increased 1 to 8 days after casting, rather than during transit of the earthworm gut, and then remained relatively constant over the remaining 56 days of the incubation. Inorganic P in bicarbonate and hydroxide was not affected significantly by casting and did not change significantly with time of incubation.

Similar patterns in P dynamics in P_i and P_o fractions were observed during *in situ* ageing of M. carimaguensis casts produced in the field although of much lower magnitude, especially for P_o fractions. The most marked changes occurred during transit of the earthworm gut whereas, after the initial increase, P_i and P_o in all fractions remained relatively constant over the 64 days of field incubation.

4. Discussion

4.1. P accumulation and mineralization in casts

The use of both field and laboratory experiments in this study helps provide a better understanding of the actual dynamics of nutrients in soil and in biogenic structures produced by soil macroinvertebrates. Although the total P contents of control soils in the field and laboratory experiments were similar, ingestion by M. carimaguensis increased total P by 53% to 114% under field conditions but by <20% under laboratory conditions. Under field conditions, M. carimaguensis apparently incorporated P from sources such as litter, undecomposed plant debris and roots and other holorganic casts that were not present in the soil under laboratory conditions. Together with mixing of soil and litter, both coprophagy and necro-rhizophagy seem to be the dominant features of the M. carimaguensis diet (Mariani et al., 2001).

Phosphorus incorporated into soil from “non-soil” sources may enter into all P fractions (organic and inorganic) but the fractionation data indicate that a greater proportion entered into labile P_i fractions, particularly under field conditions (Tables 4 and 5). It is not possible to conclude that M. carimaguensis promoted the mineralization of organic P since the proportion of P fractions in the substrate was unknown. However, the fact that under laboratory conditions, where organic litter substrate was not provided (LE1) and where the

1 increase in total P due to casting was quite small, relatively large increases especially in the
 2 secondary (NaOH- P_i and $-P_o$) pool sizes accompanied by decreases in stable fractions
 3 suggests that ingestion of the soil by the earthworm did promote movement of P from stable
 4 to more labile P forms. Moreover, the relatively large increases observed in resin- P_i in the
 5 laboratory experiment, and in all labile fractions (H_2O-P_o , resin- P_i , bic- P_i and bic- P_o) in the
 6 field experiment (all small pools), suggests some mineralization of less available P from
 7 large more stable pools where relative changes would be difficult to detect. This would agree
 8 with the interpretation that M. carimaguensis likely behaves as an endogeic (soil consumer)
 9 rather than an anecic (soil + litter consumer) species in terms of feeding regimes (Mariani et
 10 al., 2001). Since the addition of green material residues had no significant effect on either P_i
 11 or P_o fractions in the casts produced in the laboratory experiment, the hypothesis that M.
 12 carimaguensis ingests faeces from other earthworms (Mariani et al., 2001) or even may feed
 13 on their own casts (Jiménez et al., 1998b) is strengthened.

14 Although beginning with almost identical total soil P contents, casts produced *in situ* in the
 15 B. decumbens and P. phaseoloides pasture had more than twice the total P content and
 16 correspondingly higher P content in all fractions than casts produced from savanna soil
 17 (Table 4). This was probably due to higher biomass production, both aboveground and roots,
 18 of legume and deep-rooting grasses in the pasture resulting in greater litter fall and root
 19 turnover (Thomas, 1992; 1995; Thomas et al., 1992; Oberson et al., 1995; Fisher et al.,
 20 1994) as well as dung depositions from cattle.

21 Our study confirmed the results obtained by several authors who reported higher P contents
 22 in earthworm casts than in the surrounding soil from grassland ecosystems (Sharpley and
 23 Syers, 1976; Barois and Lavelle, 1986; Barois et al., 1987; James, 1991; López-Hernández et
 24 al., 1993; Brossard et al., 1996; Scheu, 1987). Barois et al. (1999) reported a significant
 25 increase of available P after transit through the earthworm gut, with the largest differences

for water extractable P, which was doubled after ingestion. Phosphorus contents were at least 30% higher in casts of several earthworm species than in the soil. In our study, water extractable P was 300% higher in casts than in the adjacent control soil under field conditions. Stabilization of P in casts of M. carimaguensis occurred between 16 and 64 days after cast deposition, whilst this lag was of 4 days in casts of the endogeic P. corethrurus (López-Hernández et al., 1993).

Sharpley and Syers (1976) estimated that the total amount of organic-P accumulated in 30 tons ha⁻¹ yr⁻¹ of surface earthworm casts in permanent pastures from a New Zealand watershed was 11-14 kg ha⁻¹ yr⁻¹. Estimated surface fresh cast production by M. carimaguensis at the study site was, respectively, 1.2 t ha⁻¹ yr⁻¹ and 13.2 t ha⁻¹ yr⁻¹ in the savanna and in the pasture, based on a density of 0.2 fresh casts m⁻² and 2.2 fresh casts m⁻² in the respective systems and taken into account the active period of the species, at least 4 months (Jiménez, 1999). The average dry weight of casts ranges from 25 g (Decaëns, 2000) to 35 g (Jiménez, unpubl. data). Thus, 0.36 kg ha⁻¹ yr⁻¹ and 5.61 kg ha⁻¹ yr⁻¹ of total P may be accumulated in fresh casts of M. carimaguensis, respectively, in the savanna and the pasture (0.13 and 1.8 kg ha⁻¹ yr⁻¹ of total P_o). This represents an important contribution to the overall P fluxes in these agroecosystems. For example, the total P uptake in above-ground biomass of a grass-legume pasture (B. humidicola plus several legumes), a maize monocrop and the native savanna was found to be 14, 18 and 4 kg P ha⁻¹ yr⁻¹ at the same site (Friesen et al., 1997). Hence, total P accumulated in casts of M. carimaguensis is ≈40% of total annual P uptake by grasses in the pasture and only ≈9% of the total P uptake by above-ground vegetation in the savanna. Nonetheless, as Decaëns et al. (1999b) showed for N dynamics in casts of the same species, the global contribution of the whole earthworm community to availability of P for plants may be even higher due to the presence of other earthworm

species in the soil and the casts deposited in the soil (Jiménez et al. 1998a). Thus, our results confirm those of Guggenberger et al. (1996) who showed that earthworm activity at Carimagua resulted in a marked effect on P availability.

A higher phosphatase activity in savanna soil than in pasture soil may be due to very low P_i availability in savanna soil (Oberson et al., 2001), explained by differences in botanical composition of both systems (Rao et al., 1997). Measurements of phosphatase activity and microbial P on field experiment samples further indicate that M. carimaguensis participates in the mineralization of available P_o fractions. In the field experiments, phosphatase activity was greater in M. carimaguensis casts than in the control soil from the savanna. The conflicting observation seen in the laboratory experiment was probably an artifact of the methodology since the organic residues added to earthworm cultured pots could have already mineralized before ingestion by the earthworms. The high values of phosphatase activity obtained in our study for all treatments show the importance of both biological and biochemical processes in P_o mineralization (Oberson et al., 1995).

López Hernández et al. (1989) observed no difference in phosphatase activity between termite mounds and surrounding soil in Venezuelan savannas. However, Satchell and Martin (1984) reported high P_i content in fresh casts due to high phosphatase activity. A strong enzymatic activity has also been reported in fresh earthworm casts from temperate regions (Sharpley and Syers, 1976) as well as tropical sites (Mulongoy and Bedoret, 1989).

Microbial activity is enhanced in casts of tropical earthworms due to strong enzymatic activity and available organic C (Mulongoy and Bedoret, 1989). In this study values of microbial P were similar in both soil and cast samples from the lab experiment, and slightly higher in samples from the pasture. In summary, therefore, the increase in total P content of casts over soils in the field can be explained by organic matter, litter (including roots) and

cast selection by M. carimaguensis while the comparative increases in labile fractions extracted with water, NaHCO₃ and NaOH are due to the reorganization or translocation of P from stable to available pools for plant uptake. Evidence for the latter is found in the increased microbial biomass P in casts.

4.2. “Soil ecosystem engineers” and P dynamics in savanna soils

Casts of M. carimaguensis are large resistant structures that persist at the soil surface from two to eleven months on average in intensive pastures (exposed or protected to cattle trampling, respectively), and 5 months in native savannas (Decaëns, 2000). Some termite species (Nasutitermitinae) colonize these compact casts and create channels and deposit faecal pellets over its surface. When these casts finally split and smaller aggregates spread in the soil surface, the nutrients that were preserved from further mineralization processes may be released (Decaëns, 2000). Thus these surface casts represent a significant source of direct P easily available for plant uptake and may also explain the increase in root biomass observed under earthworm casts (Decaëns et al., 1999a). The role of other soil ecosystem engineers present in these savannas on P dynamics via the biogenic structures they create, for example, Microcerotermes sp., Spinitermes sp. (Termitidae) and Velocitermes sp. (Nasutitermitinae) species and the ants Atta laevigata and Acromyrmex landolti (Decaëns et al., 2001) should be addressed in further studies.

The importance of M. carimaguensis activity in natural and introduced pastures on incorporating P from organic sources into soil P pools, in increasing the labile P pools and improving P cycling was demonstrated in our short-term studies under laboratory and field conditions. The ecological significance of earthworms in P cycling in the native savanna and the introduced pasture is based on the improved nutritional basis of plant litter from the

pasture. There is an enhancement of biotic processes in the pasture, since populations of this species are quite large (Jiménez et al., 1998b) and there is also higher microbial biomass P (Oberson et al., 2001). The long-term effects of earthworms on P cycling and other nutrients must therefore be tremendous and merit further investigation, since the net benefits of earthworms in soil quality improvement and fertility are still ignored.

Studies on the role that soil ecosystem engineers play in P dynamics in Neotropical savannas are restricted to a few. Soil feeding termites can increase available P in their nests two or five-fold (Anderson and Wood, 1984). In the savannas of Venezuela, for example, both the quantity and distribution of P depends upon the biological activity of the termite Nasutitermes ephratae (Holmgren) (Nasutitermitidae) as reported by López-Hernández et al. (1989). The availability of mineral P derived from litter or soil can be increased by the activity of earthworms (Mansell et al., 1981), as we found in the field study. But P availability also changes with time as casts aged in the field. Similarly, Wood et al. (1983) found that P retention declined in the biogenic structures produced by the African savanna termite Cubitermes oculatus.

Acknowledgements

We thank the staff at Carimagua station for their friendship and invaluable help during field work, G. Borrero and M. Rodríguez (CIAT) for their technical assistance with the phosphorus fractionations. We are grateful to I. Rao (CIAT), P. Lavelle (IRD, France) and M. Brossard (IRD-EMBRAPA Brazil) for their comments, criticism and useful recommendations made on a previous version of this manuscript. Thanks to R. Thomas (CIAT) for English revision. Andrés F. Rangel is acknowledged for his help during fieldwork while doing his own research and A. Feijoo for his comments and support given to

one of the coauthors. We are also deeply appreciated to two anonymous referees for their suggestions that helped us improve this article.

References

Anderson, J.M., Wood, T.G., 1984. Mound composition and soil modification by two soil-feeding termites (Termitinae, Termitidae) in a riparian Nigerian forest. *Pedobiologia* 26, 77-82.

Barois, I., Lavelle, P., 1986. Changes in respiration rates and some physicochemical properties of a tropical soil during transit through Pontoscolex corethrurus (Glossoscolecidae, Oligochaeta). *Soil Biology & Biochemistry* 18, 539-541.

Barois, I., Lavelle, P., Brossard, M., Tondoh, J., Martínez, M.A., Rossi, J.P., Senapati, B.K., Angeles, A., Fragoso, C., Jiménez, J.J., Decaëns, T., Lattaud, C., Kanyonyo, J., Blanchart, E., Chapuis, L., Brown, G.G., Moreno, A., 1999. Ecology of earthworm species with large environmental tolerance and/or extended distributions. In: Lavelle, P., Brussaard, L., Hendrix, P.F. (Eds.), *Earthworm Management in Tropical Agroecosystems*. CAB-I, Wallingford, pp. 57-85.

Barois, I., Verdier, B., Kaiser, P., Lavelle, P., Rangel, P., 1987. Influence of the tropical earthworm Pontoscolex corethrurus (Glossoscolecidae) on the fixation and mineralization of nitrogen. In : Bovincini Pagliai, A. M., Omodeo, P. (Eds.), *On Earthworms*, Vol. 2. Selected Symposia and Monographs, U.Z.I., Ed. Mucchi, Modena, pp. 151-158.

Bohlen, P.J., Parmelee, R.W., McCartney, D.A., Edwards, C.A., 1997. Earthworm effects on Carbon and Nitrogen dynamics of surface litter in corn agroecosystems. *Ecological Applications* 7(4), 1341-1349.

Bouché, M.B., 1977. Stratégies lombriciennes. In: Lohm, U., Persson, T. (Eds.), *Soil Organisms as Component of Ecosystems*. *Ecol. Bull. (Stockholm)* 25, 122-132.

Bowman R.A., 1988. A rapid method to determine total phosphorous in soils. *Soil Science Society of America Journal* 52, 1301-1304.

- 1 Bowman R.A., Cole, C., 1978. An exploratory method for fractionation of organic
2 phosphorous from grassland soils. *Soil Science* 125, 95-100.
- 3 Brossard, M., Lavelle, P., Laurent, J.Y., 1996. Digestion of a vertisol by the endogeic
4 earthworm (*Polypheretima elongata*, *Megascolecidae*) increases soil phosphate extractability.
5 *European Journal of Soil Biology* 32, 107-111.
- 6 Buse, A., 1990. Influence of earthworms on nitrogen fluxes and plant growth in cores taken
7 from variously managed upland pastures. *Soil Biology & Biochemistry* 22 (6), 775-780.
- 8 Chapuis, L., Brossard, M., 1995. Modifications et stabilité du phosphore échangeable d'un
9 ferrasol ingère par un ver géophage. *Compte Rendus de l'Académie des Sciences de Paris*
10 *Series Ila* 320, 587-592.
- 11 Coleman, D.C., Hendrix, P.F., Beare, M.H., Crossley, D.A., Hu, S., van Vliet, P.C.J., 1994.
12 The impacts of management and biota on nutrient dynamics and soil structure in sub-tropical
13 agroecosystems: Impacts on detritus food webs. In: Pankhurst, C.E., Doube, B.M., Gupta,
14 V.V.S.R., Grace, P.R. (Eds.), *Soil Biota. Management in Sustainable Farming Systems*.
15 CSIRO, Melbourne, pp. 133-143.
- 16 Cooper, D.W., 1968. The significance level in multiple tests made simultaneously. *Heredity* 23,
17 614-617.
- 18 Decaëns, T., 2000. Degradation dynamics of surface earthworm casts in grasslands of the
19 eastern plains of Colombia. *Biology and Fertility of Soils* 32, 149-156.
- 20 Decaëns, T., Galvis, J.H., Amézquita, E., 2001. Propriétés des structures produites par les
21 ingénieurs écologiques à la surface du sol d'une savane colombienne. *Compte Rendus de*
22 *l'Académie des Sciences de Paris Série III* 324(5), pp. 465-478.
- 23 Decaëns, T., Lavelle, P., Jiménez, J.J., Escobar, G., Rippstein, G., 1994., Impact of land
24 management on soil macrofauna in the Oriental Llanos of Colombia. *European Journal of*
25 *Soil Biology* 30(4), 157-168.

- 1 Decaëns, T., Mariani, L., Lavelle, P., 1999a. Soil surface macrofaunal communities
2 associated with earthworm casts in grasslands of the Eastern Plains of Colombia. *Applied*
3 *Soil Ecology* 13, 87-100.
- 4 Decaëns, T., Rangel, A.F., Asakawa, N., Thomas, R.J., Lavelle, P., 1999b. Carbon and
5 nitrogen dynamics in ageing earthworm casts in grasslands of the Eastern plains of
6 Colombia. *Biology & Fertility of Soils* 30, 20-28.
- 7 Fisher, M.J., Rao, I.M., Ayarza, M.A., Lascano, C.E., Sanz, J.I. , Thomas, R.J. and Vera,
8 R.R., 1994. Carbon storage by deep-rooted grasses in the Southamerican savannas. *Nature*
9 371, 236-238.
- 10 Friesen, D.K., Rao, I.M., Thomas, R.J., Oberson, A., Sanz, J.I., 1997. Phosphorous
11 acquisition and cycling in crop and pasture systems in low fertility tropical soils. *Plant &*
12 *Soil* 196, 289-294.
- 13 Frossard, E., Brossard, M., Hedley, M. J., Metherel, A., 1995. Reactions controlling the
14 cycling of P in soils. In: Tiessen, H. (Ed.), *Phosphorus in the Global Environment*. John
15 Wiley and Sons, New York.
- 16 Guggenberger, G., Haumaier, L., Thomas, R.J., Zech, W., 1996. Assessing the organic
17 phosphorous status of an Oxisol under tropical pastures following native savanna using ^{31}P
18 NMR spectroscopy. *Biology & Fertility of Soils* 23, 332-339.
- 19 Hedley, M.J., Stewart, J.W.B., Chauhan, B.S., 1982. Changes in inorganic and organic soil
20 phosphorous fractions induced by cultivation practices and by laboratory incubations. *Soil*
21 *Science Society of America Journal* 46, 970-976.
- 22 James, S.W. 1991. Soil, nitrogen, phosphorous and organic matter processing by earthworms
23 in tallgrass prairie. *Ecology* 72(6), 2101-2109.
- 24 Jiménez, J.J., 1999. Estructura de las Comunidades y Dinámica de las Poblaciones de
25 Lombrices de Tierra en las Sabanas Naturales y Perturbadas de Carimagua (Colombia).
26 Doctoral Thesis, Universidad Complutense, Madrid, 311 p (in Spanish).

- 1 Jiménez, J.J., Decaëns, T., 2000. Vertical distribution of earthworms in grassland soils of the
2 Colombian Llanos. *Biology & Fertility of Soils* 32, 463-473.
- 3 Jiménez, J.J., Moreno, A.G., Lavelle, P., 1999. Reproductive strategies of three native
4 earthworm species from the savannas of Carimagua (Colombia). *Pedobiologia* 43, 851-858.
- 5 Jiménez, J.J., Moreno, A.G., Lavelle, P., Decaëns, T., 1998a. Population dynamics and
6 adaptive strategies of Martiodrilus carimaguensis (Oligochaeta, Glossoscolecidae), a native
7 species from the well-drained savannas of Colombia. *Applied Soil Ecology* 9 (1-3), 153-160.
- 8 Jiménez, J.J., Brown, G.G., Decaëns, T., Feijoo, A., Lavelle, P., 2000. Differences in the timing
9 of diapause and patterns of aestivation in some tropical earthworms. *Pedobiologia* 44(6), 677-
10 694.
- 11 Jiménez, J.J., Moreno, A.G., Decaëns, T., Lavelle, P., Fisher, M.J., Thomas, R.J., 1998b.
12 Earthworm communities in native savanna and man-made pastures of the eastern plains of
13 Colombia. *Biology & Fertility of Soils* 28, 101-110.
- 14 Jones, C.G., Lawton, J.H., Shachak, M., 1994. Organisms as ecosystem engineers. *Oikos* 69,
15 373-386.
- 16 Krishnamoorthy, R.V., 1990. Mineralization of phosphorous by faecal phosphatases of some
17 earthworms of Indian tropics. *Proceedings of the Indian Academie of Sciences (Animal*
18 *Sciences)* 95, 341-351.
- 19 Krom, M., 1980. Spectrophotometric determination of ammonia, a study of modified
20 Bertheloy reaction using salicylate and dichloroisocyanurate. *Analyst* 105, 305-316.
- 21 Lal, R., 1974. No tillage effects on soil properties and maize (Zea mays L.) production in
22 western Nigeria. *Plant & Soil*, 40: 321-331.
- 23 Lal, R., 1991. Soil conservation and biodiversity. In: Hawksworth, D.L. (Ed.), *The*
24 *Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture*. CAB
25 International, Wallingford, pp. 89-103.

- 1 Lavelle, P., 1997. Faunal activities and soil processes: adaptive strategies that determine
2 ecosystem function. *Advances in Ecological Research* 27, 93-132
- 3 Lavelle, P., 2000. Ecological challenges for soil science. *Soil Science* 165, 73-86.
- 4 Lavelle, P., Martin, A., 1992. Small-scale and large-scale effects of endogeic earthworms on
5 soil organic matter dynamics in soils of the humid tropics. *Soil Biology & Biochemistry*
6 24(12), 1491-1498.
- 7 Lavelle, P., Blanchart, E., Martin, A., Spain, A.V., Martin, S., 1992. Impact of soil fauna on the
8 properties of soils in the humid tropics. In: Segoe, S. (Ed.), *Myths and Science of Soils of the*
9 *Tropics*. SSSA Special Publication No. 29, Madison, pp. 157-185.
- 10 Lilliefors, H.W., 1967. The Kolmogorov-Smirnov test for normality with mean and variance
11 unknown. *Journal of American Statistical Association* 62, 399-402.
- 12 López-Hernández, D., Lavelle, P., Fardeau, J.C., Niño, M., 1993. Phosphorous
13 transformations in two P-sorption contrasting tropical soils during transit through
14 Pontoscolex corethrurus (Glossoscolecidae: Oligochaeta). *Soil Biology & Biochemistry* 25
15 (6), 789-792.
- 16 López-Hernández, D., Fardeau, J.C., Niño, M., Nannipieri, P., Chacón, P., 1989.
17 Phosphorous accumulation in savanna termite mound in Venezuela. *Journal of Soil Science*
18 40, 635-640.
- 19 Lunt, H.A., Jacobson, G.M., 1944. The chemical composition of earthworm casts. *Soil*
20 *Science* 58, 367-375.
- 21 Mansell, G.P., Syers, J.K., Gregg P.E.H., 1981. Plant availability of phosphorous in dead
22 herbage ingested by surface-casting earthworms. *Soil Biology & Biochemistry* 13, 163-167.
- 23 Mariani, L., 2001. Impact des biostructures produites par Martiodrilus carimaguensis
24 (Oligochaeta, Glossoscolecidae) sur le fonctionnement du sol dans les savannes orientales
25 de Colombie. Thèse de Doctorat. Université Pierre et Marie Curie, Paris VI. 236p. (in
26 French).

- 1 Mariani, L., Bernier, N., Jiménez, J.J. and Decaëns, T., 2001. Régime alimentaire d'un ver
2 de terre des savanes colombiennes – une remise en question des types écologiques. *Compte*
3 *Rendus de l'Académie des Sciences de Paris Série III*, 324(8): 733-742.
- 4 Mariani, L., Decaëns, T., Jiménez, J.J., Torres, E.A., Amézquita, E., Rainfall impact on casts
5 of various ages of a tropical anecic earthworm. *Geoderma* (in press).
- 6 Marinissen, J.C.Y., de Ruiter P.C., 1993. Contribution of earthworms to carbon and nitrogen
7 cycling in agroecosystems. *Agriculture Ecosystems and Environment* 47, 59-74.
- 8 Martin, A., Marinissen, J.Y.C., 1993. Biological and physico-chemical processes in
9 excrements of soil animals. *Geoderma* 56, 331-347.
- 10 Mattingly, G.E.G., 1975. Labile phosphate in soil. *Soil Science* 119, 369-375.
- 11 Mullen, M.D., 1998. Transformations of other elements. In: Sylvia, D.M., Fuhrmann, J.J.,
12 Hartel, P.G., Zuberer, D.A. (Eds.), *Principles and Applications of Soil Microbiology*.
13 Prentice Hall, New Jersey, pp. 369-386.
- 14 Mulongoy, K., Bedoret, A., 1989. Properties of worm casts and surface soils under various
15 plant covers in the humid tropics. *Soil Biology & Biochemistry* 21, 197-203.
- 16 Murphy, J., Riley, J.P., 1962. A modified single solution method for the determination of
17 phosphate in natural waters. *Anal Chim Acta* 27, 31-36.
- 18 Nye, P.H., 1955. Some soil-forming processes in the humid tropics. IV. The action of soil
19 fauna. *Journal of Soil Science* 6(1), 73-83.
- 20 Oberson, A., Friesen, D.K., Rao, I.M., Bühler, S., Frossard, E., 2001. Phosphorous
21 transformations in an oxisol under contrasting land-use systems: the role of the soil
22 microbial biomass. *Plant & Soil* 237, 197-210.
- 23 Oberson, A., Friesen, D.K., Tiessen, H., Moir, J.O. and Borrero, G., 1995. Phosphorus
24 transformations in improved pastures, In: *Tropical Lowlands Program Annual Report 1994*.
25 Working document no. 148. CIAT, Cali, Colombia, pp. 182-187.

- 1 Oberson, A., Friesen, D.K., Tiessen, H., Morel, C., Stahel, W., 1999. Phosphorus status and
2 cycling in native savanna and improved pastures on an acid low-P Colombian Oxisol.
3 *Nutrient Cycling in Agroecosystems* 55(1), 77-88.
- 4 Olsen, S.R., Sommers, L.E., 1982. Phosphorous. In: Page, A.L., Miller, R.H., Keeney, D.R.
5 (Eds.), *Methods of Soil Analysis*, part 2, 2nd ed. ASA, Madison, Wisconsin, pp. 404-430.
- 6 Satchell, J.E., Martin, K., 1984. Phosphatase activity in earthworm faeces. *Soil Biology &*
7 *Biochemistry* 16, 191-194.
- 8 Rao, I.M., Borrero, V., Ricaurte, J., García, R., Ayarza, M.A., 1997. Adaptive attributes of
9 tropical forage species to acid soils III. Differences in phosphorus acquisition and utilization
10 as influenced by varying phosphorus supply and soil type. *Journal of Plant Nutrition* 20, 155-
11 180.
- 12 Scheu, S., 1987. Microbial activity and nutrient dynamics in earthworm casts (Lumbricidae).
13 *Biology & Fertility of Soils* 5, 230-234.
- 14 Sharpley, A.N., Syers, J.K., 1976. Potential role of earthworm casts for the Phosphorous
15 enrichment of run-off waters. *Soil Biology & Biochemistry* 8, 341-346.
- 16 Sparling, G.P., Whale, K.N., Ramsay, A.J., 1985. Quantifying the contribution from the soil
17 microbial biomass to the extractable P levels of fresh and acid air-dried soils. *Australian*
18 *Journal of Soil Research* 23, 613-621.
- 19 Stork, N.E., Eggleton, P., 1992. Invertebrates as determinants and indicators of soil quality.
20 *American Journal of Alternative Agriculture* 7, 38-47.
- 21 Swift, M.J., Heal, O.W., Anderson, J.M., 1979. *Decomposition in Terrestrial Ecosystems*.
22 Blackwell Scientific, Oxford.
- 23 Tabatabai, M.A., 1982. Soil enzymes. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.),
24 *Methods of Soil Analysis*, part 2. Chemical and Microbiological properties. ASA and SSSA
25 publication, Madison, Wisconsin, pp. 903-947.

- 1 Thomas, R.J., Lascano, C., Sanz, J.I., Ara, M.A., Spain, J.M., Vera, R.R., Fisher, M.J., 1992.
2 The role of pastures in production systems. In: CIAT. Pastures for the Tropical Lowlands.
3 CIAT's Contributions. Cali, Colombia, pp. 121-144.
- 4 Thomas, R.J., 1992. The role of the legume in the nitrogen cycle of productive and
5 sustainable pastures. Grass and Forage Science 47, 133-142.
- 6 Thomas, R.J., 1995. Role of legumes in providing N for sustainable tropical pasture systems.
7 Plant & Soil 174, 103-118.
- 8 Thomas, R.J., Fisher, M.J., Ayarza, M.A., Sanz, J.I., 1995. The role of forage grasses and
9 legumes in maintaining the productivity of acid soils in Latin America. In: Lal, R., Stewart,
10 B.A. (Eds.), Soil Management: Experimental Basis for Sustainability and Environmental
11 Quality. Advances in Soil Science series. Lewis Publishers, Boca Raton, pp. 61-83.
- 12 Thompson, L., Thomas, C.D., Radley, J.M.A., Williamson, S., Lawton, J.H., 1993. The
13 effect of earthworms and snails in a simple plant community. Oecologia 95, 171-178.
- 14 Tiessen, H., Moir, J.O., 1993. Characterization of available P by sequential extraction. In:
15 Carter, M.R. (Ed.), Soil Sampling and Methods of Analysis. Lewis Publishers, Boca Raton,
16 FL, USA. pp. 75-86.
- 17 Wood, T.G., Johnson, R.A., Anderson, J.M., 1983. Modification of soils in Nigerian
18 savanna by soil feeding Cubitermes (Isoptera, Termitidae). Soil Biology & Biochemistry 15,
19 575-579.

20

Tables

Table 1. Physico-chemical properties of soils in the native savanna and the intensive pasture[†]

Land use system	pH(H ₂ O) 1:1	Total C	Total N	Bray-II P	Exchangeable cations				
					Al	Ca	Mg	K	H
		----- mg g ⁻¹ -----			----- cmol ⁺ kg ⁻¹ -----				
Savanna	4.80	23.5	1.45	1.3	2.42	0.26	0.11	0.08	0.27
Pasture	4.96	24.9	1.67	2.2	1.90	0.89	0.23	0.09	0.26

[†] Oberson et al. (1999)

1 Table 2. Chemical properties (mean $\pm$ standard error) of soil and one day-old casts of *M. carimaguensis* in the
2 laboratory (LE1 experiment) and field experiments.

Experiment	pH (H ₂ O)	Total C (%)	Bray-II P (mg-P kg ⁻¹)	Phosphatase activity (mg kg ⁻¹ h ⁻¹) [§]	Microbial P (mg-P kg ⁻¹)	Total P [‡] (HClO ₄ dig.) (mg kg ⁻¹)
<i>Laboratory Experiment</i>						
Native savanna						
Soil	4.6 $\pm$ 0.01	2.6 $\pm$ 0.04	2.6 $\pm$ 0.19	215 $\pm$ 13.5	4.1 $\pm$ 0.07	208 $\pm$ 0.88
Casts	5.2 $\pm$ 0.02	2.5 $\pm$ 0.00	2.9 $\pm$ 0.13	120 $\pm$ 10.1	4.1 $\pm$ 0.44	225 $\pm$ 4.06
<i>B. decumbens</i> – Kudzu pasture						
Soil	4.6 $\pm$ 0.01	2.9 $\pm$ 0.02	4.2 $\pm$ 0.20	313 $\pm$ 16.6	6.0 $\pm$ 0.52	248 $\pm$ 0.33
Casts	5.2 $\pm$ 0.05	3.0 $\pm$ 0.04	4.1 $\pm$ 0.65	242 $\pm$ 6.63	5.4 $\pm$ 0.24	272 $\pm$ 0.67
<i>Field Experiment</i>						
Native savanna						
Soil	5.1 $\pm$ 0.04	2.1 $\pm$ 0.02	1.0 $\pm$ 0.06	254 $\pm$ 7.46	2.5 $\pm$ 0.13	179 $\pm$ 1.42
Casts	5.4 $\pm$ 0.04	4.1 $\pm$ 0.03	6.3 $\pm$ 0.25	312 $\pm$ 4.2	4.0 $\pm$ 0.34	267 $\pm$ 5.92
<i>B. decumbens</i> – Kudzu pasture						
Soil	5.2 $\pm$ 0.05	2.1 $\pm$ 0.04	2.5 $\pm$ 0.20	299 $\pm$ 6.89	4.1 $\pm$ 0.67	194 $\pm$ 2.65
Casts	5.8 $\pm$ 0.65	5.2 $\pm$ 0.11	11.0 $\pm$ 0.92	249 $\pm$ 7.13	10.9 $\pm$ 0.88	396 $\pm$ 28.06

3 [§] mg p-nitrophenol kg⁻¹ h⁻¹

4 [‡] from laboratory experiment 2 (LE2)

5

1 Table 3. Two-way analyses of variance (ANOVA) for phosphatase activity, microbial biomass P, pH, Bray-II P, Total P and Total C in soil and one day-old casts of M.
2 carimaguensis from incubation in the field. The F-ratios and error mean squares for each variable are indicated. Each test is significant at the Bonferroni corrected probability
3 level [overall probability / (n of variable x n of tests)] for overall significant levels of 0.05, 0.01, and 0.001.^a
4

Source	<i>df</i>	Phosphatase activity ^b	Microbial P ^b	pH ^b	Bray-II P ^c	Total P ^c	Total C ^c
Land use (A)	1	0.02 NS	44.10***	23.30**	94.58***	40.43***	62.65***
Cast vs. soil (B)	1	0.17 NS	48.47***	100.64***	551.91***	272.90***	2595.10***
AxB	1	54.72***	9.14 NS	5.14 NS	0.28 NS	20.25***	71.57***
Error mean squares	10 ^b , 24 ^c	0.00044	0.0048	0.00004	0.0037	0.0014	0.00018

^a NS = not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Table 4. Phosphorus fractions in soil and fresh (1-day-old) casts of *M. carimaguensis* collected in the field from the native savanna and the intensive pasture¹.

		H ₂ O-Po	Resin-Pi	NaHCO ₃		NaOH		1M HCl-Pi	HCl hc		Residue -Pt	Total Pi	Total Po	Total P
				Pi	Po	Pi	Po		Pi	Po				
----- mg P kg ⁻¹ soil -----														
Savanna	Soil	0.5 b	0.8 b	1.6 c	8.6 b	22 bc	42 ac	0.3 b	38 c	22 ac	59 b	63	73 b	195 c
	Cast	1.4 a	7.8 a	9.9 b	17.0 a	52 ac	68 a	0.9 b	52 b	21 ac	68 b	123	108 a	299 b
	%increase	180	875	519	98	137	64	200	36	-5	15	95	47	53
	%P added ²	1	7	8	8	29	25	1	13	-1	9	58	33	100
<i>B. decumbens</i> + Kudzu	Soil	0.8 b	1.4 b	2.8 c	9.1 b	26 b	43 bc	0.8 b	45 bc	10 bc	60 b	76	62 b	199 c
	Cast	2.0 a	12.8 a	19.0 a	18.8 a	82 a	82 a	5.6 a	83 a	32 a	88 a	202	136 a	425 a
	%increase	150	814	579	107	213	92	600	83	234	46	165	117	114
	%P added	1	5	7	4	25	17	2	17	10	12	56	32	100

¹ values within a column followed by the same letter do not differ significantly ($P < 0.05$) according to Tukey's HSD test.

² percentage of total P increase in cast over soil found in respective fraction.

Table 5. Phosphorus fractions in soil and fresh (1-day-old) casts of *M. carimaguenensis* produced in the laboratory (LE1) from soil collected in the native savanna and the intensive pasture¹.

		H ₂ O-Po	Resin-Pi	NaHCO ₃		NaOH		1M HCl-Pi	HCl hc		Residue -Pt	Total Pi	Total Po	Total P
				Pi	Po	Pi	Po		Pi	Po				
----- mg P kg ⁻¹ soil -----														
Savanna	Soil	1.6 ab	0.9 d	4.1 b	1.8	25 d	32 c	0.3	41 b	16	43	71	51 bc	165 b
	Cast	1.9 b	4.0 b	4.3 b	2.4	30 c	50 b	0.4	42 b	17	42	81	71 ac	193 bc
	%increase	19	344	5	33	23	57	33	2	6	-3	14	39	17
	%P added ²	1	11	1	2	20	64	0	2	3	-5	35	70	100
<i>B. decumbens</i> + Kudzu	Soil	1.7 ab	2.8 c	6.5 a	1.0	34 b	54 a	0.2	51 a	22	46	94	78 a	218 ac
	Cast	2.5 a	6.3 a	6.5 a	1.4	39 a	57 a	0.7	55 a	20	55	108	80 a	243 a
	%increase	47	125	0	40	14	5	250	9	-12	20	14	2	11
	%P added	3	15	0	2	20	11	2	20	-11	38	57	5	100

¹ values within a column followed by the same letter (or no letter) do not differ significantly ($P < 0.05$) according to Tukey's HSD test.

² percentage of total P increase in cast over soil found in respective fraction.

Table 6. Two-way ANOVA for the different P fractions analysed in control soil and incubated casts of *M. carimaguensis* in the field. The F-ratios and error mean squares for each variable are indicated. Each test is significant at the Bonferroni corrected probability level [overall probability / (n of variable x n of tests)] for overall significant levels of 0.05, 0.01, and 0.001.^a

Source	df	H ₂ O-P _o	Resin-P _i	NaHCO ₃ -P _o	NaHCO ₃ -P _i	NaOH-P _o	NaOH-P _i
Land use (A)	1	58.65 ***	87.49 ***	32.67 ***	118.81 ***	37.15 ***	229.31 ***
Cast age (B)	6	17.54 ***	21.35 ***	15.23 ***	32.04 ***	25.81 ***	49.98 ***
AxB	6	3.50 NS	2.89 NS	2.25 NS	3.67 *	1.42 NS	4.30 *
Error mean squares	66	0.11	4.02	7.70	7.99	81.02	47.97

^a NS = not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Table 7. Two-way ANOVA for the different P fractions analysed in control soil and incubated casts of *M. carimaguensis* in the laboratory. The F-ratios and error mean squares for each variable are indicated. Each test is significant at the Bonferroni corrected probability level [overall probability / (n of variable x n of tests)] for overall significant levels of 0.05, 0.01, and 0.001.^a

Source	df	H ₂ O-P _o	Resin-P _i	NaHCO ₃ -P _o	NaHCO ₃ -P _i	NaOH-P _o	NaOH-P _i
Land use (A)	1	9.25 NS	103.77 ***	0.18 NS	13.67 *	2.65 NS	57.72 ***
Cast vs. soil (B)	6	11.50 ***	22.66 ***	9.83 ***	1.15 NS	1.40 NS	2.40 NS
AxB	6	0.71 NS	2.13 NS	0.65 NS	1.63 NS	1.22 NS	0.93 NS
Error mean squares	66	0.15	0.27	15.98	1.72	176.32	18.99

^a NS = not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Figure captions

Figure 1. Dynamics of P_i (●) and P_o (Δ) fractions in casts of M. carimaguensis produced and incubated in the laboratory (LE1 experiment). *C* Control (non-ingested) soil. Bars indicate standard deviation.

Figure 2. Dynamics of P_i (●) and P_o (Δ) fractions in *in-situ* ageing casts of M. carimaguensis. *C* Control (non-ingested) soil. Bars indicate standard deviation.

Figure 1

1
2

3 Figure 2